

Synthetic Methods

Palladium-Catalyzed Synthesis of Ammonium Sulfinates from Aryl Halides and a Sulfur Dioxide Surrogate: A Gas- and Reductant-Free Process**



Edward J. Emmett, Barry R. Hayter, and Michael C. Willis*

Abstract: Sulfonyl-derived functional groups populate a broad range of useful molecules and materials, and despite a variety of preparative methods being available, processes which introduce the most basic sulfonyl building block, sulfur dioxide, using catalytic methods, are rare. Described herein is a simple reaction system consisting of the sulfur dioxide surrogate DABSO, triethylamine, and a palladium(0) catalyst for effective conversion of a broad range of aryl and heteroaryl halides into the corresponding ammonium sulfinates. Key features of this gas- and reductant-free reaction include the low loadings of palladium (1 mol%) and ligand (1.5 mol%) which can be employed, and the use of isopropyl alcohol as both a solvent and formal reductant. The ammonium sulfinate products are converted in situ into a variety of sulfonyl-containing functional groups, including sulfones, sulfonyl chlorides, and sulfonamides.


Sulfonyl-derived functional groups such as sulfones, sulfonamides, and sulfonic acids feature in a myriad of applications ranging from active ingredients in pharmaceuticals^[1] and agrochemicals,^[2] to key intermediates in complex chemical syntheses,^[3] to speciality materials^[4] and food additives.^[5] The syntheses of these functional groups are in general based on classic reactivity patterns. For example, sulfones are most commonly prepared by the oxidation of a precursor sulfide, itself usually obtained from a conventional nucleophile/electrophile combination.^[6] Aromatic sulfonamides are ordinarily accessed from the corresponding sulfonyl chlorides, which in turn are most often available from the parent aromatic using electrophilic aromatic substitution chemistry.^[7] Both of these approaches feature significant limitations: the use of an oxidative route precludes the presence of

oxidation-sensitive functional groups, and the corresponding sulfide precursors often require the use of foul-smelling thiols for their preparation. Using electrophilic aromatic substitution chemistry limits the accessible molecules to those that conform to the inherent electronic and steric factors which control selectivity in these processes. Alternative procedures exist for both of these functional groups;^[8] however, convenient and general methods to access these important molecules are still lacking.

The fundamental feature that links all sulfonyl-derived functional groups is the -SO₂- arrangement of atoms, thus suggesting that a conceptually simple route to these molecules is one that involves the direct catalytic introduction of SO₂. Despite the ready availability of SO₂ as a bulk feedstock, its use in transition-metal catalysis is severely limited. Pioneering reports from Pelzer and Keim established that the palladium-catalyzed introduction of SO₂ was feasible, although these early reports of sulfinic acid synthesis described reactions of poor scope and required high pressures of SO₂ and hydrogen, and the use of inconvenient aryl diazonium substrates [Eq. (1)].^[9]



This work: direct ammonium sulfinate synthesis:



We have introduced the amine-SO₂ charge-transfer complex DABSO [DABCO·(SO₂)₂] as a convenient and easy-to-handle solid which functions as a surrogate for SO₂ gas in a number of known transformations,^[10,11] and also in a new palladium-catalyzed aminosulfonylation procedure using aryl halides.^[12] Wu et al. have reported related transformations employing K₂S₂O₅ as the SO₂ source,^[13] and also the use of aryl boronic acids to access identical products.^[14] All of these processes suffer from the limitation that only hydrazine nucleophiles can be employed.^[15]

[*] E. J. Emmett, Prof. M. C. Willis
Department of Chemistry, University of Oxford
Chemistry Research Laboratory
Mansfield Road, Oxford, OX1 3TA (UK)
E-mail: michael.willis@chem.ox.ac.uk
Homepage: <http://mcwillis.chem.ox.ac.uk/MCW/Home.html>

Dr. B. R. Hayter
Oncology Innovative Medicines, AstraZeneca
Alderley Park, Macclesfield, Cheshire, SK10 4TG (UK)

[**] This work was supported by the EPSRC and AstraZeneca.

Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/anie.201404527>.

© 2014 The Authors. Published by Wiley-VCH Verlag GmbH & Co. KGaA. This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

Rather than focus on the synthesis of a single class of sulfonyl derivative, a catalytic synthesis of sulfonates would potentially allow access to a host of sulfonyl-derived functional groups through established reactions.^[16] Although we have recently shown that aminosulfonamides, prepared from the palladium-catalyzed combination of aryl halides, DABSO, and hydrazines, can be converted into the corresponding sulfonates by way of a base-promoted degradation, the process is cumbersome in that it requires the initial introduction and then expulsion of an expensive hydrazine derivative [Eq. (2)].^[17] Shavnya and Miscitti have also shown that sulfonates can be prepared using palladium catalysis, in this case, from the combination of aryl halides and $K_2S_2O_5$. However, the process requires high ligand loadings (30 mol %), the use of an excess of inorganic reductant, as well as a stoichiometric additive (TBAB).^[18] Herein we describe an operationally simple protocol which employs only DABSO, triethylamine, and isopropyl alcohol, in combination with a low palladium and ligand loading to convert aryl halides into the corresponding ammonium sulfonates [Eq. (3)]. No inorganic reducing agent is needed, with isopropyl alcohol serving as both the reaction solvent and the formal reductant. The process is high yielding across a broad range of aryl and heteroaryl halides, and the ammonium sulfonate products display good reactivity in their conversion into a number of sulfonyl derivatives.

We selected the conversion of 4-iodotoluene (**1a**) into the corresponding sulfonate (**2a**) as a model transformation (Table 1). Our initial reagent and catalyst selections were based on our earlier developed aminosulfonylation chemistry, with the exchange of the hydrazine nucleophile for a range of reductants. The desired sulfonate was only observed in significant quantities when sodium formate in isopropyl

alcohol was employed (entry 1).^[19] The use of calcium formate provided a small improvement in efficiency (entries 2 and 3), and these conditions were then used to evaluate a number of alternative phosphine ligands (entries 4–9). Significant improvements in yields were obtained with both PCy_3 and $PrBu_2Me$ (used as their HBf_4 salts), and the optimal ligand proved to be $PdAd_2Bu$ ^[20] (entry 9), which in addition to providing a small improvement in yield also delivered a practical advantage because of its improved stability.^[20] We were aware that the *i*PrOH, used as solvent, could also function as a reductant in palladium catalysis,^[21] and accordingly we then performed a duplicate experiment but without the addition of any external inorganic reducing agent. Pleasingly, an identical yield was obtained for the reductant-free transformation (entry 10). As far as we are aware, these experiments represent the first example of an alcohol being employed as a reductant in combination with a sulfinyl derivative. Finally, decreasing the palladium loading to 5 mol %, the ligand loading to 7.5 mol %, and the number of DABSO equivalents to 0.6 (1.2 equivalents of SO_2) delivered the optimized reaction conditions (entry 11). The catalyst loading could be reduced further, for example, using only 0.5 mol % palladium and 0.75 % ligand led to a 91 % conversion into the sulfonate (entry 12).

Sulfonates are versatile intermediates, but they are rarely the desired final product of a transformation. To evaluate the scope of the developed reaction we elected to derivatize the newly formed sulfonates in situ. Using α -bromo *tert*-butyl acetate as the electrophile allowed the corresponding sulfones to be readily prepared, and was preferable to attempting to isolate and purify individual ammonium sulfonate salts (Table 2). In general, a broad range of functionalized aryl iodides could be converted smoothly into the corresponding sulfonates, and then into the isolable sulfone products. Electron-rich substrates performed well, with a variety of electron-donating substituents being tolerated in all positions of the aromatic ring (**3a–f**). Notable examples include the mixed sulfide-sulfone **3d**, a product difficult to prepare using traditional sulfone methodology, and sulfones **3e** and **3f**, featuring unprotected OH and NH_2 substituents, respectively. A selection of electron-neutral substrates were employed without incident (**3g–j**). Electron-withdrawing functional groups were likewise compatible with the reaction conditions, with ester, ketone, amide, nitrile, and trifluoromethyl-substituted products being isolated in good yields (**3k–o**). The greater reactivity of aryl iodide substrates allowed sulfones featuring bromo, chloro, and fluoro substituents to be prepared (**3p–r**). Finally, although aryl iodide substrates proved to be most efficient, it was possible to employ aryl bromide starting materials, with the sulfones **3k** and **3o** being prepared from both aryl halide substrates.


We next explored the use of the heteroaryl halides as substrates (Table 3). Examples of indole, benzofuran, benzothiophene, thiophene, pyridine, and quinolone ring systems were all converted into the expected sulfone products, although in general the yields were reduced relative to the benzene-derived examples (**3s–x**). Several alkenyl iodides were also investigated, and proved to be compatible with the reaction (**3y,z**).

Table 1: Optimization of reaction conditions for the formation of the sulfonate **2a** from 4-iodotoluene.^[a]

Entry	Ligand	Reductant	Yield [%] ^[b]
1	$PrBu_3 \cdot HBF_4$	HCO_2Na	68
2	$PrBu_3 \cdot HBF_4$	HCO_2K	38
3	$PrBu_3 \cdot HBF_4$	$(HCO_2)_2Ca$	71
4	binap	$(HCO_2)_2Ca$	23
5	dppf	$(HCO_2)_2Ca$	84
6	$dccp \cdot (HBF_4)_2$	$(HCO_2)_2Ca$	25
7	$PCy_3 \cdot HBF_4$	$(HCO_2)_2Ca$	88
8	$PrBu_2Me \cdot HBF_4$	$(HCO_2)_2Ca$	95
9	$PA_{d_2}Bu$	$(HCO_2)_2Ca$	96
10	$PA_{d_2}Bu$	none	96
11 ^[c]	$PA_{d_2}Bu$	none	99
12 ^[d]	$PA_{d_2}Bu$	none	91


[a] Reaction conditions: $Pd(OAc)_2$ (10 mol %), ligand (20 mol %), DABSO (1.1 equiv), Et_3N (3.0 equiv), *i*PrOH [0.2 M], 75 °C, 16 h. [b] HPLC yield relative to an internal standard. [c] $Pd(OAc)_2$ (5 mol %), $PA_{d_2}Bu$ (7.5 mol %), DABSO (0.6 equiv). [d] $Pd(OAc)_2$ (0.5 mol %), $PA_{d_2}Bu$ (0.75 mol %). binap = 2,2'-bis(diphenylphosphanyl)-1,1'-binaphthyl, dccp = 1,3-bis(dicyclohexylphosphino)propane, dppf = 1,1'-bis(diphenylphosphino)ferrocene.

Table 2: Scope of aryl halides in the palladium-catalyzed preparation of aryl ammonium sulfonates.^[a]




[a] Reaction conditions: Pd(OAc)₂ (5 mol%), PAd₂Bu (7.5 mol%), DABSO (0.6 equiv), Et₃N (3.0 equiv), iPrOH [0.2 M], 75 °C, 16 h. DMF = N,N-dimethylformamide.

Table 3: Heteroaryl and alkenyl halides in the palladium-catalyzed preparation of ammonium sulfonates.^[a]



[a] Reaction conditions: Pd(OAc)₂ (5 mol%), PAd₂Bu (7.5 mol%), DABSO (0.6 equiv), Et₃N (3.0 equiv), iPrOH [0.2 M], 75 °C, 16 h. [b] Pd(OAc)₂ (10 mol%), PAd₂Bu (15 mol%), DABSO (1.1 equiv).


For pragmatic reasons all of the reactions shown in Tables 2 and 3 were performed using 5 mol% of palladium and 7.5 mol% of ligand. On a preparative scale (12 mmol) it was possible to achieve good reactivity at lower loadings, with the sulfone **3g** being isolated in 96% yield from a reaction employing 1 mol% of palladium and 1.5 mol% of ligand (Scheme 1). Although we have not scaled the reaction further, the use of isopropyl alcohol as both the solvent and formal reductant holds promise in this perspective, as it has been identified as a sustainable solvent suitable for process development.^[22]



Scheme 1. Preparative-scale synthesis of the sulfone **3g**.

At the outset of our investigation an efficient catalytic synthesis of sulfonates was targeted because of the wide range of established reactivity, for additional transformations, known for these molecules. In situ derivatization of sulfonates is particularly important as their purification is hampered because of their propensity to disproportionate in the presence of acid. However, the majority of these proven derivatization reactions had been developed on metal sulfinate salts. To validate our sulfinate synthesis we needed to ascertain the utility of the corresponding ammonium salts. Our initial findings, employing the ammonium sulfinate **2g**, are shown in Scheme 2. Alkylation with a range of alkyl halides was possible and was achieved by adding a DMF solution of the electrophile to the crude sulfinate salt, and examples employing α -bromo *tert*-butyl acetate and benzyl bromide are shown (**3g** and **4**). Cyclohexene oxide was efficiently opened to deliver the corresponding β -hydroxy-sulfone (**5**).^[23] Diarylsulfones were accessible by a combination with aryl iodonium salts,^[24] and treatment of **2g** with diphenyliodonium chloride provided the diarylsulfone (**6**). Aryl(heteroaryl)sulfones were available using an S_NAr-type process.^[25] Accordingly, reaction of **2g** with 2-chlorobenzothiazole provided the corresponding heterocyclic sulfone (**7**). Treatment of the ammonium sulfinate with sulfonyl chloride resulted in oxidative chlorination and allowed isolation of the derived sulfonyl chloride (**8**).^[26] The moderate yield achieved for this transformation reflects isolation issues. The derived sulfonyl chloride intermediates can be combined directly with an amine to deliver sulfonamides.^[26] For example, the morpholine derivative **9** was isolated in good yield. Finally, reduction using a Sn/HCl mixture allowed isolation of the corresponding disulfide (**10**).^[27] In general, the ammonium sulfinate products were found to deliver reactivity at least comparable to that of their metal-salt counterparts, and in several cases to show increased rates of reaction, presumably because of their increased solubility in organic solvents.

In summary, by employing the SO₂ surrogate DABSO, partnered with isopropyl alcohol as the reductant, we have



Scheme 2. Additional reactivity of ammonium sulfinate **2g**.

been able to develop a simple and efficient process for the palladium-catalyzed conversion of aryl and heteroaryl halides into the corresponding ammonium sulfonates. This effective sulfonation reaction employs low catalyst loadings, is operationally simple to perform, and has broad substrate scope. The ability to employ a simple alcohol as both reaction solvent and reductant significantly streamlines the process. The ammonium sulfonate products display good levels of reactivity and can be smoothly converted into a variety of sulfonyl derivatives. Further studies to develop related transformations, and to elucidate the mechanism of this transformation, are underway.

Received: April 21, 2014
Published online: July 25, 2014

Keywords: alcohols · arenes · palladium · sulfonamides · synthetic methods

- [1] For example: a) M. Teall, P. Oakley, T. Harrison, D. Shaw, E. Kay, J. Elliott, U. Gerhard, J. L. Castro, M. Shearman, R. G. Ball, N. N. Tsou, *Bioorg. Med. Chem. Lett.* **2005**, *15*, 2685; b) Y. Harrak, G. Casula, J. Basset, G. Rosell, S. Plescia, D. Raffa, M. G. Cusimano, R. Pouplana, M. D. Pujol, *J. Med. Chem.* **2010**, *53*, 6560; c) D. A. Smith, R. M. Jones, *Curr. Opin. Drug Discovery Dev.* **2008**, *11*, 72.
- [2] For example: a) W.-M. Xu, F.-F. Han, M. He, D.-Y. Hu, J. He, S. Yang, B.-A. Song, *J. Agric. Food Chem.* **2012**, *60*, 1036; b) Y. Noutoshi, M. Ikeda, T. Saito, H. Osada, K. Shirasu, *Front Plant Sci.* **2012**, *3*, 245; c) *Modern Crop Protection Compounds* (Eds.: W. Kramer, U. Schirmer), Wiley-VCH, Weinheim, **2007**, pp. 93–112.
- [3] N. S. Simpkins, *Sulphones in Organic Synthesis*, Pergamon Press, Oxford, **1993**.
- [4] “Polysulfones”: M. J. El-Hibri, S. A. Weinberg, *Encyclopedia of Polymer Science and Technology*, Wiley, New York, **2002**.
- [5] D. J. Ager, D. P. Pantaleone, S. A. Henderson, A. R. Katritzky, I. Prakash, D. E. Walters, *Angew. Chem.* **1998**, *110*, 1900; *Angew. Chem. Int. Ed.* **1998**, *37*, 1802.
- [6] For example: W. Su, *Tetrahedron Lett.* **1994**, *35*, 4955.
- [7] For example: D. J. Dale, P. J. Dunn, C. Golightly, M. L. Hughes, P. C. Levett, A. K. Pearce, P. M. Searle, G. Ward, A. Wood, *Org. Process Res. Dev.* **2000**, *4*, 17.
- [8] For recent alternative sulfone and sulfonamide syntheses, see: a) S. Cacchi, G. Fabrizi, A. Goggiamani, L. M. Parisi, R. Bernini, *J. Org. Chem.* **2004**, *69*, 5608; b) A. Kar, I. A. Sayyed, W. F. Lo, H. M. Kaiser, M. Beller, M. K. Tse, *Org. Lett.* **2007**, *9*, 3405; c) X. Zhao, E. Dimitrijevic, V. M. Dong, *J. Am. Chem. Soc.* **2009**, *131*, 3466; d) O. Saidi, J. Marafie, A. E. W. Ledger, P. Man Liu, M. F. Mahon, G. Kociok-Köhn, M. K. Whittlesey, C. G. Frost, *J. Am. Chem. Soc.* **2011**, *133*, 19298; e) C. G. Frost, J. P. Hartley, D. Griffin, *Synlett* **2002**, 1928; f) S. Caddick, J. D. Wilden, D. B. Judd, *J. Am. Chem. Soc.* **2004**, *126*, 1024.
- [9] G. Pelzer, W. Keim, *J. Mol. Catal. A* **1999**, *139*, 235.
- [10] a) H. Woolven, C. González-Rodríguez, I. Marco, A. L. Thompson, M. C. Willis, *Org. Lett.* **2011**, *13*, 4876; b) E. J. Emmett, B. R. Hayter, M. C. Willis, *Angew. Chem.* **2013**, *125*, 12911; *Angew. Chem. Int. Ed.* **2013**, *52*, 12679; c) A. S. Deeming, C. J. Russell, A. J. Hennessy, M. C. Willis, *Org. Lett.* **2014**, *16*, 150.
- [11] a) C. Waldmann, O. Schober, G. Haufe, K. Kopka, *Org. Lett.* **2013**, *15*, 2954; b) B. Rocke, K. B. Bahnck, M. Herr, S. Lavergne, V. Mascitti, C. Perreault, J. Polivkova, A. Shavnya, *Org. Lett.* **2014**, *16*, 154.
- [12] a) B. Nguyen, E. J. Emmett, M. C. Willis, *J. Am. Chem. Soc.* **2010**, *132*, 16372; b) E. J. Emmett, C. S. Richards-Taylor, B. Nguyen, A. Garcia-Rubia, B. R. Hayter, M. C. Willis, *Org. Biomol. Chem.* **2012**, *10*, 4007.
- [13] S. Ye, J. Wu, *Chem. Commun.* **2012**, *48*, 10037.
- [14] S. Ye, J. Wu, *Chem. Commun.* **2012**, *48*, 7753.
- [15] The Buchwald group have reported a related system in which phenyl chlorosulfate is combined with boronic acids to furnish sulfonyl chlorides: J. R. DeBergh, N. Niljianskul, S. L. Buchwald, *J. Am. Chem. Soc.* **2013**, *135*, 10638.
- [16] S. Patai, *The Chemistry of Sulphinic Acids, Esters and Their Derivatives*, Wiley, Chichester, **1990**.
- [17] C. S. Richards-Taylor, D. C. Blakemore, M. C. Willis, *Chem. Sci.* **2014**, *5*, 222.

- [18] A. Shavnya, S. B. Coffey, A. C. Smith, V. Mascitti, *Org. Lett.* **2013**, *15*, 6226.
- [19] See the Supporting Information for details.
- [20] A. Zapf, M. Beller, *Chem. Commun.* **2005**, 431.
- [21] Recent examples: a) R. L. Greenaway, C. D. Campbell, H. A. Chapman, E. A. Anderson, *Adv. Synth. Catal.* **2012**, *354*, 3187; b) J. Chen, Y. Zhang, L. Yang, X. Zhang, J. Liu, L. Li, H. Zhang, *Tetrahedron* **2007**, *63*, 4266; c) Y. Tsuchiya, Y. Hamashima, M. Sodeoka, *Org. Lett.* **2006**, *8*, 4851.
- [22] D. Prat, O. Pardigon, H.-W. Flemming, S. Letestu, V. Ducandas, P. Isnard, E. Guntrum, T. Senac, S. Ruisseau, P. Cruciani, P. Hosek, *Org. Process Res. Dev.* **2013**, *17*, 1517.
- [23] A. K. Maiti, P. Bhattacharyya, *Tetrahedron* **1994**, *50*, 10483.
- [24] N. Umierski, G. Manolikakes, *Org. Lett.* **2013**, *15*, 4972.
- [25] S. Liang, R.-Y. Zhang, L.-X. Xi, S.-Y. Chen, X.-Q. Yu, *J. Org. Chem.* **2013**, *78*, 11874.
- [26] R. Pandya, T. Murashima, L. Tedeschi, A. G. M. Barrett, *J. Org. Chem.* **2003**, *68*, 8274.
- [27] I. W. J. Still, I. D. G. Watson, *Synth. Commun.* **2001**, *31*, 1355.
-