

A Journal of the Gesellschaft Deutscher Chemiker

Angewandte Chemie

GDCh

International Edition

www.angewandte.org

Accepted Article

Title: Sulfonium Acids Loaded onto an Unusual Thiotemplate Assembly
Line Construct the Cyclopropanol Warhead of a Burkholderia
Virulence Factor

Authors: Felix Trottmann, Keishi Ishida, Jakob Franke, Aleksa Stanišić,
Mie Ishida-Ito, Hajo Kries, Georg Pohnert, and Christian
Hertweck

This manuscript has been accepted after peer review and appears as an Accepted Article online prior to editing, proofing, and formal publication of the final Version of Record (VoR). This work is currently citable by using the Digital Object Identifier (DOI) given below. The VoR will be published online in Early View as soon as possible and may be different to this Accepted Article as a result of editing. Readers should obtain the VoR from the journal website shown below when it is published to ensure accuracy of information. The authors are responsible for the content of this Accepted Article.

To be cited as: *Angew. Chem. Int. Ed.* 10.1002/anie.202003958

Link to VoR: <https://doi.org/10.1002/anie.202003958>

Sulfonium Acids Loaded onto an Unusual Thiotemplate Assembly Line Construct the Cyclopropanol Warhead of a *Burkholderia* Virulence Factor

Felix Trottmann, Keishi Ishida, Jakob Franke, Aleksa Stanišić, Mie Ishida-Ito, Hajo Kries, Georg Pohnert, and Christian Hertweck*

Abstract: Pathogenic bacteria of the *Burkholderia pseudomallei* group cause severe infectious diseases such as glanders and melioidosis. Malleicyprols were identified as important bacterial virulence factors, yet the biosynthetic origin of their cyclopropanol warhead has remained enigmatic. By a combination of mutational analysis and metabolomics we found that sulfonium acids, dimethylsulfoniumpropionate (DMSP) and gonyol, known as osmolytes and as crucial components in the global organosulfur cycle, are key intermediates en route to the cyclopropanol unit. Functional genetics and in vitro analyses uncover a specialized pathway to DMSP involving a rare prokaryotic SET-domain methyltransferase for a cryptic methylation, and show that DMSP is loaded onto the NRPS-PKS hybrid assembly line by an adenylation domain dedicated to zwitterionic starter units. Then, the megasynthase transforms DMSP into gonyol, as demonstrated by heterologous pathway reconstitution in *E. coli*.

Burkholderia pseudomallei group pathogens cause lethal infections that are difficult to treat.^[1] The best-known members of this pathogen complex are *Burkholderia mallei*, which causes the zoonotic disease glanders, and *B. pseudomallei*, the causative agent of melioidosis. The low infective dose needed and the possibility of infections through inhalation have led to the classification of *B. mallei* and *B. pseudomallei* as biological

warfare agents^[2] and a threat to global health.^[3] Since infections by these notorious pathogens are difficult to treat,^[4] novel therapeutic approaches such as anti-virulence strategies are needed. As prerequisite to disarming pathogens, it is essential to understand their virulence factors and the biosynthetic pathways involved.^[5] For *B. pseudomallei* and related pathogens such as the less virulent model organism *B. thailandensis*, various macromolecular^[6] and low-molecular-weight virulence factors^[3, 7] have been identified. Notably, all pathogens of the *B. pseudomallei* complex share a gene locus coding for an unusual modular thiotemplate assembly line with components of modular non-ribosomal peptide synthetases (NRPS) and polyketide synthases (PKS). Gene inactivation experiments unequivocally linked this gene cluster, named *bur*, to pathogenicity in nematode^[8] and mouse models.^[9] However, the first metabolite associated to this pathway - burkholderic acid^[10] *syn.* malleilactone^[8] (**1**) - did not exhibit any activity explaining the phenotypes observed in the infection models.

Recently, we found **1** to be the inactivated form of the true virulence factor, a highly reactive, cyclopropanol-substituted congener named bis-malleicyprol (**2a**, Figure 1A) formed from the monomer malleicyprol (**2b**).^[11] Nematode and toxicity assays showed dramatically increased activity of **2a** compared to **1**, implicating the cyclopropanol warhead in virulence.^[11] Thus, understanding its biosynthesis would set the basis for antivirulence strategies. According to stable-isotope labeling experiments and gene knockouts, the NRPS-PKS hybrid enzyme BurA assembles the cyclopropanol-containing fragment of **2b**, followed by dimerization to **2a**, which opens to form the inactive propanone-substituted unit of **1**, from a yet unknown methionine (**3**)-derived C3 building block and malonyl-CoA (Figure 1B).^[10] Yet, structures and biotransformations of the precursors loaded onto BurA have remained a riddle. Here we decipher the biogenetic origin of the rare cyclopropanol warhead of malleicyprol and show that a set of zwitterionic sulfonium acids initiates biosynthesis that play key roles in global sulfur cycling.

[*] F. Trottmann, Dr. K. Ishida, Dr. M. Ishida-Ito Prof. Dr. C. Hertweck
Department of Biomolecular Chemistry, Leibniz Institute for Natural
Product Research and Infection Biology (HKI)
Beutenbergstr. 11a, 07745 Jena, Germany
Fax: (+) 49 3641 5320804
E-mail: christian.hertweck@leibniz-hki.de
Homepage: www.leibniz-hki.de

A. Stanišić, Dr. H. Kries
Junior Research Group Biosynthetic Design of Natural Products,
Leibniz Institute for Natural Product Research and Infection Biology
(HKI)
Beutenbergstr. 11a, 07745 Jena, Germany

Prof. Dr. J. Franke
Institute of Botany,
Leibniz University Hannover, 30419 Hannover, Germany

Prof. Dr. G. Pohnert
Institute for Inorganic and Analytical Chemistry, Friedrich Schiller
University Jena, 07743 Jena, Germany

Prof. Dr. C. Hertweck
Natural Product Chemistry, Faculty of Biological Sciences,
Friedrich Schiller University Jena, 07743 Jena, Germany.

Supporting information and the ORCID identification number(s) for the
author(s) of this article can be found under: <https://doi.org/####>




Figure 1. A) Structures of burkholderic acid (1) and bis-malleicyprol (2a) featuring a cyclopropanol warhead. B) Isotope-labeling studies^[10] suggest acetate and methionine (3) as precursors to malleicyprol (2b), the monomer^[11] of 2a.

To elucidate the biogenetic origin of the malleicyprol warhead we first focused on the NRPS-PKS hybrid enzyme BurA and its unknown starter unit. Specifically, we tested for accumulation of intermediates in a mutant of the malleicyprol overexpressing strain *B. thailandensis* *Pbur* lacking a functional copy of *burA* (*B. thailandensis* *PburΔburA*^[10]). Candidates for the sulfur-containing pathway intermediates, however, could not be detected by routine HR-LCMS-based metabolic profiling. Only a comparative metabolomics analysis (*Pbur* vs. *PburΔburA*) searching for highly polar metabolites revealed the elusive methionine-derived starter unit. Compound 4 with *m/z* 135 accumulates in *B. thailandensis* *PburΔburA* cells (Figure 2A). Based on its exact mass (*m/z* 135.0474; [M+H]⁺) we deduced the molecular formula (C₅H₁₁O₂S) of 4. By comparison with authentic reference compounds we identified 4 as the zwitterionic compound dimethylsulfoniopropionate (DMSP; *m/z* 135.0474; [M+H]⁺; Supplementary Figure S1). This finding is intriguing since DMSP plays a pivotal role in the marine organosulfur cycle, serving as osmolyte for marine algae and as abundant carbon and sulfur source for bacteria.^[12] It is the precursor of the climate-relevant gas dimethylsulfide that is emitted at remarkably high amounts of >10⁷ tons per year into the atmosphere.^[13] Despite its wide distribution, DMSP has thus far not been implicated as a building block in natural product biosynthesis.




Figure 2. Identification of sulfonium intermediates in malleicyprol biosynthesis. A) Volcano plot analysis comparing pellet extracts of *B. thailandensis* variants *Pbur* and *PburΔburA*. B) Volcano plot analysis comparing supernatant extracts of *B. thailandensis* variants *Pbur* and *PburΔburA* (Supplementary Figure 2). C) Proposed DMSP biosynthesis in *B. thailandensis*. D) Production of two bis-malleicyprol (2a, *m/z* 611.3589) diastereomers in gene inactivation mutants of *B. thailandensis* *Pbur* monitored by UHPLC-MS (EIC in negative ion mode). E) Genomic alignment of the DMSP assembly line from *S. mobaraensis* to the *bur* biosynthetic gene cluster.

To confirm DMSP as a precursor of malleicyprols, we performed stable-isotope labeling experiments by chemically complementing suitable block mutants. Therefore, we required insight into the molecular basis of DMSP formation in *B. thailandensis*. By analogy to one of the established bacterial DMSP biosynthesis pathways^[12] (see ref.^[14] for alternative routes to DMSP) methionine would undergo S-methylation to form S-methylmethionine (5), decarboxylation, transamination and oxidation (Figure 2C). In silico analysis of the *bur* gene locus revealed candidate genes for a methyltransferase (BurB), a decarboxylase (BurI), a transaminase (BurD), and a dehydrogenase (BurE) (Figure 2E). Comparison of the deduced protein sequences to the recently published bacterial DMSP biosynthetic machinery in *Streptomyces mobaraensis*^[14b] showed BurD and BurE to be almost identical with their orthologues (96% and 98%), whereas BurB and BurI are only distantly related to their *S. mobaraensis* counterparts (35% and 48%).

To disrupt malleicyprol production, we individually inactivated each of the four putative DMSP biosynthesis genes in *B. thailandensis* *Pbur* by homologous crossover and replacement with a resistance cassette. The *ΔburD* and *ΔburE* mutants are still capable of producing the malleicyprol complex, albeit in reduced amounts (Figure 2D). Possibly, unspecific housekeeping enzymes take over transamination of sulfonium amine 6 and oxidation of the unstable aldehyde 7.^[14b] In contrast, the two main malleicyprol diastereomers are absent in the *ΔburB*

(methyltransferase) and $\Delta burI$ (decarboxylase) mutants (Figure 2D).

Closer inspection of culture extracts from $Pbur\Delta burI$ revealed enrichment of metabolite **5** (m/z 164.0744 in positive ion mode) that is identical to the unusual charged amino acid S-methylmethionine **5** (SMM; Figure 1D and Supplementary Figure S3). Since **5** cannot be detected in the mutant lacking *burB*, its gene product BurB must act in the biosynthetic pathway upstream of DMSP. BurB's role in DMSP formation is remarkable as it belongs to class V of the methyltransferase superfamilies containing a SET-domain.^[15] SET-domain-containing methyltransferases are well studied in eukaryotes and regulate gene expression through histone lysine methylation.^[15] Yet, little is known about prokaryotic SET-domain methyltransferases, and natural product modifying methyltransferases have exclusively been found in class I (Rossmann-like fold) or class III (tetrapyrrole methylase).^[16] To corroborate the function of BurB, we cloned and overexpressed *burB* in *Escherichia coli* and purified the His6-tagged protein via Ni-affinity. Incubation of purified BurB with L-methionine (**3**) and S-adenosylmethionine (SAM) generated SMM (Figure 3A), which was detected after derivatization with **8** to compound **9** (Figure 3A). Thus, BurB represents a novel C-S bond forming enzyme in secondary metabolism.^[17]

Having established the key steps to DMSP in *B. thailandensis* and with suitable null producers at hand, we performed chemical complementation. Therefore, we synthesized $^{13}\text{C}_3$ -labeled DMSP from $^{13}\text{C}_3$ acrylic acid (**10**, Figure 3C). Supplementation of the $Pbur\Delta burI$ mutant with $^{13}\text{C}_3$ -DMSP not only restored production of the malleicyprols (see Supplementary Figure S4) but also enriched ^{13}C in their cyclopropanol residue (Figure 3B). These results unequivocally confirm DMSP as a key intermediate in the formation of the malleicyprol warhead.




Figure 3. DMSP is a precursor of the cyclopropanol warhead of malleicyprol. A) Transformation of methionine (**3**) to S-methylmethionine (**5**; SMM) by BurB and derivatization with **8**; HR-LCMS detection of derivatized SMM; EIC m/z 330.0754 in positive ion mode; top: SMM derivatized with FDNB, middle: methylation of methionine with BurB, bottom: heat-inactivated BurB. B) Mass spectra of native bis-malleicyprol (top) and of ^{13}C -enriched bis-malleicyprol (bottom) C) Synthesis of ^{13}C -labeled DMSP and subsequent complementation of *B. thailandensis* $Pbur\Delta burI$ leads to incorporation of the C_3 unit into **2**.

Since DMSP accumulates in the $\Delta burA$ mutant, we reasoned that this unusual zwitterionic substrate would be activated and loaded onto the *bur* assembly line. In NRPS, adenylation (A) domains select, activate and load amino acids onto the assembly line and are thus regarded as gatekeepers.^[18] As bioinformatic substrate predictors^[19] failed on BurA-A we generated a homology model of the A domain. In this way we noted the absence of the conserved aspartate (D235 in GrsA-A) present in all α -amino acid activating A domains.^[18] Consequently, BurA-A was expected to select a non-canonical starter unit lacking the α -amino group usually bound by this aspartate. According to homology modeling, BurA-A shares important binding pocket features with the A-domain ATRR-A activating glycine betaine.^[20] In both ATRR-A and BurA-A, the loop carrying conserved D235 (GrsA-A) has been replaced with a shorter loop carrying hydrophobic residues (Figure 4A; Supplementary Figure S5). In another position, both binding pockets have an acidic residue (D606 in BurA-A) well placed to interact with a positively charged substrate moiety such as the sulfonium group of DMSP.

To verify whether BurA-A selects and activates DMSP we cloned and expressed the gene fragment for the A domain (*burA-A*) in *E. coli* BL21(DE3) and purified the His6-tagged protein via Ni-affinity. We probed the activity of BurA-A with the MesG/hydroxylamine assay, which monitors pyrophosphate released during substrate adenylation in a coupled photometric assay (Figure 4B).^[21] From a panel of carboxylic acids, BurA-A shows highest activity for DMSP. In stark contrast, neither a mixture of the proteinogenic amino acids, **5** nor 3-(methylthio)propionic acid (**11**) are activated (Figure 4C). Apparently, the positively charged sulfonium group enables substrate binding. Replacing this group with protonated nitrogen in **12**, thus maintaining the positive charge, reduces activity more than four-fold. In substrate saturation kinetics with DMSP, we determined a k_{cat} of 2.4 min^{-1} and a K_{M} of 0.15 mM (Supplementary Figure S7). Altogether, these results indicate that DMSP is the preferred substrate of BurA-A, making it the first adenylation domain that incorporates the osmolyte DMSP into a natural product assembly line.

An in silico analysis of the modular architecture of BurA suggested that DMSP, once loaded onto the thiolation (T) domain, would be elongated through Claisen condensation with malonyl-ACP, and the resulting β -keto intermediate transformed into the corresponding alcohol by the ketoreductase (KR) domain (Figure 4E). To identify the downstream product of DMSP we heterologously reconstituted the assembly line. Therefore, we cloned and expressed *burA* in *E. coli*. Only when we employed IPTG for induction and supplemented DMSP, we detected production of compound **13** with m/z 179.0742 (positive ion mode) in XAD16 extracts (Figure 4D). The presence of the same species in *B. thailandensis* $Pbur$ (Figure 1B) indicates that compound **13** is not an artifact generated in *E. coli* but actually formed in the intact *bur* pathway. By HR-LCMS and MS² comparison with an authentic reference we found that **13** is identical to the sulfonium acid gonyol.^[22] The structure of **13** agrees with our in silico prediction of the biosynthetic steps mediated by BurA. However, the discovery of this sulfonium intermediate of the malleicyprol assembly line is surprising, as **13** has been reported as a dominant zwitterion in the marine dinoflagellate *Gonyaulax polyedra*.^[22] Moreover, **13** is widely distributed as minor osmolyte in several phytoplankton groups.^[23] Previous studies have identified DMSP and acetate as the precursors of **13** in *G. polyedra*,^[22, 24] but the enzymes involved in gonyol biosynthesis have remained unknown. We now report BurA as the first enzyme involved in a gonyol biosynthetic

pathway, and unexpectedly, it is a modular NRPS-PKS hybrid. We reason that the sulfonium group represents a leaving group, likely as dimethylsulfide that enables cyclopropanol formation downstream of BurA. The enzymes and mechanisms involved in the cyclization are the subject of ongoing studies.




Figure 4. A) Sequence alignment of BurA-A with the canonical A domain GrsA and the glycine betaine accepting A domain ATRR. B) Concept of the MesG/hydroxylamine A domain assay C) Specificity profile of BurA-A; amino acid mix: all 20 proteinogenic amino acids (Supplementary Figure S6). D) Heterologous production of gonyol in *E. coli*; UHPLC-MS monitoring: EIC (*m/z* 179.0736) in positive ion mode; top, synthetic reference, middle, culture extracts of *E. coli* Rosetta2 expressing *burA* on plasmid pHIS8; bottom, *E. coli* Rosetta2 with empty pHIS8. E) Loading of DMSP onto BurA leads to production of gonyol and malleicypols.

In summary, we have uncovered crucial steps in the biosynthetic pathway to the virulence factor malleicypol employed by animal and human pathogenic *Burkholderia* species. Our findings have broad implications for ecology and synthetic biology. We describe BurB as a new C-S bond-forming enzyme in secondary metabolism that mediates a cryptic methylation to form the sulfonium group of the DMSP precursor. A role of this zwitterionic sulfonium acid in bacterial secondary metabolism is new. In contrast, DMSP is widely distributed in marine life, and metagenomics of known DMSP methyltransferase genes show that bacteria are significant producers of DMSP in marine environments.^[25] Our discovery of a new methyltransferase associated with DMSP biosynthesis and the identification of a

PKS-NRPS hybrid as a gonyol synthetase allows for genomics-based identification of ecologically relevant producer strains. From a biosynthetic perspective DMSP is noteworthy as a novel PKS primer unit,^[26] and the sulfonium-accepting adenylation domain BurA-A is an important addition to the synthetic biology toolbox that opens up new possibilities for engineering polyketides and nonribosomal peptides.

Acknowledgements

We thank A. Perner for LC-MS measurements. Financial support by the DFG (SFB 1127 ChemBioSys, and Leibniz Award to C. H.) and the European Regional Development Fund (ERDF) (MassNat) is gratefully acknowledged. J. F. acknowledges financial support by the SMART BIOTECs alliance between the Technische Universität Braunschweig and the Leibniz Universität Hannover, supported by the Ministry of and Culture (MWK) of Lower Saxony, Germany. H. K. received a fellowship from the Daimler and Benz foundation.

Conflict of interest

The authors declare no conflict of interest.

Keywords: Biosynthesis · DMSP · Mass spectrometry · NRPS · Virulence factors


- [1] G. C. Whitlock, D. Mark Estes, A. G. Torres, *FEMS Microbiol. Lett.* **2007**, 277, 115-122.
- [2] J. Gilad, I. Harary, T. Dushnitsky, D. Schwartz, Y. Amsalem, *Isr. Med. Assoc. J.* **2007**, 9, 499-503.
- [3] J. B. Biggins, H.-S. Kang, M. A. Ternei, D. DeShazer, S. F. Brady, *J. Am. Chem. Soc.* **2014**, 136, 9484-9490.
- [4] D. Dance, *Int. J. Antimicrob. Ag.* **2014**, 43, 310-318.
- [5] R. Thänert, O. Goldmann, A. Beineke, E. Medina, *Nat. Commun.* **2017**, 8, 14268.
- [6] a) A. Cruz-Migoni, G. M. Hautbergue, P. J. Artymiuk, P. J. Baker, M. Bokori-Brown, C.-T. Chang, M. J. Dickman, A. Essex-Lopresti, S. V. Harding, N. M. Mahadi, L. E. Marshall, G. W. Mobbs, R. Mohamed, S. Nathan, S. A. Ngugi, C. Ong, W. F. Ooi, L. J. Partridge, H. L. Phillips, M. F. Raih, S. Ruzhenikov, M. Sarkar-Tyson, S. E. Sedelnikova, S. J. Smither, P. Tan, R. W. Titball, S. A. Wilson, D. W. Rice, *Science* **2011**, 334, 821-824; b) W. J. Wiersinga, H. S. Virk, A. G. Torres, B. J. Currie, S. J. Peacock, D. A. B. Dance, D. Limmathurotsakul, *Nat. Rev. Dis. Prim.* **2018**, 4, 17107.
- [7] a) H. Yang, C. D. Kooi, P. A. Sokol, *Infect. Immun.* **1993**, 61, 656-662; b) A. F. Alice, C. S. López, C. A. Lowe, M. A. Ledesma, J. H. Crosa, *J. Bacteriol.* **2006**, 188, 1551-1566; c) J. Franke, K. Ishida, M. Ishida-Ito, C. Hertweck, *Angew. Chem. Int. Ed.* **2013**, 52, 8271-8275; d) J. Franke, K. Ishida, C. Hertweck, *J. Am. Chem. Soc.* **2014**, 136, 5599-5602; e) J. Franke, K. Ishida, C. Hertweck, *Chemistry* **2015**, 21, 8010-8014; f) R.-R. Wong, C. Kong, S.-H. Lee, S. Nathan, *Sci. Rep.* **2016**, 6, 27475; g) F. Trottmann, J. Franke, K. Ishida, M. García-Altare, C. Hertweck, *Angew. Chem. Int. Ed.* **2019**, 58, 200-204.
- [8] a) J. B. Biggins, M. A. Ternei, S. F. Brady, *J. Am. Chem. Soc.* **2012**, 134, 13192-13195; b) J. R. Klaus, J. Deay, B. Neuenswander, W. Hursh, Z. Gao, T. Boudhdhara, T. D. Williams, J. Douglas, K. Monize, P. Martins, C. Majerczyk, M. R. Seyedsayamdost, B. R. Peterson, M. Rivera, J. R. Chandler, *J. Bacteriol.* **2018**, 200, e00008-18.
- [9] a) M. G. Gutierrez, D. R. Yoder-Himes, J. M. Warawa, *Front. Cell. Infect. Microbiol.* **2015**, 5; b) M. G. Moule, N. Spink, S. Willcocks, J. Lim, J. A. Guerra-Assunção, F. Cia, O. L. Champion, N. J. Senior, H. S. Atkins, T. Clark, G. J.

- Bancroft, J. Cuccui, B. W. Wren, *Infect. Immun.* **2016**, *84*, 701-710.
- [10] a) J. Franke, K. Ishida, C. Hertweck, *Angew. Chem. Int. Ed.* **2012**, *51*, 11611-11615; b) Note: Met accumulation in media leads to increased malleilactone production: A. Li, D. Mao, A. Yoshimura, P. C. Rosen, W. L. Martin, E. Gallant, M. Wühr, M. R. Seyedsayamdost, *mBio* **2020**, *11*, e03210-19.
- [11] F. Trottman, J. Franke, I. Richter, K. Ishida, M. Cyrulies, H.-M. Dahse, L. Regestein, C. Hertweck, *Angew. Chem. Int. Ed.* **2019**, *58*, 14129-14133.
- [12] M. A. Moran, B. P. Durham, *Nat. Rev. Microbiol.* **2019**, *17*, 665-678.
- [13] A. J. Kettle, M. O. Andreae, *J. Geophys. Res.* **2000**, *105*, 26793-26808.
- [14] a) A. R. J. Curson, B. T. Williams, B. J. Pinchbeck, L. P. Sims, A. B. Martínez, P. P. L. Rivera, D. Kumaresan, E. Mercadé, L. G. Spurgin, O. Carrión, S. Moxon, R. A. Cattolico, U. Kuzhiumparambil, P. Guagliardo, P. L. Clode, J.-B. Raina, J. D. Todd, *Nat. Microbiol.* **2018**, *3*, 430-439; b) C. Liao, F. P. Seebeck, *Angew. Chem. Int. Ed.* **2019**, *58*, 3553-3556.
- [15] a) R. Alvarez-Venegas, *Front. Genet.* **2014**, *5*, 65-65; b) C. Nwasike, S. Ewert, S. Jovanovic, S. Haider, S. Mujtaba, *Ann. N.Y. Acad. Sci.* **2016**, *1376*, 18-28.
- [16] D. K. Liscombe, G. V. Louie, J. P. Noel, *Nat. Prod. Rep.* **2012**, *29*, 1238-1250.
- [17] K. L. Dunbar, D. H. Scharf, A. Litomska, C. Hertweck, *Chem. Rev.* **2017**, *117*, 5521-5577.
- [18] a) G. L. Challis, J. Ravel, C. A. Townsend, *Chem. Biol.* **2000**, *7*, 211-224; b) F. Kudo, A. Miyanaga, T. Eguchi, *J. Ind. Microbiol. Biotechnol.* **2019**, *46*, 515-536; c) T. Stachelhaus, H. D. Mootz, M. A. Marahiel, *Chem. Biol.* **1999**, *6*, 493-505.
- [19] a) C. N. Boddy, *J. Ind. Microbiol. Biotechnol.* **2014**, *41*, 443-450; b) M. Röttig, M. H. Medema, K. Blin, T. Weber, C. Rausch, O. Kohlbacher, *Nucleic Acids Res.* **2011**, *39*, W362-W367.
- [20] Y. Hai, A. M. Huang, Y. Tang, *Proc. Natl. Acad. Sci. U.S.A.* **2019**, *116*, 10348-10353.
- [21] B. P. Duckworth, D. J. Wilson, C. C. Aldrich, *Methods Mol. Biol.* **2016**, *1401*, 53-61.
- [22] H. Nakamura, K. Fujimaki, O. Sampei, A. Murai, *Tetrahedron Lett.* **1993**, *34*, 8481-8484.
- [23] B. Gebser, G. Pohnert, *Mar. Drugs* **2013**, *11*, 2168-2182.
- [24] H. Nakamura, T. Jin, M. Funahashi, K. Fujimaki, O. Sampei, A. Murai, T. Roenneberg, J. W. Hastings, *Tetrahedron* **1997**, *53*, 9067-9074.
- [25] L. Zha, Y. Jiang, M. T. Henke, M. R. Wilson, J. X. Wang, N. L. Kelleher, E. P. Balskus, *Nat. Chem. Biol.* **2017**, *13*, 1063-1065.
- [26] B. T. Williams, K. Cowles, A. Bermejo Martínez, A. R. J. Curson, Y. Zheng, J. Liu, S. Newton-Payne, A. J. Hind, C.-Y. Li, P. P. L. Rivera, O. Carrión, J. Liu, L. G. Spurgin, C. A. Brearley, B. W. Mackenzie, B. J. Pinchbeck, M. Peng, J. Pratscher, X.-H. Zhang, Y.-Z. Zhang, J. C. Murrell, J. D. Todd, *Nat. Microbiol.* **2019**, *4*, 1815-1825.
- [27] B. S. Moore, C. Hertweck, *Nat. Prod. Rep.* **2002**, *19*, 70-99.

COMMUNICATION

Sulfur cycle meets pathogen.

Sulfonium acids known from global sulfur cycles were elucidated as the biosynthetic origin of the cyclopropanol warhead of malleicyprols, virulence factors of human-pathogenic bacteria. The pathway involves a specialized S-methyl transferase to form DMSP, which is activated by a zwitterion-specific adenylation domain and transformed into the key intermediate gonyol.



F. Trottman, K. Ishida, J. Franke, A. Stanišić, M. Ishida-Ito, H. Kries, G. Pohnert, C. Hertweck*

Page No. – Page No.

Sulfonium Acids Loaded onto Unusual Thiotemplate Assembly Line Constitute Cyclopropanol Warhead of a *Burkholderia* Virulence Factor

Institute and/or researcher Twitter usernames: @c_hertweck; @Hertweck_Lab; @KriesLab; @LeibnizHKI