This article was downloaded by: [Moskow State Univ Bibliote] On: 15 February 2014, At: 00:25 Publisher: Taylor & Francis Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



# Organic Preparations and Procedures International: The New Journal for Organic Synthesis

Publication details, including instructions for authors and subscription information: http://www.tandfonline.com/loi/uopp20

## A CONVENIENT SYNTHESIS OF 2(5H)-FURANONE

Ruzhen Cao<sup>a</sup>, Chong Liu<sup>a</sup> & Lunzu Liu<sup>a</sup> <sup>a</sup> Institute of Elemento-Organic Chemistry, Nankai University, Tianjin, 300071, PR CHINA Published online: 18 Feb 2009.

To cite this article: Ruzhen Cao, Chong Liu & Lunzu Liu (1996) A CONVENIENT SYNTHESIS OF 2(5H)-FURANONE, Organic Preparations and Procedures International: The New Journal for Organic Synthesis, 28:2, 215-216, DOI: <u>10.1080/00304949609356524</u>

To link to this article: http://dx.doi.org/10.1080/00304949609356524

### PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <a href="http://www.tandfonline.com/page/terms-and-conditions">http://www.tandfonline.com/page/terms-and-conditions</a>

#### A CONVENIENT SYNTHESIS OF 2(5H)-FURANONE

Submitted by (07/20/95)

Ruzhen Cao, Chong Liu and Lunzu Liu\*

Institute of Elemento-Organic Chemistry, Nankai University Tianjin 300071, P. R. CHINA

2(5H)-Furanone (2) has found extensive use as an important chemical,<sup>1</sup> and recently has attracted considerable attention as a plant growth regulator,<sup>2</sup> antiulcer agent<sup>3</sup> and fish growth promoter.<sup>4</sup> It has been prepared by oxidation of furfural with hydrogen peroxide in the presence of formic acid,<sup>5,6</sup> thiourea<sup>7</sup> or cobaltous oxide.<sup>8</sup> Although these methods are one-pot reactions, they are deficient in some respects: the reagents are more expensive, the handling is inconvenient and perhaps most importantly the yield is much lower. We now describe a convenient method which involved refluxing furfural and hydrogen peroxide in dichloroethane and sodium sulfate. This reaction pathway may be described as shown in below.



The oxidation afforded a mixture of 2(3H)-furanone (1) and 2(5H)-furanone (2) which was separated by distillation. Isomerization of 1 with  $Et_3N$  afforded 2 in high yield. In order to avoid hydrolysis of 2 to organic acids, the reaction was carried out in a two phase solution (dichloroethane-sat.  $Na_2SO_4$ ) to give 2 in good yield. However, with saturated sodium chloride a very low yield of 2 was obtained presumably due to increased hydrolysis.

#### **EXPERIMENTAL SECTION**

<sup>1</sup>H NMR spectra were recorded on a JEOL FX-90Q spectrometer with TMS as the internal standard.

**Preparation of 2(5H)-Furanone (2).** In a 1L four-necked flask fitted with a reflux condenser, a mechanical stirrer, a thermometer and a dropping funnel were placed redistilled furfural (96g, 1mol), dichloroethane (200mL) and sodium sulfate (120g). Then 30% hydrogen peroxide (220mL, 2.13mol) was added dropwise with stirring at  $60~70^{\circ}$  over a period of 2 hrs. After addition, the reaction mixture was heated to reflux for 10 hrs, then cooled to room temperature and filtered. The organic phase of the filtrate was separated, and the aqueous layer was extracted with dichloroethane (2 x 50mL). Triethy-lamine (2~4g) was added to the combined dichloroethane layers containing 1 and 2. The mixture was stirred for 2 hrs at 40°. After removal of the dichloroethane, the residual liquid was distilled *in vacuo* 

to give 2 (56g, yield 67%). To separate compounds 1 and 2, the dichloroethane layer was concentrated on a rotary evaporator and the residual liquid was distilled *in vacuo* to give 2(3H)-furanone (15g, yield 18%) and 2(5H)-furanone (31g, yield 37%).

2(3H)-Furanone (1): bp. 60-62°/40mmHg. <sup>1</sup>H NMR(CDCl<sub>3</sub>):  $\delta$  3.10(t, 2H, <sup>3</sup>J = 2.4, <sup>4</sup>J = 2.4, 3-H), 5.52(dt, 1H, <sup>3</sup>J = 3.6, <sup>4</sup>J = 2.4, 5-H), 6.74(dt, 1H, <sup>3</sup>J = 2.4, <sup>3</sup>J = 3.6, 4-H). *Anal.* Calcd for C<sub>4</sub>H<sub>4</sub>O<sub>2</sub>: C, 57.14; H, 4.76. Found: C, 57.01; H, 4.62 2(5H)-Furanone (2): bp. 98-100°/40mmHg. <sup>1</sup>H NMR(CDCl<sub>3</sub>):  $\delta$  4.88(dd, 2H, <sup>4</sup>J = 2.2, <sup>3</sup>J = 1.7, 5-H), 6.10(dt, 1H, <sup>4</sup>J = 2.2, <sup>3</sup>J = 5.8, 3-H), 7.58(dt, 1H, <sup>3</sup>J = 1.7, <sup>3</sup>J = 5.8, 4-H). *Anal.* Calcd for C<sub>4</sub>H<sub>4</sub>O<sub>2</sub>: C, 57.14; H, 4.76. Found: C, 57.10; H, 4.71

#### REFERENCES

- 1. Y. S. Rao, Chem. Rev., 76, 625 (1976).
- V. V. Nemchenko and N. P. Ivanova, *Khim. Sel'sk. Khoz.*, 1, 91 (1991); *Chem. Abs.*, 115, 44136 (1991).
- 3. H. Aihara, M. Muramatsu and I. Arai, Jpn Kokai Tokkyo Koho, 225315 (1988); Chem. Abs., 110, 63776 (1989).
- 4. L. N. Tsen and V. S. Suleimanyan, Rybn. Khoz. (Moscow), 5, 42 (1982); Chem. Abs., 97, 22625 (1982).
- 5. L. A. Badovskaya, Khim. Geterotsikl. Soedin., 10, 1314 (1978); Chem. Abs., 90, 54751 (1979).
- 6. H. Jan-Anders, K. Nasman and G. Pensar, Synthesis, 786 (1985).
- 7. Zh. T. Lavlenko and S.V. Zhuravlev, Ger(East)., 274333 (1989); Chem. Abs., 113, 42808 (1990).
- 8. L. A. Badovskaya and G.F. Myzychenko, USSR, 470516 (1975); Chem. Abs., 83, 113700 (1975).

\*\*\*\*\*