

Domino Synthesis of Thiochromenes through Cu-Catalyzed Incorporation of Sulfur using Xanthate Surrogate

Pandi Muthupandi, Nallappan Sundaravelu, and Govindasamy Sekar

J. Org. Chem., **Just Accepted Manuscript** • DOI: 10.1021/acs.joc.6b02740 • Publication Date (Web): 26 Jan 2017

Downloaded from <http://pubs.acs.org> on January 27, 2017

Just Accepted


“Just Accepted” manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides “Just Accepted” as a free service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. “Just Accepted” manuscripts appear in full in PDF format accompanied by an HTML abstract. “Just Accepted” manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are accessible to all readers and citable by the Digital Object Identifier (DOI®). “Just Accepted” is an optional service offered to authors. Therefore, the “Just Accepted” Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the “Just Accepted” Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these “Just Accepted” manuscripts.

Domino Synthesis of Thiochromenes through Cu-Catalyzed Incorporation of Sulfur using Xanthate Surrogate

Pandi Muthupandi, Nallappan Sundaravelu and Govindasamy Sekar*

Department of Chemistry, Indian Institute of Technology Madras, Chennai-600036, Tamil Nadu, India

gsekar@iitm.ac.in



An efficient domino reaction has been developed for the synthesis of thiochromenes through Cu-catalyzed in situ incorporation of sulfur. This domino method avoids the use of less accessible and unpleasant arenethiols as starting materials, instead utilizes very stable aryl halides along with potassium ethyl xanthate as an odorless sulfur surrogate. The domino methodology proceeds through $\text{C}_{(\text{aryl})}\text{-S}$ coupling, thioester cleavage, sulfa-Michael addition, aldol reaction and elimination reaction sequences to provide thiochromenes in good yields


INTRODUCTION

Synthesis of sulfur-containing compounds continues to be an important task in organic chemistry because of their applications in biology, material science and food chemistry.¹⁻³ Thiochromenes, the sulfur analogues of chromenes, have been testified to exhibit many pharmaceutical properties such as anti-cancer, anti-bacterial, anti-inflammatory and anti-psychiatric activities.⁴ Conventionally, thiochromenes could be prepared by four step process such as Michael addition of thiophenol to 2-alkenoic acids followed by acid-catalyzed cyclization, reduction and dehydration.⁵ Only a handful of attempts have been made to achieve efficient method for the


1
2
3 synthesis of thiochromenes.⁶ However, most of these methods suffer from a number of
4
5 drawbacks like requirement of multistep synthesis, limited substrate scope and lower yield.
6
7

8 In the past years, domino sulfa-Michael-aldol reaction of 2-mercaptobenzaldehydes with α,β -
9
10 unsaturated compounds has been used as an efficient tool for the synthesis of chiral
11
12 thiochromanes.⁷ However, very few methods utilized cascade sulfa-Michael-aldol reaction to
13
14 synthesize chiral thiochromenes,⁸ cascade sulfa-Michael-aldol reaction of 2-
15
16 mercaptobenzaldehydes with simple chalcones has remained unexplored. Despite the progress
17
18 towards synthesis of thiochromenes, all the reported methods utilize arenethiols as starting
19
20 materials which are less accessible, produce unpleasant smell and concern environmental safety.
21
22 Therefore, development of a catalytic method which overcomes all above mentioned difficulties
23
24
25
26
27 would be highly desirable.

28
29 In recent years, considerable effort has been devoted towards transition metal catalyzed C_(aryl)-S
30
31 bond formation by means of in situ generation of sulfur.⁹ As a result, construction of sulfur-
32
33 containing heterocyclic compounds involving concomitant formation of C-S bonds has been
34
35 developed to some extent.¹⁰ We have previously used xanthates as coupling partner with aryl
36
37 halides under copper catalysis as a means of incorporating sulfur.¹¹ We envisioned a domino
38
39 process for the synthesis of thiochromenes through Cu-catalyzed in situ incorporation of sulfur
40
41 using easily available and odourless potassium ethyl xanthate as source of sulfur (Scheme 1). In
42
43 this strategy, coupling of 2-halobenzaldehyde and xanthate can be achieved by copper catalyst.
44
45 Thioester cleavage could occur in the presence of base to generate 2-formylbenzenethiolate
46
47 which can undergo sulfa-Michael addition with chalcone followed by aldol reaction and then
48
49 base can facilitate the dehydration.
50
51
52
53
54
55
56
57
58
59
60

Scheme 1. Our strategy for Cu-catalyzed domino synthesis of thiochromenes**RESULTS AND DISCUSSION**

Initially, 2-iodobenzaldehyde was reacted with simple chalcone in the presence of 2 equivalents of potassium ethyl xanthate, 10 mol% of $\text{Cu}(\text{OAc})_2$ and 2 equivalents of NaOH in DMSO solvent at 100 °C. The reaction proceeded to completion after 14 h, and the desired thiochromene was isolated in 23% yield (Table 1, entry 1). The structure of **3a** was unambiguously confirmed by single-crystal XRD analysis (see the Supporting Information, Figure 1). Encouraged by this result, optimization of the reaction was performed with respect to Cu source, solvents and bases. When $\text{Cu}(\text{OAc})_2$ was replaced with other copper salts such as CuI, CuCl, CuBr_2 and CuCl_2 , the yield was not improved (entries 2-5).

Table 1. Optimization of reaction conditions^a


Entry	Cu salt	Additive	Solvent	Time (h)	Yield (%) ^b
1	$\text{Cu}(\text{OAc})_2$	NaOH	DMSO	14	23 ^c
2	CuI	NaOH	DMSO	24	8 ^c
3	CuCl	NaOH	DMSO	24	7 ^c
4	CuBr_2	NaOH	DMSO	24	11 ^c
5	CuCl_2	NaOH	DMSO	24	16 ^c
6	$\text{Cu}(\text{OAc})_2$	NaOH	PhMe	24	nd ^d
7	$\text{Cu}(\text{OAc})_2$	NaOH	Dioxane	24	nd ^d
8	$\text{Cu}(\text{OAc})_2$	NaOH	CH_3CN	24	nd ^d
9	$\text{Cu}(\text{OAc})_2$	NaOH	DMF	14	38 ^c
10	$\text{Cu}(\text{OAc})_2$	KOH	DMF	14	34 ^c
11	$\text{Cu}(\text{OAc})_2$	LiOH	DMF	18	25 ^c
12	$\text{Cu}(\text{OAc})_2$	Na_2CO_3	DMF	24	21 ^c
13	$\text{Cu}(\text{OAc})_2$	K_3PO_4	DMF	24	28 ^c
14	$\text{Cu}(\text{OAc})_2$	NaOAc	DMF	14	62 ^c
15	$\text{Cu}(\text{OAc})_2$	NaO^tBu	DMF	3	- ^e
16	$\text{Cu}(\text{OAc})_2$	Et_2NH	DMF	5	- ^e
17	$\text{Cu}(\text{OAc})_2$	DABCO	DMF	5	- ^e
18	$\text{Cu}(\text{OAc})_2$	AgOAc	DMF	9	79
19	$\text{Cu}(\text{OAc})_2$	AgOAc	DMF	7	92^f

^aReaction conditions: **1a** (0.5 mmol), **2a** (0.5 mmol), xanthate (1 mmol), $\text{Cu}(\text{OAc})_2$ (10 mol%), additive (1 mmol), solvent (2 mL), 100 °C. ^bIsolated yield. ^cThe reaction generated a complex mixture from which pure product could nevertheless be isolated. ^dStarting materials remained. ^eThe reaction generated an complex, inseparable mixture from which pure product could not be isolated. ^f1.5 mmol of xanthate and 0.75 mmol of AgOAc were used.

1
2
3 When the reaction was conducted in different solvents such as toluene, dioxane and acetonitrile,
4
5 the thiochromene formation was not observed in these solvents (entries 6-8). Interestingly, the
6
7 thiochromene was isolated in 38% yield, when the reaction was carried out in DMF solvent
8
9 (entry 9). Further optimization was performed in DMF solvent, as it provided better yield than
10
11 DMSO solvent. Replacement of NaOH with other bases such as KOH and LiOH resulted in
12
13 lower yield (entries 10 and 11). The starting materials were not consumed completely when
14
15 Na₂CO₃ and K₃PO₄ were used as base (entries 12 and 13). To our delight, thiochromene was
16
17 obtained in 62% yield, when NaOAc was used (entry 14). Other bases such as NaO^tBu, Et₂NH
18
19 and DABCO gave inseparable complex reaction mixtures (entries 15-17). Remarkable
20
21 improvement in yield (81%) was observed when AgOAc was employed as base (entry 18). After
22
23 studying the reaction condition with respect to temperature, equivalents of Cu(OAc)₂, xanthate
24
25 and base, the domino reaction provided 92% yield for thiochromene in the presence of 10 mol%
26
27 Cu(OAc)₂, 3 equivalents of xanthate and 1.5 equivalents AgOAc in DMF at 100 °C (entry 19).

28
29
30
31
32
33
34 Next, substrate scope of this domino reaction was explored under the optimized reaction
35
36 conditions. The domino reaction proceeded smoothly with a variety of other substrates, and the
37
38 corresponding thiochromenes were isolated in good yields (Scheme 2). Electron-releasing
39
40 substituents on the aryl ring that is attached to alkene part of chalcone facilitated the domino
41
42 reaction, and thiochromenes **3b-3e** were obtained in good to excellent yields. Chalcones bearing
43
44 chloro, bromo and fluoro substituents, importantly at *ortho* position of aryl ring reacted
45
46 effortlessly and provided the corresponding thiochromenes (**3f-3h**) in good yields.
47
48
49
50
51
52
53
54
55
56
57
58
59
60


Scheme 2. Scope of the domino reaction with respect to 2-iodobenzaldehyde



Electron-withdrawing group containing chalcone was also tolerated in the domino reaction and afforded thiochromene **3i** in 82% yield. Replacement of the phenyl group that is attached to the alkene part of the chalcone with naphthyl, thiophenyl and pyridyl groups gave the corresponding thiochromenes **3j-3l** in good to excellent yields. Similarly, electron-releasing and electron-withdrawing substituents on the aryl ring that is attached to carbonyl part of chalcone were well tolerated in this domino reaction, and the corresponding thiochromenes **3m-3r** were isolated in decent yields. Chalcones having naphthyl, thiophenyl and methyl groups attached to carbonyl part reacted easily to give thiochromene **3s**, **3t** and **3u**. Notably, cinnamaldehydes and crotonaldehyde were found to be effective substrates for this domino reaction (**3v-3x**). Also, 2-iodobenzaldehyde having bromo, chloro or methyl substituents offered thiochromenes **3y**, **3z** and **3aa** in excellent yields.

Subsequently, the domino synthesis of thiochromenes was performed using comparatively less reactive 2-bromobenzaldehyde **4** as starting material. The domino reaction gave thiochromene in 67% yield under the standard optimized conditions. However, the yield was increased to 81% when the reaction was conducted at 120 °C (Scheme 3).


Scheme 3. Scope of the domino reaction with respect to 2-bromobenzaldehyde



The generality of domino synthesis of thiochromenes using less reactive 2-bromobenzaldehyde **4** as starting material was explored with various substituted chalcones. All the substituents were well tolerated and the corresponding thiochromenes (**3a**, **3ab**, **3i**, **3d**, **3ac** and **3ad**) were obtained in good yields.


In order to showcase the efficacy of domino synthesis of thiochromenes on gram scale, the reaction was performed in 5 mmol (1.16 g) scale under standard optimized condition without altering any reaction parameters (Scheme 4). The reaction proceeded smoothly and provided the thiochromene in 86% yield.

Scheme 4. Gram-scale synthesis



To gain insight into the possible reaction pathway, control experiments were performed. When the domino reaction was performed without AgOAc, formation of thiochromene **3a** was not observed, instead thiochromane **5a** was isolated in 19% yield as mixture of two diastereomers along with diarylsulfide **6a** (32% yield) (Scheme 5a). Also, thiochromene **3a** formation was not observed in the absence of both Cu(OAc)₂ and AgOAc (Scheme 5b).


Scheme 5. Control experiments



These experiments reveal that Cu(OAc)_2 promotes $\text{C}_{(\text{aryl})}\text{-S}$ coupling but the Lewis acidity is not adequate to activate chalcone to undergo sulfa-Michael-aldol reaction. Thus, coupling of in situ formed 2-formylbenzenethiolate with 2-iodobenzaldehyde might be responsible for the formation of diarylsulfide **6a**. In order to understand the importance of AgOAc , the domino reaction was performed with other silver salts. The domino reaction with AgBr and Ag_2CO_3 did not provide thiochromene **3a** instead thiochromane **5a** and diarylsulfide **6a** (Scheme 5c) were isolated. The dehydration of thiochromane **5a** in the presence of AgOAc gave 11% yield for product **3a** and thiochromane **5a** remained without reacting even after 12 hours (Scheme 5d). These control experiments show that Cu(OAc)_2 and AgOAc are essential for the domino reaction to complete. From these observations, a possible reaction pathway has been proposed as shown in Scheme 6. Oxidative addition of copper acetate with 2-iodobenzaldehyde may lead to intermediate **A** which undergoes ligand exchange with potassium ethyl xanthate to generate

intermediate **B**. Subsequently, reductive elimination of intermediate **B** affords intermediate **C** which may undergo thioester cleavage and the reaction of resulting thiolate with chalcone **2a** by Michael-aldol-elimination reaction sequence in the presence of AgOAc to give the product **3a**.

Scheme 6. Possible reaction pathway



CONCLUSION

In summary, we have accomplished the domino synthesis of thiochromenes using Cu-catalyzed in situ incorporation of sulfur as a key step. Stable and odourless potassium ethyl xanthate has been utilized as an efficient sulfur source. The domino methodology proceeds through C_(aryl)-S coupling, thioester cleavage, sulfa-Michael addition, aldol reaction and elimination reaction sequences to provide thiochromenes in good yields. A variety of thiochromenes have been synthesized in good to excellent yields from commercially available and easily accessible starting materials and catalyst. The gram scale synthesis and avoidance of foul-smelling arenethiols make this domino process practical and environmentally friendly.

EXPERIMENTAL SECTION

General information. All reactions were carried out in oven-dried reaction tubes. Reactions were monitored by thin-layer chromatography (TLC) using silica gel 60 F₂₅₄ precoated plates

1
2
3 (0.25 mm) and visualized by UV fluorescence. Column chromatography was performed using
4
5 Silica gel (particle size: 100-200 mesh) and eluted with hexanes/ethyl acetate mixture. All the
6
7 reactions were carried out in temperature controlled magnetic stirrers. ^1H and ^{13}C NMR spectra
8
9 were recorded on 400 MHz and 500 MHz (100 MHz and 125 MHz for ^{13}C) instrument. ^1H NMR
10
11 spectra were reported relative to residual CDCl_3 (δ 7.26 ppm). When the residual peak is
12
13 overlapping with compound, spectra reported to residual TMS. ^{13}C NMR were reported relative
14
15 to CDCl_3 (δ 77.16 ppm). Chemical shifts were reported in parts per million and coupling
16
17 constants (J) are reported in Hertz. Melting points were recorded on melting point apparatus and
18
19 are corrected with benzoic acid as reference. Infrared spectra were recorded on a FTIR 4000
20
21 Series spectrometer using dry KBr pellet. The wave numbers of recorded IR signals are quoted in
22
23 cm^{-1} . High resolution mass spectra (HRMS) were recorded on Q-TOF Micro mass spectrometer.
24
25 $\text{Cu}(\text{OAc})_2$ and potassium ethyl xanthogenate were obtained from commercial source and used
26
27 directly as received.
28
29
30
31
32
33
34

35 **General procedure for the synthesis of thiochromenes.** A oven dried reaction tube was loaded
36
37 with 2-iodobenzaldehyde (0.5 mmol), chalcone (0.5 mmol), potassium ethyl xanthate (1.5
38
39 mmol), AgOAc (1.5 mmol) and $\text{Cu}(\text{OAc})_2$ (0.05 mmol) and then DMF (2mL) was added. The
40
41 reaction tube was closed with glass-stopper and stirred at 100 °C (120 °C in case of 2-
42
43 bromobenzaldehyde) for recommended time. The reaction mixture was brought to room
44
45 temperature and diluted with ethyl acetate and then washed with brine. The aqueous layer was
46
47 extracted twice with ethyl acetate and the combined organic extractions were dried over
48
49 anhydrous Na_2SO_4 . Solvent was removed under reduced pressure and the residue was purified by
50
51 silica gel column using hexanes/ethyl acetate (90:10, v/v) mixture to afford the thiochromene.
52
53
54
55
56
57
58
59
60

1
2
3 [2-(Phenyl)-2H-thiochromen-3-yl](phenyl)methanone (3a). 152mg, 92% yield; pale yellow solid;
4
5 mp 156-158 °C; R_f 0.52 (10% ethyl acetate in hexanes); ^1H NMR (CDCl_3 , 400 MHz) δ 5.48 (s,
6
7 1H), 7.10-7.15 (m, 1H), 7.17-7.30 (m, 8H), 7.39 (s, 1H), 7.46 (t, $J = 7.6$ Hz, 2H), 7.56 (tt, $J = 7.6$
8
9 Hz, 2.4 Hz, 1H), 7.69 (d, $J = 7.8$ Hz, 2H); ^{13}C NMR (CDCl_3 , 100 MHz) δ 40.3, 125.8, 126.7,
10
11 127.8, 127.9, 128.5, 128.8, 129.3, 130.5, 130.9, 131.2, 132.0, 132.8, 133.3, 138.2, 140.1, 142.0,
12
13 195.8; FTIR (KBr) 3056, 1644, 1586, 1559, 1490, 758 cm^{-1} ; HRMS (m/z) calculated for
14
15 $\text{C}_{22}\text{H}_{17}\text{OS}$ [$\text{M}+\text{H}$] $^+$: 329.1000; found: 329.1018.
16
17
18
19

20
21 [2-(*p*-Tolyl)-2H-thiochromen-3-yl](phenyl)methanone (3b). 139 mg, 81% yield; pale yellow
22
23 solid; mp 117-119 °C; R_f 0.41 (5% ethyl acetate in hexanes); ^1H NMR (CDCl_3 , 400 MHz) δ 2.17
24
25 (s, 3H), 5.36 (s, 1H), 6.93 (d, $J = 6.4$ Hz, 2H), 7.04 (td, $J = 6.2$ Hz, 1.2 Hz, 1H), 7.08 (d, $J = 6.4$
26
27 Hz, 2H), 7.13-7.18 (m, 3H), 7.30 (s, 1H), 7.38 (t, $J = 6.4$ Hz, 2H), 7.48 (tt, $J = 6.0$ Hz, 1.6 Hz,
28
29 1H), 7.62 (d, $J = 6.4$ Hz, 2H); ^{13}C NMR (CDCl_3 , 100 MHz) δ 21.2, 40.1, 125.7, 126.6, 127.8,
30
31 128.5, 129.3, 129.5, 130.5, 130.9, 131.1, 132.0, 132.8, 133.5, 137.5, 138.2, 139.1, 140.0, 195.9;
32
33 FTIR (KBr) 3054, 1641, 1507, 1441, 754 cm^{-1} ; HRMS (m/z) calculated for $\text{C}_{23}\text{H}_{19}\text{OS}$ [$\text{M}+\text{H}$] $^+$:
34
35 343.1157; found: 343.1135.
36
37
38
39

40
41 [2-(4-(Methylthio)phenyl)-2H-thiochromen-3-yl](phenyl)methanone (3c). 162 mg, 86% yield;
42
43 yellowish viscous liquid; R_f 0.39 (10% ethyl acetate in hexanes); ^1H NMR (CDCl_3 , 400 MHz) δ
44
45 2.39 (s, 3H), 5.42 (s, 1H), 7.08 (d, $J = 8.4$ Hz, 2H), 7.13 (dd, $J = 6.6$ Hz, 2.4 Hz, 1H), 7.19 (d, $J =$
46
47 8.4 Hz, 2H), 7.22-7.28 (m, 3H), 7.38 (s, 1H), 7.46 (t, $J = 8.0$ Hz, 2H), 7.56 (t, $J = 7.2$ Hz, 1H),
48
49 7.69 (d, $J = 7.2$ Hz, 2H); ^{13}C NMR (CDCl_3 , 100 MHz) δ 15.8, 39.9, 125.8, 126.8, 127.2, 127.8,
50
51 128.5, 129.3, 130.5, 130.9, 131.2, 132.0, 132.6, 133.2, 138.0, 138.1, 138.8, 140.1, 195.8; FTIR
52
53 (KBr) 3059, 1642, 1583, 1486, 1556, 754 cm^{-1} ; HRMS (m/z) calculated for $\text{C}_{23}\text{H}_{18}\text{ONaS}_2$
54
55 [$\text{M}+\text{Na}$] $^+$: 397.0698; found: 397.0691.
56
57
58
59
60

[2-(3-Methoxyphenyl)-2H-thiochromen-3-yl](phenyl)methanone (3d). 164 mg, 91% yield; yellow solid; mp 147-149 °C; R_f 0.36 (10% ethyl acetate in hexanes); ^1H NMR (CDCl_3 , 400 MHz) δ 3.68 (s, 3H), 5.45 (s, 1H), 6.71 (dd, $J = 8.4$ Hz, 2.4 Hz, 1H), 6.82 (s, 1H), 6.87 (d, $J = 8.0$ Hz, 1H), 7.08-7.15 (m, 2H), 7.19-7.27 (m, 3H), 7.38 (s, 1H), 7.46 (t, $J = 7.6$ Hz, 2H), 7.55 (t, $J = 7.6$ Hz, 1H), 7.69 (d, $J = 7.2$ Hz, 2H); ^{13}C NMR (CDCl_3 , 100 MHz) δ 40.2, 55.2, 112.6, 113.1, 119.1, 125.8, 127.7, 128.5, 129.3, 129.8, 130.5, 130.9, 131.2, 132.0, 132.8, 133.3, 138.2, 140.2, 143.5, 159.8, 195.8; FTIR (KBr) 3056, 2834, 1641, 1600, 1556, 1487, 755 cm^{-1} ; HRMS (m/z) calculated for $\text{C}_{23}\text{H}_{19}\text{O}_2\text{S}$ $[\text{M}+\text{H}]^+$: 359.1106; found: 359.1125

[2-(3,4,5-Trimethoxyphenyl)-2H-thiochromen-3-yl](phenyl)methanone (3e). 157 mg, 75% yield; brownish viscous liquid; R_f 0.54 (30% ethyl acetate in hexanes); ^1H NMR (CDCl_3 , 400 MHz) δ 3.67 (s, 6H), 3.76 (s, 3H), 5.40 (s, 1H), 6.51 (s, 2H), 7.12-7.17 (m, 1H), 7.26-7.29 (m, 2H), 7.32 (d, $J = 7.2$ Hz, 1H), 7.39 (s, 1H), 7.48 (t, $J = 8.0$ Hz, 2H), 7.58 (tt, $J = 7.6$ Hz, 1.6 Hz, 1H), 7.71 (dd, $J = 8.4$ Hz, 1.6 Hz, 2H); ^{13}C NMR (CDCl_3 , 100 MHz) δ 40.6, 56.0, 60.9, 103.7, 125.9, 127.9, 128.6, 129.3, 130.6, 131.2, 132.1, 132.9, 133.8, 137.5, 137.6, 138.1, 139.9, 153.2, 195.9; FTIR (KBr) 3051, 2862, 1647, 1592, 1541, 1436, 751 cm^{-1} ; HRMS (m/z) calculated for $\text{C}_{25}\text{H}_{22}\text{O}_4\text{NaS}$ $[\text{M}+\text{Na}]^+$: 441.1137; found: 441.1153.

[2-(4-Chlorophenyl)-2H-thiochromen-3-yl](phenyl)methanone (3f): 151 mg, 83% yield; yellow viscous liquid; R_f 0.65 (10% ethyl acetate in hexanes); ^1H NMR (CDCl_3 , 400 MHz) δ 5.41 (s, 1H), 7.13-7.23 (m, 5H), 7.24-7.27 (m, 3H), 7.41 (s, 1H), 7.47 (t, $J = 7.6$ Hz, 2H), 7.57 (tt, $J = 7.6$ Hz, 2.0 Hz, 1H), 7.69 (d, $J = 7.2$ Hz, 2H); ^{13}C NMR (CDCl_3 , 100 MHz) δ 39.8, 126.0, 127.8, 128.1, 128.6, 128.9, 129.3, 130.4, 131.0, 131.4, 132.1, 132.4, 133.0, 133.6, 138.0, 140.4, 140.5, 195.7; FTIR (KBr) 3038, 1651, 1613, 1578, 1476, 818, 757 cm^{-1} ; HRMS (m/z) calculated for $\text{C}_{22}\text{H}_{15}\text{OClNaS}$ $[\text{M}+\text{Na}]^+$: 385.0424; found: 385.0432.

[2-(2-Bromophenyl)-2H-thiochromen-3-yl](phenyl)methanone (3g). 155 mg, 76% yield; pale yellow solid; mp 148-150 °C; R_f 0.38 (10% ethyl acetate in hexanes); ^1H NMR (CDCl_3 , 400 MHz) δ 5.86 (s, 1H), 7.00-7.08 (m, 3H), 7.11-7.23 (m, 3H), 7.29 (d, $J = 7.2$ Hz, 1H), 7.48 (t, $J = 7.2$ Hz, 2H), 7.54-7.62 (m, 3H), 7.72 (d, $J = 7.2$ Hz, 2H); ^{13}C NMR (CDCl_3 , 100 MHz) δ 40.0, 122.9, 126.0, 127.6, 127.7, 128.2, 128.6, 129.1, 129.3, 130.5, 131.0, 131.4, 132.1, 132.2, 132.5, 133.8, 137.9, 139.6, 141.8, 195.2; FTIR (KBr) 3059, 1641, 1585, 1556, 1461, 1109, 754 cm^{-1} ; HRMS (m/z) calculated for $\text{C}_{22}\text{H}_{15}\text{OBrNaS}$ [$\text{M}+\text{Na}$] $^+$: 428.9919; found: 428.9929.

[2-(2-Fluorophenyl)-2H-thiochromen-3-yl](phenyl)methanone (3h). 137 mg, 79% yield; yellowish viscous liquid; R_f 0.41 (10% ethyl acetate in hexanes); ^1H NMR (CDCl_3 , 400 MHz) δ 5.77 (s, 1H), 6.88 (t, $J = 7.6$ Hz, 1H), 7.00-7.07 (m, 2H), 7.10-7.16 (m, 2H), 7.18-7.23 (m, 2H), 7.27 (d, $J = 7.0$ Hz, 1H), 7.43-7.52 (m, 3H), 7.56 (t, $J = 7.6$ Hz, 1H), 7.72 (d, $J = 6.8$ Hz, 2H); ^{13}C NMR (CDCl_3 , 100 MHz) δ 33.3 (d, $J = 3.9$ Hz), 116.0 (d, $J = 21.6$ Hz), 124.1 (d, $J = 3.7$ Hz), 125.9, 127.7 (d, $J = 3.0$ Hz), 127.9, 128.5, 128.6 (d, $J = 11.5$ Hz), 129.3 (d, $J = 7.8$ Hz), 129.4, 130.4, 130.9, 131.3, 131.7, 132.1, 132.6, 137.9, 141.3, 159.1 (d, $J = 246.9$ Hz), 195.2; FTIR (KBr) 3056, 1641, 1598, 1583, 1436, 1264, 754 cm^{-1} ; HRMS (m/z) calculated for $\text{C}_{22}\text{H}_{15}\text{OFNaS}$ [$\text{M}+\text{Na}$] $^+$: 369.0725; found: 369.0722.

[2-(4-(Trifluoromethyl)phenyl)-2H-thiochromen-3-yl](phenyl)methanone (3i). 140 mg, 82% yield; yellowish viscous liquid; R_f 0.46 (10% ethyl acetate in hexanes); ^1H NMR (CDCl_3 , 400 MHz) δ 5.39 (s, 1H), 7.06-7.12 (m, 1H), 7.18-7.21 (m, 3H), 7.32 (d, $J = 8.4$ Hz, 2H), 7.37-7.44 (m, 5H), 7.51 (tt, $J = 7.6$ Hz, 2.4 Hz, 1H), 7.63 (dd, $J = 8.2$ Hz, 1.6 Hz, 2H); ^{13}C NMR (CDCl_3 , 100 MHz) δ 40.0, 124.1 (q, $J = 270.2$ Hz), 125.7 (q, $J = 3.7$ Hz), 126.1, 127.1, 127.8, 128.6, 129.3, 129.8 (q, $J = 32.2$ Hz), 130.4, 131.1, 131.5, 132.16, 132.19, 132.6, 137.9, 140.8, 145.8,

1
2
3 195.5; FTIR (KBr) 3064, 1641, 1617, 1597, 1446, 1325, 1120, 754 cm^{-1} ; HRMS (m/z) calculated
4 for $\text{C}_{23}\text{H}_{15}\text{OSNaF}_3$ $[\text{M}+\text{Na}]^+$: 419.0693; found: 419.0705.
5
6
7

8
9 [2-(Naphthalen-1-yl)-2H-thiochromen-3-yl](phenyl)methanone (3j). 178 mg, 94% yield; yellow
10 viscous liquid; R_f 0.40 (5% ethyl acetate in hexanes); ^1H NMR (CDCl_3 , 400 MHz) δ 6.34 (s,
11 1H), 7.09-7.25 (m, 5H), 7.34 (dd, $J = 7.2$ Hz, 1.6 Hz, 1H), 7.46-7.61 (m, 4H), 7.62-7.68 (m, 2H),
12 7.70 (d, $J = 7.0$ Hz, 1H), 7.77 (d, $J = 7.6$ Hz, 2H), 7.85 (d, $J = 8.4$ Hz, 1H), 8.33 (d, $J = 8.4$ Hz,
13 1H); ^{13}C NMR (CDCl_3 , 100 MHz) δ 35.8, 123.4, 124.4, 125.2, 125.88, 125.93, 126.7, 128.2,
14 128.5, 128.6, 129.2, 129.4, 129.7, 130.4, 131.1, 131.2, 132.0, 132.8, 133.2, 134.7, 135.7, 138.2,
15 141.6, 195.5; FTIR (KBr) 3056, 1641, 1594, 1554, 1441, 754 cm^{-1} ; HRMS (m/z) calculated for
16 $\text{C}_{26}\text{H}_{18}\text{ONaS}$ $[\text{M}+\text{Na}]^+$: 401.0971; found: 401.0973.
17
18
19
20
21
22
23
24
25
26
27

28
29 [2-(Thiophen-2-yl)-2H-thiochromen-3-yl](phenyl)methanone (3k). 125 mg, 74% yield;
30 brownish viscous liquid; R_f 0.35 (10% ethyl acetate in hexanes); ^1H NMR (CDCl_3 , 400 MHz) δ
31 5.76 (s, 1H), 6.81 (t, $J = 3.6$ Hz, 1H), 6.91 (d, $J = 2.4$ Hz, 1H), 7.04 (d, $J = 5.2$ Hz, 1H), 7.15 (t, J
32 = 7.6 Hz, 1H), 7.24-7.30 (m, 2H), 7.31-7.36 (m, 2H), 7.48 (t, $J = 7.6$ Hz, 2H), 7.57 (t, $J = 7.2$ Hz,
33 1H), 7.73 (d, $J = 7.2$ Hz, 2H); ^{13}C NMR (CDCl_3 , 100 MHz) δ 35.7, 124.8, 125.0, 126.0, 126.8,
34 128.1, 128.5, 129.4, 130.4, 131.1, 131.2, 132.1, 132.5, 133.8, 138.1, 139.8, 145.4, 195.4; FTIR
35 (KBr) 3059, 1640, 1590, 1553, 1445, 754 cm^{-1} ; HRMS (m/z) calculated for $\text{C}_{20}\text{H}_{14}\text{ONaS}_2$
36 $[\text{M}+\text{Na}]^+$: 357.0378; found: 357.0381.
37
38
39
40
41
42
43
44
45
46
47

48
49 [2-(Pyridin-2-yl)-2H-thiochromen-3-yl](phenyl)methanone (3l). 148 mg, 90% yield; brownish
50 viscous liquid; R_f 0.41 (20% ethyl acetate in hexanes); ^1H NMR (CDCl_3 , 400 MHz) δ 5.57 (s,
51 1H), 7.05-7.13 (m, 2H), 7.19 (dd, $J = 7.6$ Hz, 1.6 Hz, 1H), 7.22-7.28 (m, 3H), 7.43 (s, 1H), 7.45-
52 7.54 (m, 3H), 7.57 (tt, $J = 7.2$ Hz, 2.0 Hz, 1H), 8.48 (ddd, $J = 4.8$ Hz, 1.6 Hz, 0.8 Hz, 1H), 7.79
53
54
55
56
57
58
59
60

1
2
3 (dd, $J = 8.2$ Hz, 1.6 Hz, 2H); ^{13}C NMR (CDCl_3 , 100 MHz) δ 42.4, 121.0, 122.3, 125.8, 127.6,
4
5 128.5, 129.6, 130.7, 130.9, 131.0, 132.1, 132.5, 132.9, 136.8, 138.1, 140.6, 149.7, 160.2, 195.9;
6
7 FTIR (KBr) 3056, 1641, 1585, 1465, 751 cm^{-1} ; HRMS (m/z) calculated for $\text{C}_{21}\text{H}_{15}\text{ONNaS}$
8
9 $[\text{M}+\text{Na}]^+$: 352.0767; found: 352.0772.
10
11

12
13 (2-Phenyl-2H-thiochromen-3-yl)(p-tolyl)methanone (3m). 147 mg, 86% yield; yellow viscous
14
15 liquid; R_f 0.36 (5% ethyl acetate in hexanes); ^1H NMR (CDCl_3 , 400 MHz) δ 2.43 (s, 3H), 5.46
16
17 (s, 1H), 7.12 (td, $J = 6.0$ Hz, 1.2 Hz, 1H), 7.16-7.22 (m, 3H), 7.23-7.29 (m, 7H), 7.38 (s, 1H),
18
19 7.62 (d, $J = 6.8$ Hz, 2H); ^{13}C NMR (CDCl_3 , 100 MHz) δ 21.7, 40.5, 125.8, 126.7, 127.72,
20
21 127.74, 128.8, 129.2, 129.6, 130.6, 130.8, 131.0, 132.7, 133.5, 135.4, 139.5, 142.1, 142.8, 195.6;
22
23 FTIR (KBr) 3046, 1642, 1596, 1543, 1472, 753 cm^{-1} ; HRMS (m/z) calculated for $\text{C}_{23}\text{H}_{18}\text{ONaS}$
24
25 $[\text{M}+\text{Na}]^+$: 365.0976; found: 365.0965.
26
27
28
29

30
31 (2-Phenyl-2H-thiochromen-3-yl)(3-bromophenyl)methanone (3n). 159 mg, 78% yield; brownish
32
33 viscous liquid; R_f 0.45 (10% ethyl acetate in hexanes); ^1H NMR (CDCl_3 , 400 MHz) δ 5.43 (s,
34
35 1H), 7.11-7.16 (m, 1H), 7.17-7.22 (m, 3H), 7.23-7.28 (m, 5H), 7.33 (t, $J = 6.0$ Hz, 1H), 7.38 (s,
36
37 1H), 7.58 (d, $J = 6.0$ Hz, 1H), 7.67 (d, $J = 6.4$ Hz, 1H), 7.80 (s, 1H); ^{13}C NMR (CDCl_3 , 100
38
39 MHz) δ 40.2, 122.8, 125.9, 126.7, 127.7, 127.8, 127.9, 128.8, 130.0, 130.3, 131.1, 131.5, 132.0,
40
41 132.9, 133.0, 134.8, 140.1, 140.8, 141.8, 194.2; FTIR (KBr) 3059, 1643, 1617, 1557, 1436,
42
43 1112, 758 cm^{-1} ; HRMS (m/z) calculated for $\text{C}_{22}\text{H}_{16}\text{OBrS}$ $[\text{M}+\text{H}]^+$: 407.0100; found: 407.0103.
44
45
46
47
48

49 (2-Phenyl-2H-thiochromen-3-yl)(4-chlorophenyl)methanone (3o). 150 mg, 82% yield; yellow
50
51 solid; mp 144-146 $^\circ\text{C}$; R_f 0.48 (5% ethyl acetate in hexanes); ^1H NMR (CDCl_3 , 400 MHz) δ 5.43
52
53 (s, 1H), 7.13 (td, $J = 8.4$ Hz, 2.4 Hz, 1H), 7.16-7.23 (m, 3H), 7.23-7.28 (m, 5H), 7.35 (s, 1H),
54
55 7.44 (d, $J = 8.4$ Hz, 2H), 7.64 (d, $J = 8.4$ Hz, 2H); ^{13}C NMR (CDCl_3 , 100 MHz) δ 40.3, 125.9,
56
57
58
59
60

1
2
3 126.7, 127.8, 127.9, 128.8, 130.3, 130.7, 131.0, 131.4, 132.9, 133.1, 136.5, 138.4, 140.1, 141.9,
4
5 194.6; FTIR (KBr) 3060, 1648, 1616, 1586, 1487, 828, 761 cm^{-1} ; HRMS (m/z) calculated for
6
7 $\text{C}_{22}\text{H}_{15}\text{OClNaS}$ [$\text{M}+\text{Na}$] $^{+}$: 385.0424; found: 385.0429.
8
9

10
11 (2-Phenyl-2H-thiochromen-3-yl)(4-fluorophenyl)methanone (3p). 129 mg, 74% yield; yellow
12
13 viscous liquid; R_f 0.45 (5% ethyl acetate in hexanes); ^1H NMR (CDCl_3 , 400 MHz) δ 5.35 (s,
14
15 1H), 7.01-7.07 (m, 3H), 7.07-7.13 (m, 3H), 7.14-7.20 (m, 5H), 7.26 (s, 1H), 7.61-7.67 (m, 2H);
16
17 ^{13}C NMR (CDCl_3 , 100 MHz) δ 40.5, 115.7 (d, $J = 21.7$ Hz), 125.8, 126.7, 127.7, 127.8, 128.8,
18
19 130.4, 130.9, 131.2, 131.8 (d, $J = 9.0$ Hz), 132.8, 133.2, 134.3 (d, $J = 3.3$ Hz), 139.7, 141.9,
20
21 165.1 (d, $J = 251.8$ Hz), 194.4; FTIR (KBr) 3054, 1643, 1560, 1421, 751 cm^{-1} ; HRMS (m/z)
22
23 calculated for $\text{C}_{22}\text{H}_{15}\text{OFNaS}$ [$\text{M}+\text{Na}$] $^{+}$: 369.0725; found: 369.0727.
24
25
26
27

28
29 (2-Phenyl-2H-thiochromen-3-yl)[4-(trifluoromethyl)phenyl]methanone (3q). 132 mg, 66%
30
31 yield; yellow viscous liquid; R_f 0.48 (10% ethyl acetate in hexanes); ^1H NMR (CDCl_3 , 400
32
33 MHz) δ 5.39 (s, 1H), 7.05-7.10 (m, 1H), 7.11-7.17 (m, 3H), 7.18-7.21 (m, 5H), 7.28 (s, 1H),
34
35 7.64-7.72 (m, 4H); ^{13}C NMR (CDCl_3 , 125 MHz) δ 40.0, 123.7 (q, $J = 270.9$ Hz), 125.5 (q, $J =$
36
37 3.6 Hz), 125.9, 126.6, 127.7, 127.9, 128.8, 129.4, 130.1, 131.1, 131.6, 132.8, 132.9, 133.2 (q, $J =$
38
39 32.5 Hz), 141.1, 141.4, 141.8, 194.5; FTIR (KBr) 3058, 1647, 1617, 1585, 1406, 1324, 1129,
40
41 752 cm^{-1} ; MS (m/z): HRMS (m/z) calculated for $\text{C}_{23}\text{H}_{15}\text{OF}_3\text{NaS}$ [$\text{M}+\text{Na}$] $^{+}$: 419.0688; found:
42
43 419.0693.
44
45
46
47

48
49 (2-Phenyl-2H-thiochromen-3-yl)(naphthalen-2-yl)methanone (3r). 160 mg, 84% yield; yellow
50
51 viscous liquid; R_f 0.70 (10% ethyl acetate in hexanes); ^1H NMR (CDCl_3 , 400 MHz) δ 5.51 (s,
52
53 1H), 7.08 (t, $J = 7.2$ Hz, 1H), 7.15-7.27 (m, 6H), 7.32 (d, $J = 7.2$ Hz, 2H), 7.45 (s, 1H), 7.49-
54
55 7.59 (m, 2H), 7.76 (d, $J = 8.4$ Hz, 1H), 7.84-7.92 (m, 3H), 8.18 (s, 1H); ^{13}C NMR (CDCl_3 , 100
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

MHz) δ 40.5, 125.6, 125.8, 126.7, 126.9, 127.7, 127.8, 127.9, 128.2, 128.5, 128.8, 129.3, 130.2, 130.5, 130.9, 131.1, 132.4, 132.8, 133.5, 135.0, 135.4, 140.1, 142.0, 195.7; FTIR (KBr) 3056, 1640, 1584, 1555, 1490, 758 cm^{-1} ; HRMS (m/z) calculated for $\text{C}_{26}\text{H}_{19}\text{OS}$ $[\text{M}+\text{H}]^+$: 379.1157; found: 379.1164.

(2-Phenyl-2H-thiochromen-3-yl)(thiophen-2-yl)methanone (3s). 115 mg, 68% yield; yellow viscous liquid; R_f 0.40 (10% ethyl acetate in hexanes); ^1H NMR (CDCl_3 , 400 MHz) δ 5.38 (s, 1H), 7.12-7.20 (m, 5H), 7.22-7.29 (m, 4H), 7.33 (d, $J = 7.6$ Hz, 1H), 7.65 (dd, $J = 5.0$ Hz, 1.2 Hz, 1H), 7.68 (s, 1H), 7.71 (dd, $J = 3.6$ Hz, 1.2 Hz, 1H); ^{13}C NMR (CDCl_3 , 100 MHz) δ 40.9, 125.9, 126.8, 127.8, 127.9, 128.8, 130.5, 130.7, 131.0, 132.6, 133.4, 133.6, 133.7, 137.8, 141.8, 143.1, 186.7; FTIR (KBr) 3058, 1617, 1584, 1553, 1411, 759 cm^{-1} ; HRMS (m/z) calculated for $\text{C}_{20}\text{H}_{15}\text{OS}_2$ $[\text{M}+\text{H}]^+$: 335.0564; found: 335.0557.

1-(2-Phenyl-2H-thiochromen-3-yl)ethan-1-one (3t). 112 mg, 84% yield; yellow viscous liquid; R_f 0.42 (10% ethyl acetate in hexanes); ^1H NMR (CDCl_3 , 400 MHz) δ 2.48 (s, 3H), 5.36 (s, 1H), 7.14-7.20 (m, 6H), 7.23-7.26 (m, 2H), 7.37 (d, $J = 7.2$ Hz, 1H), 7.67 (s, 1H); ^{13}C NMR (CDCl_3 , 100 MHz) δ 25.7, 38.9, 125.8, 126.7, 127.7, 127.8, 128.7, 130.6, 130.9, 131.2, 133.0, 134.1, 137.5, 142.1, 196.6; FTIR (KBr) 3058, 2924, 1659, 1622, 1584, 1438, 754 cm^{-1} ; HRMS (m/z) calculated for $\text{C}_{17}\text{H}_{14}\text{ONaS}$ $[\text{M}+\text{Na}]^+$: 289.0663; found: 289.0687.

2-Phenyl-2H-thiochromene-3-carbaldehyde (3u).^{8a} 123 mg, 97% yield; pale yellow viscous liquid; R_f 0.44 (10% ethyl acetate in hexanes); ^1H NMR (CDCl_3 , 400 MHz) δ 5.21 (s, 1H), 7.16-7.22 (m, 6H), 7.27 (t, $J = 7.2$ Hz, 2H), 7.41 (d, $J = 7.6$ Hz, 1H), 7.46 (s, 1H), 9.67 (s, 1H); ^{13}C NMR (CDCl_3 , 100 MHz) δ 38.5, 126.0, 126.6, 127.8, 128.0, 128.8, 130.0, 131.1, 131.9, 134.3,

1
2
3 135.0, 141.8, 145.3, 191.3; FTIR (KBr) 3058, 1668, 1583, 1554, 1437, 751 cm^{-1} ; HRMS (m/z)
4
5 calculated for $\text{C}_{16}\text{H}_{12}\text{OKS}$ $[\text{M}+\text{K}]^+$: 291.0246; found: 291.0245.
6
7

8
9 2-Phenyl-2H-thiochromene-3-carbaldehyde (3v). 131 mg, 98% yield; yellow viscous liquid; R_f
10 0.47 (10% ethyl acetate in hexanes); ^1H NMR (CDCl_3 , 400 MHz) δ 2.16 (s, 3H), 5.10 (s, 1H),
11 6.92 (d, $J = 7.6$ Hz, 2H), 7.01 (d, $J = 8.4$ Hz, 2H), 7.08-7.13 (m, 1H), 7.16-7.21 (m, 2H), 7.33 (d,
12 $J = 7.2$ Hz, 1H), 7.37 (s, 1H), 9.58 (s, 1H); ^{13}C NMR (CDCl_3 , 100 MHz) δ 21.2, 38.3, 125.9,
13 126.5, 127.8, 129.5, 130.0, 131.1, 131.8, 134.3, 135.0, 137.8, 138.9, 145.2, 191.4; FTIR (KBr)
14 3056, 2934, 1665, 1593, 1564, 1427, 753 cm^{-1} ; HRMS (m/z) calculated for $\text{C}_{17}\text{H}_{14}\text{ONaS}$
15
16 $[\text{M}+\text{Na}]^+$: 289.0663; found: 289.0670.
17
18
19
20
21
22
23
24
25

26 2-Methyl-2H-thiochromene-3-carbaldehyde (3w). 77 mg, 81% yield; yellow viscous liquid; R_f
27 0.50 (10% ethyl acetate in hexanes); ^1H NMR (CDCl_3 , 400 MHz) δ 1.29 (d, $J = 7.2$ Hz, 3H),
28 4.08 (q, $J = 7.2$ Hz, 1H), 7.18 (td, $J = 7.4$ Hz, 1.2 Hz, 1H), 7.23 (s, 1H), 7.30 (td, $J = 7.2$ Hz, 1.6
29 Hz, 1H), 7.33-7.37 (m, 2H), 9.61 (s, 1H); ^{13}C NMR (CDCl_3 , 100 MHz) δ 22.3, 30.4, 125.8,
30 128.4, 130.0, 131.0, 131.5, 134.2, 138.0, 144.3, 191.3; FTIR (KBr) 3057, 2813, 1673, 1584,
31 1556, 1439, 751 cm^{-1} ; HRMS (m/z) calculated for $\text{C}_{11}\text{H}_{10}\text{ONaS}$ $[\text{M}+\text{Na}]^+$: 213.0350; found:
32 213.0348.
33
34
35
36
37
38
39
40
41
42
43

44 (6-Bromo-2-phenyl-2H-thiochromen-3-yl)(phenyl)methanone (3x). 176 mg, 86% yield; yellow
45 viscous liquid; R_f 0.58 (10% ethyl acetate in hexanes); ^1H NMR (CDCl_3 , 400 MHz) δ 5.44 (s,
46 1H), 7.09 (d, $J = 8.4$ Hz, 1H), 7.15-7.23 (m, 5H), 7.26 (s, 1H), 7.29 (d, $J = 8.4$ Hz, 1H), 7.37 (s,
47 1H), 7.42 (t, $J = 7.6$ Hz, 2H), 7.53 (t, $J = 7.6$ Hz, 1H), 7.63 (d, $J = 8.0$ Hz, 2H); ^{13}C NMR
48 (CDCl₃, 100 MHz) δ 40.4, 118.8, 126.7, 128.0, 128.6, 128.8, 128.9, 129.1, 129.3, 131.8, 132.3,
49 133.1, 133.7, 134.5, 137.9, 138.4, 141.5, 195.6; FTIR (KBr) 3057, 1643, 1537, 1491, 1448,
50
51
52
53
54
55
56
57
58
59
60

1
2
3 1084, 728 cm^{-1} ; HRMS (m/z) calculated for $\text{C}_{22}\text{H}_{15}\text{OBrNaS}$ $[\text{M}+\text{Na}]^+$: 428.9925; found:
4
5 428.9945.
6
7

8
9 (6-Chloro-2-phenyl-2H-thiochromen-3-yl)(phenyl)methanone (3y). 146 mg, 80% yield; yellow
10 viscous liquid; R_f 0.32 (5% ethyl acetate in hexanes); ^1H NMR (CDCl_3 , 400 MHz) δ 5.48 (s,
11 1H), 7.18-7.29 (m, 8H), 7.31 (s, 1H), 7.46 (t, $J = 7.6$ Hz, 2H), 7.56 (tt, $J = 7.6$ Hz, 2.4 Hz, 1H),
12 7.68 (dd, $J = 8.2$ Hz, 1.6 Hz, 2H); ^{13}C NMR (CDCl_3 , 100 MHz) δ 40.4, 126.7, 128.0, 128.6,
13 128.8, 128.9, 129.3, 130.2, 130.8, 131.1, 131.2, 131.9, 132.3, 134.5, 137.8, 138.6, 141.4, 195.6;
14 FTIR (KBr) 3054, 1641, 1564, 1501, 1468, 821, 736 cm^{-1} ; HRMS (m/z) calculated for
15 $\text{C}_{22}\text{H}_{15}\text{OCINaS}$ $[\text{M}+\text{Na}]^+$: 385.0430; found: 385.0449.
16
17
18
19
20
21
22
23
24
25

26 (7-Methyl-2-phenyl-2H-thiochromen-3-yl)(phenyl)methanone (3z). 151 mg, 88% yield; yellow
27 viscous liquid; R_f 0.62 (10% ethyl acetate in hexanes); ^1H NMR (CDCl_3 , 400 MHz) δ 2.19 (s,
28 3H), 5.37 (s, 1H), 6.94-7.00 (m, 2H), 7.03-7.13 (m, 4H), 7.20 (d, $J = 7.2$ Hz, 2H), 7.28 (s, 1H),
29 7.37 (t, $J = 7.6$ Hz, 2H), 7.46 (t, $J = 6.8$ Hz, 1H), 7.60 (d, $J = 8.0$ Hz, 2H); ^{13}C NMR (CDCl_3 , 100
30 MHz) δ 20.9, 40.3, 126.7, 127.6, 127.7, 128.4, 128.7, 129.2, 129.3, 130.4, 131.5, 131.9, 132.16,
31 133.4, 135.5, 138.2, 140.5, 142.0, 195.9; FTIR (KBr) 3052, 2920, 1673, 1584, 1555, 1438, 751
32 cm^{-1} ; HRMS (m/z) calculated for $\text{C}_{23}\text{H}_{19}\text{OS}$ $[\text{M}+\text{H}]^+$: 343.1157; found: 343.1129.
33
34
35
36
37
38
39
40
41
42
43
44
45
46

47 [2-(4-Fluorophenyl)-2H-thiochromen-3-yl](phenyl)methanone (3aa). 122 mg, 70% yield; yellow
48 viscous liquid; R_f 0.50 (10% ethyl acetate in hexanes); ^1H NMR (CDCl_3 , 400 MHz) δ 5.36 (s,
49 1H), 6.80 (tt, $J = 10.8$ Hz, 2.4 Hz, 2H), 7.03-7.08 (m, 1H), 7.13-7.19 (m, 5H), 7.32 (s, 1H), 7.39
50 (t, $J = 7.2$ Hz, 2H), 7.48 (tt, $J = 7.2$ Hz, 1.2 Hz, 1H), 7.61 (dd, $J = 8.2$ Hz, 1.6 Hz, 2H); ^{13}C NMR
51 (CDCl_3 , 100 MHz) δ 39.7, 115.6 (d, $J = 21.4$ Hz), 125.9, 127.8, 128.4 (d, $J = 8.1$ Hz), 128.5,
52
53
54
55
56
57
58
59
60

1
2
3 129.3, 130.4, 131.0, 131.3, 132.1, 132.5, 133.3, 137.9 (d, $J = 3.2$ Hz), 138.1, 140.2, 162.3 (d, $J =$
4 244.7 Hz), 195.8; FTIR (KBr) 3053, 1648, 1598, 1564, 1423, 1198, 751 cm^{-1} ; HRMS (m/z)
5
6 calculated for $\text{C}_{22}\text{H}_{15}\text{OFNaS}$ $[\text{M}+\text{Na}]^+$: 369.0720; found: 369.0721.
7
8
9

10
11 [2-(6-Methoxynaphthalen-1-yl)-2H-thiochromen-3-yl](phenyl)methanone (3ab). 159 mg, 78%
12 yield; brownish viscous liquid; R_f 0.32 (10% ethyl acetate in hexanes); ^1H NMR (CDCl_3 , 400
13 MHz) δ 3.78 (s, 3H), 5.53 (s, 1H), 6.93-7.01 (m, 2H), 7.05 (td, $J = 6.8$ Hz, 1.6 Hz, 1H), 7.11-
14 7.17 (m, 2H), 7.21(d, $J = 7.6$ Hz, 1H), 7.31-7.41 (m, 4H), 7.44-7.55 (m, 4H), 7.63 (d, $J = 7.6$ Hz,
15 2H); ^{13}C NMR (CDCl_3 , 100 MHz) δ 40.5, 55.4, 105.7, 118.9, 125.2, 125.76, 125.79, 127.7,
16 127.8, 128.5, 128.7, 129.3, 129.7, 130.5, 131.0, 131.2, 132.0, 132.8, 133.2, 134.2, 136.9, 138.2,
17 140.2, 157.9, 195.9; FTIR (KBr) 3056, 2933, 1638, 1505, 1481, 1441, 754 cm^{-1} ; HRMS (m/z)
18 calculated for $\text{C}_{27}\text{H}_{20}\text{O}_2\text{NaS}$ $[\text{M}+\text{Na}]^+$: 431.1082; found: 431.1079.
19
20
21
22
23
24
25
26
27
28
29
30

31 (2-Phenyl-2H-thiochromen-3-yl)(benzo[d][1,3]dioxol-5-yl)methanone (3ac). 138 mg, 74%
32 yield; brownish viscous liquid; R_f 0.35 (10% ethyl acetate in hexanes); ^1H NMR (CDCl_3 , 400
33 MHz) δ 5.34 (s, 1H), 5.98 (q, $J = 0.8$ Hz, 2H), 6.80 (d, $J = 6.4$ Hz, 1H), 7.07 (td, $J = 5.8$ Hz, 1.2
34 Hz, 1H), 7.10-7.15 (m, 3H), 7.15-7.20 (m, 6H), 7.25 (dd, $J = 6.4$ Hz, 1.2 Hz, 1H), 7.29 (s, 1H);
35 ^{13}C NMR (CDCl_3 , 100 MHz) δ 40.8, 101.9, 108.0, 109.7, 125.5, 125.8, 126.7, 127.7, 127.8,
36 128.8, 130.5, 130.7, 131.0, 132.3, 132.6, 133.5, 138.7, 142.0, 148.0, 151.3, 194.2; FTIR (KBr)
37 3064, 2920, 1636, 1605, 1503, 1439, 761 cm^{-1} ; HRMS (m/z) calculated for $\text{C}_{23}\text{H}_{17}\text{O}_3\text{S}$ $[\text{M}+\text{H}]^+$:
38 373.0898; found: 373.0879.
39
40
41
42
43
44
45
46
47
48
49
50

51 ACKNOWLEDGMENT

52 We thank the DST New Delhi (Project No. SB/S1/OC-72/2013) for financial support. NS thanks
53 CSIR for fellowship.
54
55
56
57
58
59
60

SUPPORTING INFORMATION

¹H and ¹³C NMR spectra. Crystallography data.

REFERENCES

- (1) (a) *Sulphur-Containing Drugs and Related Organic Compounds*, Damani, L. A., Ed.; Wiley, New York, 1989. (b) *Sulfur Compounds: Advances in Research and Application*, Acton, A. Q., Ed.; Scholarly Editions, Atlanta, GA, 2012. (c) Nielsen, S. F. E.; Nielsen, O.; Olsen, G.M.; Liljefors, T.; Peters, D. *J. Med. Chem.* **2000**, *43*, 2217. (d) Smith, B. R.; Eastman, C. M.; Njardarson, J. T. *J. Med. Chem.* **2014**, *57*, 9764.
- (2) (a) Mishra, A.; Ma, C. Q.; Bäuerle, P. *Chem. Rev.* **2009**, *109*, 1141. (b) Takimiya, K.; Osaka, I.; Mori, T.; Nakano, M. *Acc. Chem. Res.* **2014**, *47*, 1493.
- (3) (a) Lin, D. Y.; Zhang, S. Z.; Block, E.; Katz, L. C. *Nature* **2005**, *434*, 470. (b) Joyce, N. I.; Eady, C. C.; Silcock, P.; Perry, N. B.; Van Klink, J. W. *J. Agric. Food Chem.* **2013**, *61*, 1449.
- (4) (a) Geissler, J. F.; Roesel, J. L.; Meyer, T.; Trinks, U. P.; Traxler, P.; Lydon, N. B. *Cancer Res.* **1992**, *52*, 4492. (b) Sugita, Y.; Hosoya, H.; Terasawa, K.; Yokoe, I.; Fujisawa, S.; Sakagami, H.; *Anticancer Res.* **2001**, *21*, 2629. (c) Brown, M. J.; Carter, P. S.; Fenwick, A. E.; Fosberry, A. P.; Hamprecht, D. W.; Hibbs, M. J.; Jarvest, R. L.; Mensah, L.; Milner, P. H.; O'Hanlon, P. J.; Pope, A. J.; Richardson, C. M.; West, A.; Witty, D. R. *Bioorg. Med. Chem. Lett.* **2002**, *12*, 3171. (d) Quaglia, W.; Pignini, M.; Piergentili, A.; Giannella, M.; Gentili, F.; Marucci, G.; Carrieri, A.; Carotti, A.; Poggesi, E.; Leonardi, A.; Melchiorre, C. *J. Med. Chem.* **2002**, *45*, 1633. (e) Zimmer, L.; Fournet, G.; Joseph, B.; Guillaumet, G.; Bars, D. L. *Nucl. Med. Biol.* **2003**, *30*, 541. (f) Kaye, P. T.; Musa, M. A.; Nchinda, A. T.; Nocanda, X. W. *Synth. Commun.* **2004**, *34*, 2575. (g) Yoneya, T.; Taniguchi, K.; Nakamura, R.; Tsunenari, T.; Ohizumi, I.; Kanbe, Y.; Morikawa, K.; Kaiho, S.-I.; Yamada-Okabe, H. *Anticancer Res.* **2010**, *30*, 873.

- 1
2
3 (5) (a) Camps, F.; Colomina, O.; Coll, J.; Messeguer, A. *J. Heterocyclic chem.*, **1983**, *20*, 1115.
4
5 (b) Tercio, J.; Ferreira, B.; Catani, V.; Comasseto, J. V. *Synthesis* **1987**, 149.
6
7
8 (6) (a) Arnoldi, A.; Carughi, M. *Synthesis* **1988**, 155. (b) Takido, T.; Takagi, Y.; Itabashi, K. *J.*
9
10 *Heterocycl. Chem.* **1995**, *32*, 687. (c) Kobayashi, K.; Nakahashi, R.; Shimizu, A.; Kitamura, T.;
11
12 Morikawa, O.; Konishi, H. *J. Chem. Soc., Perkin Trans. 1*, **1999**, 1547. (d) Kaye, P. T.; Nocanda,
13
14 X. W. *Synthesis* **2001**, 2389. (e) Ali, A.; Ahmad, V. U.; Liebscher, J. *Eur. J. Org. Chem.* **2001**,
15
16 529. (f) Pal, S. C.; Bhunia, S. C. *Eur. J. Org. Chem.* **2013**, 4816. (g) Roy, R.; Rakshit, S.;
17
18 Bhowmik, T.; Khan, S.; Ghatak, A.; Bhar, S. *J. Org. Chem.* **2014**, *79*, 6603.
19
20
21 (7) Selected examples; (a) Zu, L.; Wang, J.; Li, H.; Xie, H.; Jiang, W.; Wang, W. *J. Am. Chem.*
22
23 *Soc.* **2007**, *129*, 1036. (b) Gao, Y.; Ren, Q.; Wu, H.; Li, M.; Wang, J. *Chem. Commun.* **2010**,
24
25 9232. (c) Wu, L.; Wang, Y.; Song, H.; Tang, L.; Zhou, Z.; Tang, C. *Adv. Synth. Catal.* **2013**, *355*,
26
27 1053. (d) Arai, T.; Yamamoto, Y. *Org. Lett.* **2014**, *16*, 1700. (e) Chauhan, P.; Mahajan, S.; Enders,
28
29 D. *Chem. Rev.* **2014**, *114*, 8807.
30
31
32
33 (8) (a) Wang, W.; Li, H.; Wang, J.; Zu, L. *J. Am. Chem. Soc.* **2006**, *128*, 10354. (b) Rios, R.;
34
35 Sunden, H.; Ibrahim, I.; Zhao, G.; Cordova, A. *Tetrahedron Lett.* **2006**, *47*, 8679. (c)
36
37 Choudhury, A. R.; Mukherjee, S. *Adv. Synth. Catal.* **2013**, *355*, 1989.
38
39
40 (9) (a) Kondo, T.; Mitsudo, T. A. *Chem. Rev.* **2000**, *100*, 3205. (b) Lyons, T.; Sanford, W. *Chem.*
41
42 *Rev.* **2010**, *110*, 1147. (c) Beletskaya, I. P.; Ananikov, V. P. *Chem. Rev.* **2011**, *111*, 1596.
43
44
45 (10) Selected examples; (a) Hou, C.; He, Q.; Yang, C. *Org. Lett.* **2014**, *16*, 5040. (b) Qiao, Z.;
46
47 Wei, J.; Jiang, X. *Org. Lett.* **2014**, *16*, 1212. (c) Zhang, X.; Zeng, W.; Yang, Y.; Huang, H.; Liang,
48
49 Y. *Org. Lett.* **2014**, *16*, 876. (d) Li, Y.; Pu, J.; Jiang, X. *Org. Lett.* **2014**, *16*, 2692. (e) Dang, P.;
50
51 Zeng, W.; Liang, Y. *Org. Lett.* **2015**, *17*, 34. (f) Zhang, Y.; Li, Y.; Zhang, X.; Jiang, X. *Chem.*
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Commun. **2015**, *51*, 941. (g) Liu, Y.; Zhang, J. L.; Song, R. J.; Li, J. H. *Org. Lett.* **2014**, *16*, 5838.

(h) Reddy, A. S.; Swamy, K. C. K. *Org. Lett.* **2015**, *17*, 2996.

(11) (a) Prasad, D. J. C.; Sekar, G. *Org. Lett.* **2011**, *13*, 1008. (b) Prasad, D. J. C.; Sekar, G. *Org. Biomol. Chem.* **2013**, *11*, 1659. (c) Sangeetha, S.; Muthupandi, P.; Sekar, G. *Org. Lett.* **2015**, *17*, 6006.