

Silver-Catalyzed Late-Stage Fluorination

Pingping Tang, Takeru Furuya, and Tobias Ritter*


Department of Chemistry and Chemical Biology, Harvard University, 12 Oxford Street,
Cambridge, Massachusetts 02138

Received July 1, 2010; E-mail: ritter@chemistry.harvard.edu

Abstract: Carbon–fluorine bond formation by transition metal catalysis is difficult, and only a few methods for the synthesis of aryl fluorides have been developed. All reported transition-metal-catalyzed fluorination reactions for the synthesis of functionalized arenes are based on palladium. Here we present silver catalysis for carbon–fluorine bond formation. Our report is the first example of the use of the transition metal silver to form carbon–heteroatom bonds by cross-coupling catalysis. The functional group tolerance and substrate scope presented here have not been demonstrated for any other fluorination reaction to date.

Introduction

Fluorinated aromatic compounds are used as pharmaceuticals, agrochemicals, materials, and tracers for positron emission tomography (PET).¹ Fluorine incorporation often improves the properties of functional molecules; for example, fluorine substituents can increase the metabolic stability and the rate and extent of blood–brain barrier penetration of pharmaceuticals.² Selective fluorination is important but challenging, especially when the desired fluorinated molecules are complex, and carbon–fluorine bond formation must occur at a late stage of their synthesis.³ In this manuscript we present a silver-catalyzed late-stage fluorination reaction of complex small molecules, including polypeptides, polyketides, and alkaloids (eq 1). Our report is the first example of silver catalysis for carbon–heteroatom bond formation by cross-coupling chemistry. We propose that silver-catalyzed late-stage fluorination proceeds by a mechanism distinct from conventional cross-coupling chemistry.



The synthesis of fluorinated arenes is challenging due to the properties of fluorine.⁴ Electrophilic fluorination is often unselective.⁵ Nucleophilic fluorination is complicated by strong hydrogen bonding and the high hydration energy of the fluoride anion, which results in the low nucleophilicity of hydrated

fluoride and high basicity of dry fluoride.⁶ Carbon–fluorine bond formation by transition metal catalysis is difficult,⁷ in part due to strong ionic metal–fluoride bonding, and only a few examples have been reported.⁸ Reductive elimination, the product forming event in cross-coupling catalysis, becomes increasingly difficult and requires harsher reaction conditions in the bond formation series C–C, C–N, C–O, C–F.^{8d,9} Sanford^{8b} and Yu^{8c} have reported directed palladium-catalyzed electrophilic aromatic fluorination. The advantage of both transformations is the ability to convert C–H bonds into C–F bonds. Challenges include the high reaction temperature (120–150 °C) and the need for directing and blocking groups on the arene

- (1) (a) Yamazaki, T.; Taguchi, T.; Ojima, I. In *Fluorine in Medicinal Chemistry and Chemical Biology*; Ojima, I., Ed.; Wiley-Blackwell: United Kingdom, 2009. (b) Jeschke, P. *ChemBioChem* **2004**, *5*, 570. (c) Babudri, F.; Farinola, G. M.; Naso, F.; Ragni, R. *Chem. Commun.* **2007**, 1003. (2) Müller, K.; Faeh, C.; Diederich, F. *Science* **2007**, *317*, 1881. (3) (a) Wilkinson, J. A. *Chem. Rev.* **1992**, *92*, 505. (b) O'Hagan, D. *Chem. Soc. Rev.* **2008**, *37*, 308. (c) Furuya, T.; Klein, J. E. M. N.; Ritter, T. *Synthesis* **2010**, *11*, 1804.

- (4) For selected fluorination reactions, see: (a) Balz, G.; Schiemann, G. *Ber. Deutsch. Chem. Ges.* **1927**, *60*, 1186. (b) Adams, D. J.; Clark, J. H. *Chem. Soc. Rev.* **1999**, *28*, 225. (c) Sun, H.; DiMaggio, S. G. *Angew. Chem., Int. Ed.* **2006**, *45*, 2720. (d) Sandford, G. J. *Fluorine Chem.* **2007**, *128*, 90. (e) Yamada, S.; Gavryushin, A.; Knochel, P. *Angew. Chem., Int. Ed.* **2010**, *49*, 2215. (f) Anbarasan, P.; Neumann, H.; Beller, M. *Angew. Chem., Int. Ed.* **2010**, *49*, 2219. (5) Baudoux, J.; Cahard, D. *Org. React.* **2007**, *69*, 347. (6) Furuya, T.; Kutruff, C. A.; Ritter, T. *Curr. Opin. Drug Discovery Dev.* **2008**, *11*, 803. (7) For stoichiometric and catalytic transition-metal-mediated C–F bond formation reactions, see: (a) Tius, M. A.; Kawakami, J. K. *Synth. Commun.* **1992**, *22*, 1461. (b) Tius, M. A.; Kawakami, J. K. *Synlett* **1993**, 207. (c) Tius, M. A.; Kawakami, J. K. *Tetrahedron* **1995**, *51*, 3997. (d) Akana, J. A.; Bhattacharyya, K. X.; Muller, P.; Sadighi, J. P. *J. Am. Chem. Soc.* **2007**, *129*, 7736. (e) Kaspi, A. W.; Yahav-Levi, A.; Goldberg, I.; Vigalok, A. *Inorg. Chem.* **2008**, *47*, 5. (f) Furuya, T.; Kaiser, H. M.; Ritter, T. *Angew. Chem., Int. Ed.* **2008**, *47*, 5993. (g) Gorske, B. C.; Mbofana, C. T.; Miller, S. J. *Org. Lett.* **2009**, *11*, 4318. (h) Wu, T.; Yin, G.; Liu, G. *J. Am. Chem. Soc.* **2009**, *131*, 16355. (8) (a) Subramanian, M. A.; Manzer, L. E. *Science* **2002**, *297*, 1665. (b) Hull, K. L.; Anani, W. Q.; Sanford, M. S. *J. Am. Chem. Soc.* **2006**, *128*, 7134. (c) Wang, X.; Mei, T. S.; Yu, J. Q. *J. Am. Chem. Soc.* **2009**, *131*, 7520. (d) Watson, D. A.; Su, M.; Teverovskiy, G.; Zhang, Y.; García-Fortanet, J.; Kinzel, T.; Buchwald, S. L. *Science* **2009**, *325*, 1661. (9) (a) de Meijere, A.; Diederich, F. *Metal-Catalyzed Cross-Coupling Reactions*; Wiley-VCH: Weinheim, 2004. (b) Muci, A. R.; Buchwald, S. L. In *Cross-Coupling Reactions*; Miyaura, N., Ed.; Springer: Berlin, 2002. (c) Hartwig, J. F. *Nature* **2008**, *455*, 314. (d) Furuya, T.; Ritter, T. *J. Am. Chem. Soc.* **2008**, *130*, 10060. (e) Ball, N. D.; Sanford, M. S. *J. Am. Chem. Soc.* **2009**, *131*, 3796. (f) Furuya, T.; Benitez, D.; Tkatchouk, E.; Strom, A. E.; Tang, P.; Goddard, W. A., III; Ritter, T. *J. Am. Chem. Soc.* **2010**, *132*, 3793.

for regioselective fluorination. A breakthrough discovery in 2009 by Buchwald^{8d} extended palladium-catalyzed cross-coupling chemistry to nucleophilic fluorination. The reaction temperature of 80–130 °C in combination with the basicity of fluoride is a current challenge for the synthesis of molecules with protic functional groups and the synthesis of complex aryl fluorides by this method. In some cases, mixtures of regioisomers were observed. A general, functional group-tolerant late-stage fluorination of complex arenes has not yet been achieved by either nucleophilic or electrophilic fluorination reactions.

Silver is used in heterogeneous oxidative catalysis on an industrial scale.¹⁰ In homogeneous catalysis, silver is typically used as a nonredox active Lewis acid, and its redox catalysis is not well understood.¹¹ Cross-coupling catalysis most commonly relies on mononuclear complexes derived from transition metals like palladium, with predictable two-electron redox pathways. Silver undergoes one-electron redox chemistry and previously has not been employed for carbon–heteroatom bond formation by cross-coupling catalysis.


Results and Discussion

We have previously reported the electrophilic fluorination of arylsilver complexes, which required two to three equivalents of silver salt to obtain synthetically useful yields of aryl fluorides.¹² Here, we address the conceptual disadvantage of our earlier results and present silver catalysis. The development of silver catalysis was complicated by protodestannylation as shown in Table 1. Addition of the electrophilic fluorinating reagent F-TEDA-PF₆ to arylstannane **1** in the presence of 10 mol % silver catalyst produced 68% of the undesired protodestannylation product **2a** and only 30% of the desired fluorinated product **2**.


We hypothesized that tributyltin triflate, generated by transmetalation from arylstannanes to AgOTf, is hydrolyzed in situ to form triflic acid (TfOH),¹³ which is responsible for protodestannylation. Fluorination of purified arylsilver complexes, without tributyltin triflate contamination, did not afford a protodestannylation byproduct. Because common drying reagents did not suppress protodestannylation, we evaluated bases to neutralize acids that may form during catalysis and identified sodium bicarbonate (NaHCO₃), which, unlike many other bases (see Supporting Information), was compatible with F-TEDA-PF₆ under the reaction conditions. The use of NaHCO₃ had the additional, though unanticipated, benefit that most of the formed tributyltin bicarbonate precipitated and could be conveniently removed by filtration. Precipitation of the tin residue facilitated product purification, which is often cumbersome when organotin reagents are used.¹⁴

Several soluble Ag(I) salts catalyzed fluorination with equal efficiency, but silver oxide (Ag₂O) was selected as the catalyst of choice because it afforded the fluorinated products in the

Table 1. Silver Catalysis for Carbon–Fluorine Bond Formation



Ag(I) catalyst	Base (2.0 equiv)	Additive	Temp time	Yield ^a 2	Yield ^a 2a
10 mol% AgOTf	none	none	65 °C 3 h	30%	(68%)
5 mol% Ag ₂ O	NaHCO ₃	none	65 °C 5 h	87%	(9%)
5 mol% Ag ₂ O	NaHCO ₃	1.0 equiv NaOTf	65 °C 3 h	90%	(5%)
5 mol% Ag₂O	NaHCO₃	1.0 equiv NaOTf 5.0 equiv MeOH	65 °C 3 h	92%	(2%)
1 mol% Ag ₂ O	NaHCO ₃	1.0 equiv NaOTf	90 °C 18 h	92%	(2%)
1 mol% Ag ₂ O	NaHCO ₃	1.0 equiv NaOTf 5.0 equiv MeOH	90 °C 18 h	75%	(20%)




^a Yields were determined by ¹⁹F and ¹H NMR spectroscopy with 1-fluoro-3-nitrobenzene as a standard. Yields of protodestannylation products are given in parentheses. ^b 5.0 mol % Ag₂O, 1.5 equiv of F-TEDA-PF₆, 2.0 equiv of NaHCO₃, 1.0 equiv of NaOTf, 5.0 equiv of MeOH, acetone, 65 °C, 4 h.

same range of yields as other silver salts and, in contrast to many soluble Ag(I) salts, is inexpensive (\$1.0/g) and not hygroscopic. The addition of NaOTf to the Ag₂O-catalyzed reaction increased the rate of fluorination, possibly due to the conversion of insoluble Ag₂O into a soluble Ag(I) compound under the reaction conditions. While Ag₂O is not soluble in acetone itself, we confirmed the presence of an active, soluble silver catalyst: Upon filtration of the reaction mixture after 5 h, followed by addition of the same amount of all reagents except the silver source, 80% of fluorinated product was observed, based on total added stannane. We determined gravimetrically by precipitation of AgCl that 92% of the silver remained in solution as Ag(I) after fluorination. In the absence of silver catalyst, less than 5% fluorination was observed at 90 °C. The catalyst loading could be reduced to 1 mol % Ag₂O with no loss in yield; however, a longer reaction time and higher reaction temperature were required.

Fluorination using 5 mol % of Ag₂O afforded electron-deficient (**4**), electron-rich (**5**), halogenated (**6**), and *ortho,ortho* disubstituted arenes (**7**). The addition of 5 equiv of methanol decreased the amount of protodestannylation product formed to 2–5% when 5 mol % of Ag₂O was used; addition of MeOH reduced the yield of fluorinated product at 1 mol % catalyst loading. The role of MeOH is currently not understood. All fluorination reactions were performed under ambient atmosphere and in reagent grade acetone as solvent. The reaction temperature of 65 °C is slightly above the boiling temperature of acetone, and therefore, reactions were performed in closed

- (10) Linic, S.; Barteau, M. A. In *Handbook of Heterogeneous Catalysis*; Ertl, G., Knözinger, H., Schüth, F., Weitkamp, J., Eds.; Wiley-VCH: Weinheim, 2008.
- (11) (a) Naodovic, M.; Yamamoto, H. *Chem. Rev.* **2008**, *108*, 3132. (b) Weibel, J. M.; Blanc, A.; Pale, P. *Chem. Rev.* **2008**, *108*, 3149. (c) Álvarez-Corral, M.; Muñoz-Dorado, M.; Rodríguez-García, I. *Chem. Rev.* **2008**, *108*, 3174. (d) Yamamoto, Y. *Chem. Rev.* **2008**, *108*, 3199.
- (12) (a) Furuya, T.; Strom, A. E.; Ritter, T. *J. Am. Chem. Soc.* **2009**, *131*, 1662. (b) Furuya, T.; Ritter, T. *Org. Lett.* **2009**, *11*, 2860.
- (13) Mehring, M.; Nolde, C.; Schürmann, M. *Inorg. Chim. Acta* **2009**, *362*, 745.
- (14) Davis, A. G.; Gielen, M.; Pannell, K. H.; Tieckink, E. R. T. *Tin Chemistry-Fundamentals, Frontiers and Applications*; Wiley: United Kingdom, 2008.

Table 2. Ag₂O-Catalyzed Fluorination of Complex Small Molecules

^a 20 mol % of AgOTf, 2 equiv of NaOTf, and 2 equiv of F-TEDA-PF₆ were used at 90 °C. ^b 20 mol % of AgOTf, 2 equiv of NaOTf, and 5 equiv of MeOH were used. The names under the molecules in Table 2 refer to the parent molecules and not to fluorinated analogs shown.


vessels. The reaction could also be performed at reflux with yields decreased by about 10% and reaction times that are 3 h longer.

We selected a group of small molecules that contain several functional groups and found that the Ag₂O-catalyzed fluorination reaction is general beyond simple arenes (Table 2). Biologically active molecules such as carbohydrates, peptides, polyketides, and alkaloids are compatible with fluorination, and molecules derived from complex natural products such as taxol and rifamycin were successfully fluorinated. It is worth mentioning that fluorination can proceed in the presence of several functional groups, including a vinyl ether, a dienone, alcohols, an allylic alcohol, ethers and esters, and an oxetane. To date, no other fluorination reaction has been shown to have a substrate scope as broad as that shown in here. The practical reaction conditions allowed for the gram-scale synthesis of dipeptide **15** in 92% yield. For all fluorinated molecules shown in Table 2, carbon–fluorine bond formation was performed as the final synthetic step. The yield of protodesstannylated byproducts was less than 10% in all cases, which simplified purification of the fluorinated molecules. Fluorination always occurred regioselectively; no other regioisomers were detected. General high regioselectivity of the silver-catalyzed fluorination is an additional advantage over the known palladium-catalyzed fluorination reactions.

Limitations of the Ag-catalyzed fluorination reaction are shown in Chart 1. We identified that basic functional groups that react with the electrophilic fluorinating reagent unproductively are often not tolerated. Certain amines (**19**) may react with F-TEDA-PF₆ to form *N*-fluoro compounds that subsequently eliminate HF. Amines that lack β-hydrogen atoms for

Chart 1. Limitations of the Presented Fluorination Reaction^a


Amines (dimethylbenzylamine vs bridgehead amine, quinine)



Sulfur-containing substrates (sulfide vs sulfone)



Protic functional groups (carboxylic acid vs alcohol)



^a The fluorination was attempted from the corresponding arylstannane: 5.0 mol % Ag₂O, 1.5 equiv of F-TEDA-PF₆, 2.0 equiv of NaHCO₃, 1.0 equiv of NaOTf, 5.0 equiv of MeOH, acetone, 65 °C, 4 h. ^b 20 mol % of AgOTf, 2 equiv of NaOTf, and 2 equiv of F-TEDA-PF₆ were used at 90 °C.

E2 elimination of HF participate successfully in fluorination: For example, both the quinine and the bridgehead tertiary amine of quinine (**11**) did not impede aryl fluorination. In contrast to sulfides, which likely react as nucleophiles with F-TEDA-PF₆, sulfones are compatible with the reaction conditions (**20**, **21**). Fluorination of substrates with protic functional groups that can be deprotonated by NaHCO₃ proved problematic (**22**). Carboxylate anions generated by the deprotonation of carboxylic acids may react with AgOTf to form silver carboxylates, which are less effective silver salts for silver-mediated

tolerance and substrate scope presented here have not been demonstrated for any other fluorination reaction to date. A current disadvantage of the method is the need for the preparation of arylstannanes as starting materials. In addition, the fluorination reaction reported by Buchwald can afford aryl fluorides from triflates directly, while our fluorination reaction requires an additional synthetic step from triflate or halide to stannane. On the other hand, advantages of our fluorination reaction compared to the Buchwald method are milder reaction conditions, absence of constitutional isomers, and tolerance toward protic functional groups, which make our fluorination reaction particularly useful for late-stage fluorination of complex small molecules, which may not be suitable for other fluorination reactions.

The Ag-catalyzed fluorination reaction likely proceeds through a mechanism distinct from traditional cross-coupling mecha-

nisms. Noteworthy is the C–F bond formation under comparatively mild reaction conditions, by reductive elimination from a putative multinuclear high-valent silver fluoride complex. Silver catalysis, with potential redox synergy between two or more metals, is a new avenue for carbon–heteroatom cross-coupling chemistry.

Acknowledgment. This work was supported by the NSF (CHE-0952753) and NIH-NIGMS (GM088237). We thank Air Products and Chemicals, Inc. for a generous donation of F-TEDA-BF₄.

Supporting Information Available: Experimental procedures and spectroscopic data for all new compounds. This material is available free of charge via the Internet at <http://pubs.acs.org>.

JA105834T