



SAHAquines, Novel Hybrids Based on SAHA and Primaquine Motifs, as Potential Cytostatic and Antiplasmodial Agents

Maja Beus,^[a] Zrinka Rajić,^{*[a]} Dusica Maysinger,^[b] Zvonimir Mlinarić,^[a] Maja Antunović,^[c] Inga Marijanović,^[c] Diana Fontinha,^[d] Miguel Prudêncio,^[d] Jana Held,^[e] Sureyya Olgen,^[f] and Branka Zorc^{*[a]}

We report the synthesis of SAHAquines and related primaquine (PQ) derivatives. SAHAquines are novel hybrid compounds that combine moieties of suberoylanilide hydroxamic acid (SAHA), an anticancer agent with weak antiplasmodial activity, and PQ, an antimalarial drug with low antiproliferative activity. The preparation of SAHAquines is simple, cheap, and high yielding. It includes the following steps: coupling reaction between primaquine and a dicarboxylic acid monoester, hydrolysis, a new coupling reaction with *O*-protected hydroxylamine, and deprotection. SAHAquines **5** a–d showed significant reduction in cell viability. Among the three human cancer cell lines (U2OS,

HepG2, and MCF-7), the most responsive were the MCF-7 cells. The antibodies against acetylated histone H3K9/H3K14 in MCF-7 cells revealed a significant enhancement following treatment with *N*-hydroxy-*N*'-{4-[(6-methoxyquinolin-8-yl)amino]pentyl}-pentanediamide (**5 b**). Ethyl (2*E*)-3-({4-[(6-methoxyquinolin-8-yl)amino]pentyl}carbamoyl)prop-2-enoate (**2 b**) and SAHA-quines were the most active compounds against both the hepatic and erythrocytic stages of *Plasmodium* parasites, some of them at sub-micromolar concentrations. The results of our research suggest that SAHAquines are promising leads for new anticancer and antimalarial agents.

1. Introduction

Despite extensive efforts and significant progress in cancer treatment, therapeutic interventions are not yet satisfactory. Almost all available cytostatic drugs cause undesirable side effects, whereas drug resistance presents an additional problem.^[1-3] In another field of medicine, an ongoing battle is being fought against malaria, a parasitic disease caused by *Plasmodium* species.^[4,5] Malaria still poses a great health and economic burden to the highly populated countries in the tropical and subtropical parts of the world. The need for new, effective antimalarials arises from several factors, including the absence of an effective vaccine, insufficient vector control, and the emergence of multidrug-resistant *Plasmodium* strains.^[6-8]

modial compounds include the development of analogues of existing drugs, resistance reversers, and novel compounds with new mechanisms of action.^[6,9]

A number of studies have shown a relationship between cancer and malaria in regard to diagnostics, drug research, treatment, prevention, and epidemiology.^[10-13] Different classes of antimalarial drugs display direct or adjuvant activity against cancer cell lines, are known as sensitivity reversers of resistant tumor cell lines or inhibitors of drug resistance development, or have synergistic action with known anticancer drugs.^[14-20] Although their exact mode of action against cancer is still not completely understood, various mechanisms have been proposed.^[21-23]

[a]	M. Beus, Prof. Z. Rajić, Z. Mlinarić, Prof. B. Zorc Faculty of Pharmacy and Biochemistry, University of Zagreb A. Kovačića 1, 10 000 Zagreb (Croatia) E-mail: zrajic@pharma.hr bzorc@pharma.hr	[e] [f]	Dr. J. Held Institute of Tropical Medicine, University of Tübingen Wilhelmstraße 27, 72074 Tübingen (Germany) Prof. S. Olgen Erscultu of Pharmacy, Biruni University
[b]	Prof. D. Maysinger		10th street No: 45, 34010 Topkapi Istanbul (Turkey)
	Department of Pharmacology and Therapeutics, McGill University 23655 Promenade Sir-William-Osler, McIntyre Medical Sciences Building Montreal, Quebec H3G 1Y6 (Canada)	Þ	Supporting Information and the ORCID identification number(s) for the author(s) of this article can be found under: https://doi.org/10.1002/open.201800117.
[c]	Dr. M. Antunović, Prof. I. Marijanović Faculty of Science, University of Zagreb Horvatovac 102A, 10 000 Zagreb (Croatia)	ס	© 2018 The Authors. Published by Wiley-VCH Verlag GmbH & Co. KGaA. This is an open access article under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs License, which permits use and
[d]	Dr. D. Fontinha, Dr. M. Prudêncio Instituto de Medicina Molecular, Faculdade de Medicina Universidade de Lisboa Av. Prof. Egas Moniz, 1649-028 Lisboa (Portugal)		distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.





Such observations prompted us, and others, to design and prepare novel derivatives of known antimalarial drugs and evaluate their cytostatic potential.[24-34] Our efforts have been focused on primaguine (PQ), which is an old drug with many flaws (e.g., induction of hemolytic anemia in individuals lacking glucose-6-phosphate dehydrogenase, guick metabolism, degradation to inactive carboxyprimaquine, altering the treatment outcome in dependence of CYP 2D6 enzyme activity)^[35] but is currently the only available *Plasmodium* hypnozoitocide.^[36] We previously showed that various PQ derivatives of amides, ureas, bis-ureas, semicarbazides, and acylsemicarbazide-type derivatives possessed significant cytostatic activity against a panel of cancer cell lines or high selectivity towards the breast adenocarcinoma cell line (MCF-7).^[24-30] On the other hand, cytostatic agents of different classes, including histone deacetylase (HDAC) inhibitors, have been shown to exert antiplasmodial activity.^[6,37-42] Suberoylanilide hydroxamic acid (SAHA, vorinostat) was the first marketed pan-HDAC inhibitor, and it was followed by other drugs from the same class (e.g., romidepsin, belinostat, and panobinostat.[43,44] Their hydroxamic group chelates zinc ion found in the active site of Zn-dependent HDACs (classes I, II, and IV), and this leads to the accumulation of acetylated histones and other proteins.^[45] HDACs were also identified as transcription regulators in *P. falciparum*.^[46,47] Out of five P. falciparum HDACs, three are Zn-dependent enzymes, prone to inhibition by SAHA, and thus represent viable targets for drug development.^[48] Several studies have shown the antimalarial activity of SAHA and related HDAC inhibitors.[49,50]

In this study, we employed one of the classical medicinal chemists' tools—the combination of two pharmacophores in one molecule.^[7,51-54] The concept of hybrid drugs is also a valuable strategy to overcome the limitations of a combined therapy, as the resulting molecules could exhibit inhibitory activities on multiple targets.^[55] The hybrid compounds described here, SAHAquines, combine motifs of SAHA, an anticancer agent with weak antiplasmodial activity, and PQ, an antimalarial drug with low antiproliferative activity. Other here-reported PQ derivatives differ in the linker length/type and/or functional groups: compounds **2** are esters, compounds **3** are carboxylic



Figure 1. Design of SAHAquines 5 and other PQ derivatives.

acids, and compounds **4** and **6** are *O*-benzyl- and *O*-methylsubstituted hydroxamic acids. Similar to the known HDAC inhibitors, hydroxamic acids **5** consist of a capping group (quinoline ring), a linker, and a Zn-binding group (hydroxamic acid) (Figure 1). Given that hydroxamic acid is a strong binding group for metal ions that might lead to poor selectivity and confer undesired properties, such as poor pharmacokinetics,^[56,57] we introduced other functional groups instead: an ester, carboxylic acid, or *O*-protected hydroxamic group.

Herein, we report the synthesis of SAHAquines **5** a–d as well as 20 novel PQ derivatives, their chemical characterization, the assessment of their cytostatic activity, and the evaluation of their activity against the erythrocytic and hepatic stages of *Plasmodium*.

2. Results and Discussion

2.1. Chemistry

The objective of our research was to prepare SAHAquines 5 and related PQ derivatives, which differ in the type/length of the spacer and/or functional groups: compounds 2 are esters and compounds 3 are carboxylic acids, whereas compounds 4 and 6 are O-protected hydroxamic acids. Scheme 1 shows the synthetic pathway leading to derivatives 2-6. In the first reaction step, dicarboxylic acid monoesters 1a-e were coupled with PQ to give derivatives 2a-e by using 1-[bis(dimethylamino)methylene]-1H-1,2,3-triazolo[4,5-b]pyridinium 3-oxid hexafluorophosphate (HATU) as the coupling reagent, along with Hünig's base (N,N-diisopropylethylamine, DIEA). The following dicarboxylic acid monoesters were used: 4-methoxy-4-oxobutanoic acid (monomethyl hydrogen succinate) (1 a), (E)-4ethoxy-4-oxobut-2-enoic acid (monoethyl fumarate) (1b), 5methoxy-5-oxopentanoic acid (monomethyl glutarate) (1 c), 6methoxy-6-oxohexanoic acid (monomethyl adipate) (1 d), and 4-(methoxycarbonyl)benzoic acid (monomethyl terephtalate) (1 e). Notably, an analogous coupling reaction between PQ and monomethyl malonate failed in our hands. Amide formation between these two compounds by using thionyl chloride or benzotriazolide^[58] was also unsuccessful and gave a mixture of products. Classical activation of the carboxylic group with thionyl chloride worked well with other dicarboxylic acid monoesters. Hydrolysis of 2a-e with lithium hydroxide afforded corresponding acids 3a-e, which were transformed into O-benzylhydroxamic acids 4a-e and O-methylhydroxamic acids 6a-e by means of O-benzyl- and O-methylhydroxylamine, respectively. Again, HATU/DIEA was used as the coupling system. Free hydroxamic acids **5** a-d were obtained by catalytic hydrogenation of O-benzyl derivatives 4a-e. Selective deprotection of the benzyl group in fumaric derivative 4b failed. As a result of double-bond hydrogenation, succinylhydroxamic acid 5a was obtained instead of fumarylhydroxamic acid.

All new compounds were fully characterized by conventional spectroscopy and analytical methods (IR, ¹H NMR, and ¹³C NMR spectroscopy; MS; and elemental analyses). The data were consistent with the proposed structures and are given in short in







Scheme 1. Synthesis of PQ derivatives 2–6. Reagents and conditions: a) HATU, DIEA, CH₂Cl₂, RT, 1 h; b) LiOH, MeOH, H₂O, RT, 1 h; c) H₂NOBn, HATU, DIEA, CH₂Cl₂, RT, 2 h; d) H₂, Pd/C, MeOH, RT, 2–4 h; NH₂OMe, HATU, DIEA, CH₂Cl₂, RT, 2 h.

the Experimental Section and in detail in the Supporting Information.

To evaluate the drug-like properties of our novel compounds, a common set of physicochemical parameters were calculated: topological polar surface area (TPSA), number of atoms, molecular weight (MW), partition coefficient (log *P*), number of H-bond donors (HBDs), number of H-bond acceptors (HBAs), and molar refractivity (MR). The parameters were calculated with Chemicalize.org software and are presented in Table 1.^[59] All compounds **2–6** (except **4e**, which showed minimal aberration) are fully in agreement with Lipinski's and Gelovani's rules for prospective small-molecule drugs (MW \leq 500, log $P \leq$ 5, number of H-bond donors \leq 5, number of H-bond acceptors \leq 10, TPSA < 140 Å², MR within the range of 40 and 130 cm³ mol⁻¹, number of atoms 20–70).^[60]

Compd	Molecular formula	Number of atoms	MW	log P	HBD ^[a]	HBA ^[b]	Lipinski score ^[c]	$MR^{[d]}$ [cm ³ mol ⁻¹]	TPSA ^[e] [Å ²
2 a	$C_{20}H_{27}N_{3}O_{4}$	54	373.453	1.43	2	5	4	103.62	89.55
2 b	$C_{21}H_{27}N_3O_4$	55	385.464	2.38	2	5	4	109.44	89.55
2 c	$C_{21}H_{29}N_3O_4$	57	387.480	1.88	2	5	4	108.22	89.55
2 d	$C_{22}H_{31}N_3O_4$	60	401.507	2.32	2	5	4	112.82	89.55
2 e	C ₂₄ H ₂₇ N ₃ O ₄	58	421.497	3.32	2	5	4	120.65	89.55
3 a	$C_{19}H_{25}N_3O_4$	51	359.426	0.80	3	6	4	98.85	100.55
3 b	$C_{19}H_{23}N_3O_4$	49	357.410	0.90	3	6	4	99.92	100.55
3 c	C ₂₀ H ₂₇ N ₃ O ₄	54	373.453	1.29	3	6	4	103.45	100.55
3 d	$C_{21}H_{29}N_3O_4$	57	387.480	1.74	3	6	4	108.05	100.55
3 e	$C_{23}H_{25}N_3O_4$	55	407.470	2.23	3	6	4	115.89	100.55
4a	$C_{26}H_{32}N_4O_4$	66	464.566	2.58	3	6	4	131.53	101.58
4b	$C_{26}H_{30}N_4O_4$	64	462.515	2.94	3	6	4	132.60	101.58
4 c	$C_{27}H_{34}N_4O_4$	69	478.580	3.02	3	6	4	136.13	101.58
4 d	$C_{28}H_{36}N_4O_4$	72	492.620	3.47	3	6	4	140.74	101.58
4e	$C_{30}H_{32}N_4O_4$	70	512.600 ^[b]	4.27	3	6	3 ^[f]	148.57	101.58
5 a	$C_{19}H_{26}N_4O_4$	53	374.441	0.48	4	6	4	102.44	112.58
5 b	$C_{20}H_{28}N_4O_4$	56	388.468	0.92	4	6	4	107.04	112.58
5 c	$C_{21}H_{30}N_4O_4$	59	402.495	1.37	4	6	4	111.64	112.58
5 d	$C_{23}H_{26}N_4O_4$	57	422.485	2.16	4	6	4	119.47	112.58
бa	$C_{20}H_{28}N_4O_4$	56	388.468	0.85	3	6	4	106.92	101.58
6 b	$C_{20}H_{26}N_4O_4$	54	386.452	1.21	3	6	4	107.99	101.58
бc	$C_{21}H_{30}N_4O_4$	59	402.495	1.30	3	6	4	111.52	101.58
6 d	$C_{22}H_{32}N_4O_4$	62	416.522	1.74	3	6	4	116.12	101.58
бe	C ₂₄ H ₂₈ N ₄ O ₄	60	436.512	2.54	3	6	4	123.95	101.58

626



2.2. Biological Evaluation

Synthesized compounds 2-6 were tested for their anticancer activity on three human cancer cell lines (bone osteosarcoma U2OS, hepatocellular carcinoma HepG2, and breast adenocarcinoma MCF-7) and human embryonic kidney (Hek293) cells. The cells were treated with the different compounds at different concentrations, and the median inhibitory concentration (IC₅₀) values were determined (Table 2).

The data from Table 2 clearly indicate that the SAHAquines were the most potent in the selected cancer cell lines. The $\mathrm{IC}_{\mathrm{50}}$ values towards MCF-7 cells were in the low micromolar concentrations (1.6-5.4 µm). Compounds from the other subclasses were, in general, less active than compounds 5. Compounds 4d and 4e were the most effective O-benzylhydroxamic acids. In contrast, O-methylhydroxamic acids 6 exhibited weak activity, except for succinic acid derivative 6a. All compounds from the ester series, except 2c, showed moderate activity, whereas carboxylic acid derivatives **3**a-e were practically inactive.

MCF-7 cells were sensitive to 15 of the 24 tested compounds, namely, 5a-d, 2d, 2e, 4a, 4c-e, and 6a-d. Similar results were obtained in our previous studies with various PQ

Table 2. Antiproliferative screening of novel compounds 2–6 towards human cancer cell lines (U2OS, HepG2, MCF-7) and human embryonic kidney cell line (Hek293) in vitro. Compd Structure IC₅₀^[а] [µм] U2OS HepG2 MCF-7 Hek293 6.7 ± 3.1 2 a > 50> 50> 50 2b 14.9 ± 2.2 17.0±1.7 >50 >50 2 c >50 >50 >50 > 50 2 d 24.3 ± 1.5 >50 5.3 ± 3.2 > 50 2 e >50 >50 7.1 ± 3.0 > 50 > 50 3 a > 50>50 > 503b > 50 >50 >50 > 50 3 c 33.8±2.9 >50 >50 > 50 3 d > 50 >50 >50 >50





Table 2. (Continued)							
Compd	Structure	IC ₅₀ ^[а] [µм] U2OS	HepG2	MCF-7	Hek293		
3 e		>50	>50	15.2±0.5	>50		
4a		> 50	> 50	12.0±1.4	> 50		
4 b		> 50	> 50	> 50	> 50		
4c		> 50	> 50	16.6±0.8	> 50		
4 d		32.8±3.0	9.9±1.1	13.9±1.5	> 50		
4e		7.1±1.2	> 50	11.5±0.1	13.2±2.6		
5a		16.9±2.2	18.0±3.3	5.4±0.1	25.5±0.8		
5 b		3.3±0.05	20.0±0.2	1.6±0.8	8.5±0.5		
5 c		11.6±2.0	28.4±2.1	5.0±0.4	13.7±3.4		
5 d		11.6±2.1	21.7±0.2	4.7±0.1	6.1±1.9		
ба		6.6±1.2	>50	5.9 ± 0.6	8.6±2.0		
6b		>50	>50	10.0±2.8	>50		





Table 2. (Con	tinued)				
Compd	Structure	IC ₅₀ ^[а] [µм] U2OS	HepG2	MCF-7	Hek293
6c		> 50	> 50	7.1±0.4	> 50
6 d		> 50	>50	9.8±0.4	> 50
бe		> 50	> 50	> 50	> 50
PQ ^[b] SAHA ^[c] Cis ^[d]	U	$\begin{array}{c} 12.0 \pm 0.7 \\ 5.7 \pm 0.8 \\ 2.5 \pm 0.7 \end{array}$	37.7±5.8 4.0±0.1 -	$\begin{array}{c} 13.8 \pm 1.4 \\ 2.8 \pm 0.7 \\ 3.2 \pm 1.0 \end{array}$	$\begin{array}{c} 8.1 \pm 1.3 \\ 7.4 \pm 0.9 \\ 2.0 \pm 0.6 \end{array}$
[a] The conce	ntration required to decrease viability by 50%. [b) Primaquine. [c] Suberoy	anilide hydroxamic acid.	[d] Cisplatin.	

derivatives.^[24-30] On the other hand, HepG2 cells were very robust and only responded towards **4d** and compounds from subclass **5**. The non-cancer human cell line Hek293 was sensitive to **2b**, **4e**, **5a**–**d**, and **6a** but to a lesser extent than the cancer cell lines. Selectivity indices ranged from 2 to 12, depending on the tested compound and the cancer cell line employed.

To show the loss of MCF-7 cells following their exposure to the most potent compounds from subclasses ${\bf 4}$ and ${\bf 5}$, that is,

4e and **5b**, we labeled the cell nuclei with Hoechst 33342 and assessed the number of cells (Figure 2). The results from these studies complement the data from the MTT [3-(4,5-dimethylth-iazol-2-yl)-2,5-diphenyltetrazolium bromide] assay (Table 2), showing a correlation between significant loss of mitochondrial metabolic activity and a concentration-dependent decrease in the number of cells. A reduction in the number of MCF-7 cells was found for treatment over periods of 24 and 72 h. The IC₅₀ values for **5b** and **4e** obtained from these assays were



Figure 2. MCF-7 cell viability following treatment with **4e**, **5b**, primaquine, SAHA, and cisplatin for 24 or 72 h at various concentrations ranging from 50 to 0.001 μ M. Cell viability was measured by counting Hoechst 33342 labeled nuclei imaged by using a fluorescence microscope. Shown are average percentage cell viability compared to the untreated controls \pm SD from two independent experiments.

ChemistryOpen 2018, 7, 624-638





within a comparable micromolar concentration range (**5 b**: 0.5 and 1.6 μ M; **4e**: 9.0 and 11.5 μ M). Determination of the number of cells is a more direct way of showing cell loss.

As SAHAquines **5** with a free hydroxamic acid have the potential to chelate metal ions in Zn-dependent HDACs and to increase the content of acetylated histones, we performed a immunocytochemical assay for histone acetylation. MCF-7 cells were treated with **5b** and acetylated H3K9/H3K14 was measured. SAHA, PQ, and **4e** were used as controls in the histone acetylation experiments. Indeed, the results show that **5b** caused a significant accumulation of acetylated H3 histone (Figure 3), which suggested that HDAC inhibition was a possible contributing mechanism for SAHAquines **5** but not for the other PQ derivatives. Given that HDAC inhibition correlated with cellular histone acetylation but not with cell loss, we did not extensively study the kinetics of HDAC inhibition.^[61] An indepth mechanistic study should be performed to explain the mode of cell death caused by the SAHAquines.



Figure 3. MCF-7 cells were treated with PQ, SAHA, **4e**, or **5b** (1 μ M) for 24 h. a) Representative fluorescence micrographs of MCF-7 histone H3 acetylation (red) at lysine 9/lysine 14 in response to treatments. Cells were incubated with primary antibody (rabbit anti-acetyl-Histone-H3 polyclonal antibody, 500× dilution) for 24 h and then with secondary antibody (Alexa Fluor 647 goat anti-rabbit IgG -2 mg mL^{-1} ; 500× dilution) for 1 h in the dark. Cells were imaged by using a fluorescence microscope with a CY5 filter. Fluorescence was analyzed in ImageJ. Scale bar represents 20 μ M. b) Averages of fluorescence per cell ± SD (as fold increase in untreated control = 1) from at least two independent experiments. * p < 0.05, *t*-test.

Literature data on the antimalarial activity of SAHA and related HDAC inhibitors^[49, 50] prompted us to evaluate SAHAquines and their synthetic precursors for their in vitro activity against P. falciparum erythrocytic stages. The IC₅₀ values of SAHAquines 2-6 against the erythrocytic stage of two P. falciparum strains, 3D7 and Dd2, were determined (Table 3). In general, the 3D7 strain was more sensitive than the Dd2 strain. Again, the most active compounds were SAHAquines 5, the hydroxamic acid subclass. Compound **5b** had the lowest IC_{50} value (0.4 μ M for the Pf3D7 strain and 1.9 μ M for the PfDd2 strain), followed by 5d (IC_{50}\!=\!0.6 and 1.2 $\mu \text{m},$ respectively), 5c (IC_{50}\!=\!3.7 and 13.6 μ M, respectively), and finally **5a** (IC₅₀ = 15.8 and 27.1 μ M, respectively). Derivatives 2b and 2d were the most active compounds from the ester subclass, and 4e was the most active from the O-benzylhydroxamic acids. Carboxylic acids 3a-e were inactive.

We further evaluated the in vitro activity of compounds **2–6** against the hepatic stages of *P. berghei*. Two concentrations of SAHAquines were tested: 1 and 10 μ M. For comparison, the same concentrations of PQ were included in the assay as a positive control (IC₅₀=8.4 \pm 3.4 μ M), and DMSO was used as a negative control (Figure 4).

As shown in Figure 4, only ester **2b** and SAHAquines **5a–d** were active against the *P. berghei* hepatic stages, whereas the other tested compounds were completely inactive. All compounds from hydroxamic acid subclass **5** exhibited strong antiplasmodial activity at both concentrations tested, with IC_{50} values ranging from 0.3 to 1.25 μ M and without any noticeable effects on host-cell confluency (Figure 5). Our results show that the free hydroxamic acid moiety was crucial for antiplasmodial activity, as activity was lost if this group was protected or replaced by a carboxylic acid. The activity of compound **2b** from the ester subclass was probably due to the α , β -unsaturated carbonyl group (Michael acceptor moiety), which is capable of conjugate addition.

The relationship between antimalarial and anticancer activity is complex and is discussed in detail in a review by Duffy, Wade, and Chang.^[62] Both main groups of existing antimalarial

Table 3. IC ₅₀ values for compounds 2–6 against erythrocytic stage of two <i>P. falciparum</i> strains.							
Compd	IC ₅₀ [µм] <i>Pf</i> 3D7	Pf3Dd2	Compd	IC ₅₀ [µм] <i>Pf</i> 3D7	Pf3Dd2		
2a	100.0 ± 11.0	$74.1 \pm 5.3 \\ 7.2 \pm 1.9 \\ 80.1 \pm 31.0 \\ 50.4 \pm 4.6 \\ 25.0 \pm 2.8 \\ > 111 \\ > 111 \\ > 111 \\ > 111$	4d	> 27.7	> 27.7		
2b	2.9 \pm 0.2		4e	6.1 \pm 0.1	> 27.7		
2c	81.1 \pm 3.4		5a	15.8 \pm 1.7	27.1 ± 0.8		
2d	6.6 \pm 0.2		5b	0.4 \pm 0.1	1.9 ± 0.8		
2e	27.1 \pm 0.6		5c	3.7 \pm 1.3	13.6 ± 0.2		
3a	> 111		5d	0.6 \pm 0.2	1.2 ± 0.04		
3b	> 111		6a	> 111	94.4 ± 8.6		
3c	> 111		6b	8.3 \pm 0.9	14.8 ± 0.1		
3 d	> 111	> 111	6 c	> 111	> 111		
3 e	> 111	> 111	6 d	> 111	> 111		
4 a	> 55	50.0 ± 5.0	6 e	> 25	> 27.7		
4 b	> 27.7	> 27.7	PQ ^[a]	1.5 ± 0.02	4.3 ± 1.5		
4 c	39.5 ± 15.5	35.5 ± 1.2	CQ ^[b]	1.6×10^{-3}	0.265 ± 0.003		

630

ChemistryOpen 2018, 7, 624–638







Figure 4. Activity of compounds 2–6 against *P. berghei* hepatic stages at concentrations of 1 and 10 µm. Anti-infective activity (infections scale, bars) are shown.

agents, the quinolines and the sesquiterpene lactones, have a range of known anticancer properties that could either be distinct from or overlap with their antimalarial properties. Although our results show a similar pattern of activity, it should be noted that the activities of **5 b** and **5 d** are an order of magnitude higher towards the *P. falciparum* erythrocytic stage than towards the tested cancer cell lines and the human embryonic kidney cells (Hek293).

Having in mind that very good safety profiles are needed for antimalarial drugs, as they should be delivered, amongst others, to two particularly vulnerable populations, small children and pregnant women,^[63] we are aware that our finding can only be a starting point for the development of a clinically applicable antimalarial drug. As the antiplasmodial activities of compounds **2b** and **5a-d** were comparable to the activity of PQ, they might potentially replace PQ against PQ-resistant *Plasmodium* strains or in the treatment of glucose-6-phosphate dehydrogenase deficient patients and poor metabolizers, but this hypothesis still remains to be evaluated. Notably, the synthesis of SAHAquines is cheap and short (1–4 h) with good to excellent yields (43–97 %), which is beneficial for the development of antimalarial agents.^[64]

3. Conclusions

Four SAHAquines based on SAHA and PQ motifs and 20 other PQ derivatives were synthesized and evaluated in vitro against three human cancer cell lines and a human embryonic kidney







Figure 5. IC₅₀ of SAHAquines 5 a-d against *P. berghei* hepatic stages.

cell line. SAHAquines **5a-d** displayed cytostatic activity at low micromolar concentrations. A few compounds from other subclasses were also effective, but less so than **5a-d**. We showed that the most active hydroxamic acid, that is, **5b**, caused a significant accumulation of acetylated histone H3K9/H3K14, a downstream target of class I HDACs. The results from the antiplasmodial activity of SAHAquines against the erythrocytic stages of the 3D7 and Dd2 *P. falciparum* strains and against *P. berghei* hepatic stages correlated with their cytostatic activity of SAHAquines suggest that these novel compounds could constitute a basis for the development of effective anticancer or malaria prophylactic/curative agents with improved potency and selectivity.

Experimental Section

General Methods

Melting points were measured with a Stuart Melting Point (SMP3) apparatus (Barloworld Scientific, UK) in open capillaries. IR spectra were recorded with a FTIR PerkinElmer Spectrum One, and UV/Vis spectra were recorded with a Lambda 20 double-beam spectrophotometer (PerkinElmer, UK). All NMR (¹H and ¹³C) spectra were obtained at 25 °C by using an NMR Avance 600 spectrometer (Bruker, Germany) at 300 and 150 MHz for ¹H and ¹³C nuclei, respectively. Chemical shifts (δ) are reported in parts per million (ppm) relative to tetramethylsilane in the ¹H spectra and relative to $[D_6]DMSO$ in the ¹³C spectra (δ = 39.51 ppm). Coupling constants (J) are reported in hertz. Mass spectra were collected with a HPLC-MS/MS instrument (HPLC, Agilent Technologies 1200 Series; MS, Agilent Technologies 6410 Triple Quad). Mass determination was realized by using electrospray ionization (ESI) in the positive mode. Elemental analyses were performed with a CHNS LECO analyzer (LECO Corporation, USA). All compounds were routinely checked by TLC with Merck silica gel 60F-254 glass plates by using the following solvent systems: dichloromethane/methanol 9:1, 9.5:0.5, and 9.6:0.4; cyclohexane/ethyl acetate 1:1; and cyclohexane/ethyl acetate/methanol 3:1:0.5. Spots were visualized by short-wave UV light and iodine vapor. Column chromatography was performed on silica gel 0.063-0.200 mm. All chemicals and solvents were of analytical grade and were purchased from commercial sources. PQ diphosphate, 4-methoxy-4-oxobutanoic acid (monomethyl succinate), (E)-4-ethoxy-4-oxobut-2-enoic acid (monoethyl fumarate), 5methoxy-5-oxopentanoic acid (monomethyl glutarate), 6-methoxy-6-oxohexanoic acid (monomethyl adipate), 4-(methoxycarbonyl)benzoic acid (monomethyl terephtalate), DIEA, HATU, O-benzylhydroxylamine, O-methylhydroxylamine, triethylamine, MTT, cisplatin (Cis), and Hoechst 33342 were purchased from Sigma-Aldrich; rabbit anti-acetyl-Histone-H3 polyclonal antibody was purchased from Millipore, Alexa Fluor 647 goat anti-rabbit IgG was purchased from Life Technologies, and SAHA was acquired from Cayman Chemicals. PQ was prepared from PQ diphosphate prior to use. All reactions with PQ were performed under light-protected conditions.

Synthesis

General Procedure for the Preparation of Esters 2a-e

Method A: A solution of dicarboxylic acid monoester 1a-e (1.4 mmol), HATU (0.532 g, 1.4 mmol), and DIEA (0.362 g, 2.8 mmol) in dichloromethane (8 mL) was stirred at room temperature. After 10 min, a solution of PQ (0.401 g, 1.5 mmol) in dichloromethane (7 mL) was added. The mixture was stirred at room temperature for 1 h and was then concentrated under reduced pressure. The residue was dissolved in ethyl acetate and extracted with brine (3×). The organic layer was dried with sodium sulfate, filtered, and concentrated under reduced pressure.

Method B: A solution of dicarboxylic acid monoester 1a-e (1.8 mmol) in thionyl chloride (7 mL) was kept overnight and concentrated under reduced pressure. The residue was triturated with dichloromethane (2×), and the solvent was evaporated again. A solution of PQ (0.401 g, 1.5 mmol) and Et₃N (0.152 g, 1.5 mmol) in dichloromethane (8 mL) was added dropwise to the carboxylic acid chloride dissolved in dichloromethane (10 mL). The mixture was stirred at room temperature for 1 h and extracted with brine (3×). The organic layer was dried with sodium sulfate, filtered, and concentrated under reduced pressure.





Methyl 3-({4-[(6-methoxyquinolin-8-yl)amino]pentyl}carbamoyl)propanoate (2a): Method A, from the reaction of monomethyl succinate (1a, 0.185 g) and after purification by column chromatography (dichloromethane/methanol 9.5:0.5) and crystallization (ether), 2a was obtained as a pale-yellow solid (0.460 g, 88%): m.p. 71-73 °C; ¹H NMR ([D₆]DMSO): δ = 8.55–8.53 (dd, J=1.6, 4.2 Hz, 1 H, 10), 8.09–8.06 (dd, J=1.5, 8.3 Hz, 1 H, 12), 7.86 (t, J=5.3 Hz, 1 H, 1), 7.45–7.40 (m, 1 H, 11), 6.47 (d, J=2.4 Hz, 1 H, 16), 6.26 (d, J=2.4 Hz, 1H, 14), 6.12 (d, J=8.7 Hz, 1H, 7), 3.82 (s, 3H, 17), 3.66-3.57 (m, 1H, 5), 3.56 (s, 1H, 5'), 3.08–3.02 (m, 2H, 2), 2.48 (t, J=6.6 Hz, 2H, 2'), 2.33 (t, J=6.8 Hz, 2H, 3'), 1.70-1.58, 1.58-1.42 (2m, 4H, 3,4), 1.20 ppm (d, J = 6.3 Hz, 3 H, 6); ¹³C NMR ([D₆]DMSO): $\delta = 173.30$, $170.87, \ 159.00, \ 144.62, \ 144.23, \ 134.79, \ 134.52, \ 129.57, \ 122.09,$ 96.09, 91.58, 54.97, 51.21, 46.97, 38.46, 33.37, 29.82, 28.82, 25.95, 20.19 ppm; IR (ATR): $\tilde{\nu} = 2278$, 3306, 3060, 2959, 2927, 2857, 1732, 1636, 1619, 1552, 1518, 1454, 1387, 1224, 1200, 1170, 1052, 1031, 825, 793, 679 cm⁻¹; MS (ESI): *m/z*: 374.2 [*M*+1]⁺; elemental analysis calcd (%) for $C_{20}H_{27}N_{3}O_{4}$ (373.45): C 64.32, H 7.21, N 11.25; found: C 64.02, H 7.47, N 11.00.

Ethyl (2E)-3-({4-[(6-methoxyquinolin-8-yl)amino]pentyl}carbamoyl)prop-2-enoate (2b): Method B, from the reaction of monoethyl fumarate (1b, 0.202 g) and after purification by column chromatography (dichloromethane/methanol 9.6:0.4) and crystallization (ether), 2b was obtained as a pale-yellow solid (0.378 g, 70%): m.p. 71–73 °C; ¹H NMR ([D₆]DMSO): δ = 8.54–8.53 (dd, J = 1.6, 4.2 Hz, 1 H, 10), 8.51 (t, J=5.5 Hz, 1 H, 1), 8.08-8.06 (dd, J=1.5, 8.3 Hz, 1 H, 12), 7.43-7.41 (m, 1H, 11), 6.98 (d, J=15.5 Hz, 1H, 2'), 6.55 (d, J= 15.5 Hz, 1 H, 3'), 6.47 (d, J=2.4 Hz, 1 H, 16), 6.27 (d, J=2.4 Hz, 1 H, 14), 6.13 (d, J=8.8 Hz, 1 H, 7), 4.20–4.16 (q, J=7.1 Hz, 2 H, 5'), 3.82 (s, 3 H, 17), 3.66-3.61 (m, 1 H, 5), 3.20-3.17 (dd, J=6.4, 12.3 Hz, 2 H, 2), 1.71–1.65, 1.62–1.51 (2m, 4H, 3,4), 1.23 (t, J=7.1 Hz, 3H, 6'), 1.22 ppm (d, J = 6.3 Hz, 3 H, 6); ¹³C NMR ([D₆]DMSO): $\delta = 165.05$, 162.63, 158.98, 144.60, 144.20, 137.60, 134.76, 134.51, 129.55, 128.08, 122.06, 96.11, 91.51, 60.60, 54.94, 46.93, 38.81, 33.36, 25.63, 20.18, 13.97 ppm; IR (ATR): \tilde{v} = 3391, 3325, 3079, 2969, 2939, 1716, 1645, 1616, 1554, 1520, 1457, 1424, 1388, 1365, 1334, 1302, 1224, 1205, 1163, 1159, 1038, 999, 831, 819, 790, 681, 660 cm⁻¹; MS (ESI): m/z: 386.2 $[M+1]^+$; elemental analysis calcd (%) for C₂₁H₂₇N₃O₄ (385.46): C 65.44, H 7.06, N 10.90; found: C 65.25, H 7.09, N 10.70.

Methyl 4-({4-[(6-methoxyguinolin-8-yl)amino]pentyl}carbamoyl)butanoate (2 c): Method A, from the reaction of monomethyl glutarate (1 c, 0.206 g) and after purification by column chromatography (dichloromethane/methanol 9.5:0.5), 2c was obtained as an oil (0.439 g, 81%): ¹H NMR ([D₆]DMSO): $\delta = 8.53$ (d, J = 2.6 Hz, 1 H, 10), 8.07 (d, J=8.2 Hz, 1 H, 12), 7.78 (t, J=3 Hz, 1 H, 1), 7.43-7.41 (m, 1 H, 11), 6.47 (d, J=2.2 Hz, 1 H, 16), 6.26 (d, J=2.2 Hz, 1 H, 14), 6.11 (d, J=8.8 Hz, 1 H, 7), 3.82 (s, 3 H, 17), 3.64-3.60 (m, 1 H, 5), 3.57 (s, 3 H, 6'), 3.05 (t, J=6.0 Hz, 2 H, 2), 2.27 (t, J=3.0 Hz, 2 H, 2'), 2.07 (t, J=7.3 Hz, 2 H, 4'), 1.74-1.69 (m, 2 H, 3'), 1.67-1.62, 1.54-1.44 (2 m, 4H, 3, 4), 1.21 ppm (d, J=6.2 Hz, 3H, 6); ¹³C NMR ([D₆]DMSO): δ = 173.00, 171.18, 158.98, 144.60, 144.20, 134.76, 134.50, 129.55, 122.05, 96.08, 91.59, 54.95, 51.14, 46.97, 38.36, 34.28, 33.40, 32.64, 25.93, 20.62, 20.17 ppm; IR (ATR): $\tilde{v} =$ 3376, 3246, 3086, 2993, 2961, 2927, 1736, 1632, 1612, 1574, 1520, 1458, 1423, 1386, 1219, 1203, 1172, 1161, 1052, 992, 817, 788, 751, 720, 676 cm⁻¹; MS (ESI): *m/z*: 388.3 $[M+1]^+$; elemental analysis calcd (%) for C₂₁H₂₉N₃O₄ (387.47): C 65.09, H 7.54, N 10.84; found: C 65.35, H 7.33, N 10.99.

Methyl 5-({4-[(6-methoxyquinolin-8-yl)amino]pentyl}carbamoyl)pent-

anoate (2 d): Method A, from the reaction of monomethyl adipate (1 d, 0.224 g) and after purification by column chromatography (dichloromethane/methanol 9.5:0.5) and crystallization (ether/petroleum ether), **2d** was obtained as a pale-yellow solid (0.483 g, 86%): m.p. 56–58 °C; ¹H NMR ([D_6]DMSO): δ =8.54–8.52 (dd, J=4.2, 1.6 Hz, 1H, 10), 8.09–8.06 (dd, J=8.3, 1.5 Hz, 1H, 12), 7.78 (t, J=5.4 Hz, 1H, 1), 7.45–7.40 (m, 1H, 11), 6.48 (d, J=2.4 Hz, 1H, 16), 6.26 (d, J=2.4 Hz, 1H, 14), 6.12 (d, J=8.8 Hz, 1H, 7), 3.82 (s, 3H, 17), 3.66–3.59 (m, 1H, 5), 3.57 (s, 3H, 7'), 3.08–3.04 (m, 2H, 2), 2.28 (t, J=6.9 Hz, 2H, 2'), 2.04 (t, J=6.9 Hz, 2H, 5'), 1.68–1.57, 1.57–1.42 (2m, 8H, 3, 4, 3', 4'), 1.20 ppm (d, J=6.3 Hz, 3H, 6); ¹³C NMR ([D_6]DMSO): δ =173.22, 171.60, 159.00, 144.63, 144.23, 134.79, 134.53, 129.57, 122.09, 96.09, 91.58, 54.97, 51.15, 46.98, 38.36, 35.00, 33.43, 32.98, 25.99, 24.73, 24.03, 20.18 ppm; IR (ATR): $\tilde{\nu}$ = 3373, 3311, 2943, 2876, 1730, 1640, 1618, 1518, 1508, 1461, 1425, 1389, 1265, 1222, 1169, 1055, 821, 680, 625 cm⁻¹; MS (ESI): *m/z*: 402.3 [*M*+1]⁺; elemental analysis calcd (%) for C₂₂H₃₁N₃O₄ (401.50): C 65.81, H 7.78, N 10.47; found: C 65.66, H, 7.66, N 10.31.

Methyl 4-({4-[(6-methoxyquinolin-8-yl)amino]pentyl}carbamoyl)benzoate (2e): Method A, from the reaction of monomethyl terephtalate (1 e, 0.252 g) and after purification by column chromatography (dichloromethane/methanol 9.5:0.5) and crystallization (ether), 2e was obtained as a pale-yellow solid (0.489 g, 83%): m.p. 112-113 °C; ¹H NMR ([D₆]DMSO): $\delta = 8.65$ (t, J = 5.5 Hz, 1 H, 1), 8.54–8.52 (dd, J=4.2, 1.6 Hz, 1 H, 10), 8.08-8.06 (dd, J=8.2, 1.1 Hz, 1 H, 12), 8.01 (d, J=8.3 Hz, 2H, 3', 7'), 7.94 (d, J=8.3 Hz, 2H, 4', 6'), 7.43-7.41 (m, 1 H, 11), 6.47 (d, J=2.2 Hz, 1 H, 16), 6.28 (d, J=2.1 Hz, 1 H, 14), 6.15 (d, J=8.7 Hz, 1 H, 7), 3.88 (s, 3 H, 9'), 3.81 (s, 3 H, 17), 3.69-3.65 (m, 1H, 5), 3.32-3.29 (m, 2H, 2), 1.76-1.65, 1.65-1.57 (2m, 4H, 3, 4), 1.23 ppm (d, J=6.3 Hz, 3 H, 6); ^{13}C NMR ([D_6]DMSO): $\delta\!=$ 165.69, 165.24, 158.97, 144.59, 144.18, 138.76, 134.73. 134.51, 131.54, 129.53, 128.99, 127.47, 122.03, 96.09, 91.59, 54.91, 52.26, 47.00, 39.27, 33.39, 25.79, 20.15 ppm; IR (ATR): $\tilde{v} =$ 3403, 3318, 3060, 2936, 1714, 1634, 1615, 1519, 1425, 1386, 1278, 1225, 1169, 1111, 820, 791, 736, 703 cm⁻¹; MS (ESI): *m/z*: 422.2 [*M*+1]⁺; elemental analysis calcd (%) for $C_{24}H_{27}N_3O_4$ (421.49): C 68.39, H 6.46, N 9.97; found: C 68.25, H 6.76, N 10.08.

General procedure for the preparation of carboxylic acids 3 a-e: A solution of lithium hydroxide monohydrate (0.126 g, 3 mmol) in water (10 mL) was added to a solution of ester **2** (6 mmol) in methanol (10 mL). The mixture was stirred at room temperature for 1 h. Methanol was evaporated under reduced pressure, and the aqueous residue was neutralized with 10% HCl and extracted with dichloromethane (3×). The organic layer was dried with sodium sulfate, filtered, and concentrated under reduced pressure.

3-({4-[(6-Methoxyquinolin-8-yl)amino]pentyl}carbamoyl)propanoic acid (3 a): From the reaction of ester 2 a (0.224 g) and after crystallization (ether), 3a was obtained as a pale-yellow solid (0.209 g, 97%): m.p. 146–148°C; ¹H NMR ([D₆]DMSO): δ = 12.06 (s, 1 H, 5'), 8.55-8.53 (dd, J=4.2, 1.6 Hz, 1 H, 10), 8.09-8.06 (dd, J=8.3, 1.6 Hz, 1 H, 12), 7.85–7.82 (t, J=5.4 Hz, 1 H, 1), 7.45–7.41(m, 1 H, 11), 6.48 (d, J=2.4 Hz, 1 H, 16), 6.26 (d, J=2.4 Hz, 1 H, 14), 6.12 (d, J=8.7 Hz, 1H, 7), 3.82 (s, 3H, 17), 3.66-3.57 (m, 1H, 5), 3.08-3.03 (m, 2H, 2), 2.41 (t, J=6.5 Hz, 2H, 2'), 2.29 (t, J=6.7 Hz, 2H, 3'), 1.70-1.58, 1.58–1.38 (2m, 4H, 3, 4), 1.2 ppm (d, J=6.3 Hz, 3H, 6); ¹³C NMR ([D₆]DMSO): $\delta = 173.87$, 170.73, 159.00, 144.63, 144.24, 134.80, 134.52, 129.58, 122.10, 96.10, 91.59, 54.97, 46.99, 38.48, 33.38, 30.04, 29.21, 25.96, 20.19 ppm; IR (ATR): \tilde{v} = 3454, 3268, 3094, 3009, 2962, 2928, 2863, 1713, 1649, 1615,1584, 1562, 1524, 1452, 1386, 1346, 1227, 1203, 1156, 1161, 824, 785, 746, 674 cm⁻¹; MS (ESI): *m*/ *z*: 360.2 $[M+1]^+$; elemental analysis calcd (%) for C₁₉H₂₅N₃O₄ (359.42): C 63.49, H 7.01, N 11.69; found: C 63.22, H 7.30, N 11.45.

(2*E*)-3-({4-[(6-Methoxyquinolin-8-yl)amino]pentyl}carbamoyl) prop-2enoic acid (**3b**): From the reaction of ester **2b** (0.231 g) and after

Chemistry Open 2010 , 7, 024 – 030	ChemistryOpen	2018,	7, 624 -	638
---	---------------	-------	----------	-----





crystallization (ether), **3 b** was obtained as a pale-yellow solid (0.178 g, 83%): m.p. 147–149°C; ¹H NMR ([D₆]DMSO): δ = 12.83 (s, 1H, 5'), 8.55–8.53 (dd, *J* = 1.6, 4.2 Hz, 1 H, 10), 8.49 (t, *J* = 5.5 Hz, 1 H, 1), 8.09–8.06 (dd, *J* = 1.5, 8.3 Hz, 1 H, 12), 7.45–7.40 (m, 1 H, 11), 6.91 (d, *J* = 15.5 Hz, 1 H, 2'), 6.50 (d, *J* = 15.5 Hz, 1 H, 3'), 6.47 (d, *J* = 2.4 Hz, 1 H, 16), 6.27 (d, *J* = 2.4 Hz, 1 H, 14), 6.14 (d, *J* = 8.7 Hz, 1 H, 7), 3.82 (s, 3 H, 17), 3.68–3.58 (m, 1 H, 5), 3.22–3.16 (m, 2 H, 2), 1.71–1.49 (m, 4 H, 3, 4), 1.21 ppm (d, *J* = 6.3 Hz, 3 H, 6); ¹³C NMR ([D₆]DMSO): δ = 166.49, 162.95, 159.00, 144.62, 144.22, 137.11, 134.79, 134.52, 129.57, 129.33, 122.09, 96.13, 91.62, 54.96, 46.96, 38.80, 33.40, 25.71, 20.19 ppm; IR (ATR): $\tilde{\nu}$ = 3453, 3271, 3087, 2937, 1716, 1654, 1613, 1564, 1527, 1380, 1337, 1269, 1236, 1213, 1198, 1171, 981, 909, 820, 785, 767, 673 cm⁻¹; MS (ESI): *m/z*: 358.2 [*M*+1]⁺; elemental analysis calcd (%) for C₁₉H₂₃N₃O₄ (357.40): C 63.85, H 6.49, N 11.76; found: C 63.77, H 6.72, N 11.99.

4-({4-[(6-Methoxyquinolin-8-yl)amino]pentyl}carbamoyl)butanoic

acid (3 c): From the reaction of ester 2 c (0.232 g) and after crystallization (ether), 3c was obtained as a pale-yellow solid (0.186 g, 88%): m.p. 95–96 °C; ¹H NMR ([D₆]DMSO): $\delta = 12.00$ (s, 1 H, 6'), 8.55-8.53 (dd, J=4.2, 1.6 Hz, 1 H, 10), 8.10-8.07 (dd, J=8.3, 1.5 Hz, 1 H, 12), 7.80 (t, J=5.4 Hz, 1 H, 1), 7.45–7.41 (m, 1 H, 11), 6.48 (d, J= 2.4 Hz, 1 H, 16), 6.27 (d, J=2.4 Hz, 1 H, 14), 6.13 (d, J=8.7 Hz, 1 H, 7), 3.83 (s, 3 H, 17), 3.69-3.55 (m, 1 H, 5), 3.09-3.05 (m, 2 H, 2), 2.20 (t, J=7.4 Hz, 2H, 2'), 2.08 (t, J=7.4 Hz, 2H, 4'), 1.75-1.43 (m, 6H, 3', 3, 4), 1.21 ppm (d, J=6.3 Hz, 3 H, 6); 13 C NMR ([D₆]DMSO): $\delta =$ $147.15, \ 171.36, \ 159.00, \ 144.63, \ 144.23, \ 134.79, \ 134.53, \ 129.57,$ 122.09, 96.11, 91.60, 54.97, 46.98, 38.38, 34.46, 33.41, 33.04, 25.97, 20.71, 20.19 ppm; IR (ATR): \tilde{v} = 3453, 3324, 2933, 1720, 1643, 1613, 1580, 1524, 1386, 1227, 1204, 1161, 1139, 1058, 822, 786, 675 cm⁻¹; MS (ESI): m/z: 374.2 $[M+1]^+$; elemental analysis calcd (%) for $C_{20}H_{27}N_{3}O_{4}$ (373.45): C 64.32, H 7.29, N 11.25; found: C 64.21, H 7.15, N 11.48.

5-({4-[(6-Methoxyquinolin-8-yl)amino]pentyl}carbamoyl)pentanoic

acid (3d): From the reaction of ester 2d (0.241 g) and after crystallization (ether), 3d was obtained as a pale-yellow solid (0.198 g, 85%): m.p. 68–69 °C; ¹H NMR ([D₆]DMSO): δ = 11.99 (s, 1 H, 7'), 8.54 (d, J=2.9 Hz, 1 H, 10), 8.08 (d, J=8.1 Hz, 1 H, 12), 7.78 (t, J=6.0 Hz, 2H, 1), 7.45-7.40 (m, 1H, 11), 6.48 (d, J=3.0 Hz, 1H, 16), 6.26 (d, J=3.0 Hz, 1 H, 14), 6.12 (d, J=8.6 Hz, 1 H, 7), 3.82 (s, 3 H, 17), 3.69-3.56 (m, 1H, 5), 3.09-3.0 (m, 2H, 2), 2.18 (t, J=6.0 Hz, 2H, 5'), 2.04 (t, J=6.0 Hz, 2H, 2'), 1.70-1.57, 1.57-1.42 (2m, 8H, 3, 4, 3', 4'), 1.20 ppm (d, J = 6.2 Hz, 3 H, 6); ¹³C NMR ([D₆]DMSO): $\delta = 174.37$, 171.67, 159.00, 144.63, 144.23, 134.79, 134.52, 129.57, 122.09, 96.10, 91.59, 54.97, 46.98, 38.36, 35.11, 33.38, 25.99, 24.84, 24.13, 20.19 ppm; IR (ATR): v = 3453, 3370, 3294, 2943, 2863, 1731, 1609, 1562, 1519, 1455, 1424, 1385, 1224, 1203, 1166, 1156, 1133, 1053, 819, 790, 765, 677 cm⁻¹; MS (ESI): *m/z*: 388.2 [*M*+1]⁺; elemental analysis calcd (%) for $C_{21}H_{29}N_3O_4$ (387.47): C 65.10, H 7.54, N 10.84; found: C 65.44, H 7.79, N 10.63.

4-({4-[(6-Methoxyquinolin-8-yl)amino]pentyl}carbamoyl)benzoic

acid (**3** e): From the reaction of ester **2** e (0.253 g) and after crystallization (ether), **3** e was obtained as a pale-yellow solid (0.210 g, 86%): m.p. 142–144°C; ¹H NMR ([D₆]DMSO): δ = 13.12 (s, 1H, 9'), 8.64 (t, *J* = 5.5 Hz, 1H, 1), 8.54–8.53 (dd, *J* = 4.2, 1.6 Hz, 1H, 10), 8.09–8.06 (dd, *J* = 8.3, 1.5 Hz, 1H, 12), 8.00 (d, *J* = 8.4 Hz, 2H, 3', 7'), 7.92 (d, *J* = 8.4 Hz, 2H, 4', 6'), 7.45–7.41 (m, 1H, 11), 6.47 (d, *J* = 2.4 Hz, 1H, 16), 6.28 (d, *J* = 2.4 Hz, 1H, 14), 6.16 (d, *J* = 8.4 Hz, 1H, 7), 3.81 (s, 3H, 17), 3.71–3.63 (m, 1H, 5), 3.34–3.28 (m, 2H, 2), 1.76–1.57 (m, 4H, 3, 4), 1.22 ppm (d, *J* = 7.3 Hz, 3H, 6); ¹³C NMR ([D₆]DMSO): δ = 166.80, 165.42, 159.00, 144.61, 144.22, 138.45, 134.80, 134.52, 132.79, 129.58, 129.16, 127.36, 122.09, 96.14, 91.59, 54.96, 47.02, 39.28, 33.40, 25.86, 20.19 ppm; IR (ATR): $\tilde{\nu}$ = 3413,

3308, 2941, 2864 1727, 1691, 1636, 1613, 1575, 1519, 1454, 1422, 1385, 1276, 1220, 1201, 1156, 818, 786, 674 cm⁻¹; MS (ESI): *m/z*: 408.2 $[M+1]^+$; elemental analysis calcd (%) for C₂₃H₂₅N₃O₄ (407.46): C 67.80, H 6.18, N 10.31; found: C 67.93, H 6.02, N 10.68.

General procedure for the preparation of *O*-benzylhydroxamic acids **4a–e**: A solution of acid **3** (0.6 mmol), DIEA (0.155 g, 1.2 mol), and HATU (0.228 g, 0.6 mmol) in dichloromethane (6 mL) was stirred at room temperature. After 10 min, *O*-benzylhydroxylamine hydrochloride (0.112 g, 0.7 mmol) and Et₃N (0.071 g, 0.7 mmol) were added. The mixture was stirred at room temperature for 2 h and concentrated under reduced pressure. The residue was dissolved in ethyl acetate (20 mL) and extracted with brine (3×). The organic layer was dried with sodium sulfate, filtered, and concentrated under reduced pressure.

N-(Benzyloxy)-N'-{4-[(6-methoxyquinolin-8-yl)amino]pentyl}butanediamide (4a): From the reaction of 3a (0.216 g) and after purification by column chromatography (dichloromethane/methanol 9.5:0.5) and crystallization (ether), 4a was obtained as a paleyellow solid (0.111 g, 40%): m.p. 121-123°C; ¹H NMR ([D₆]DMSO): $\delta =$ 10.99 (s, 1 H, 5'), 8.54 (d, J=2.7 Hz, 1 H, 10), 8.08 (d, J=7.2 Hz, 1 H, 12), 7.85 (s, 1 H, 1), 7.44-7-35 (m, 6 H, 11, 8'-12'), 6.47 (d, J =2.2 Hz, 1 H, 16), 6.26 (d, J=2.0 Hz, 1 H, 14), 6.12 (d, J=8.6 Hz, 1 H, 7), 4.76 (s, 2H, 6'), 3.82 (s, 3H, 17), 3.66-3.57 (m, 1H, 5), 3.08-3.02 (m, 2H, 2), 2.30 (t, J=6.9 Hz, 2H, 3'), 2.18 (t, J=6.8 Hz, 2H, 2'), 1.70–1.57, 1.57–1.43 (2 m, 4 H, 3, 4), 1.20 ppm (d, J=6.2 Hz, 3 H, 6); ^{13}C NMR ([D_6]DMSO): $\delta\!=\!170.71,\;168.80,\;159.00,\;144.62,\;144.23,$ 136.06, 134.79, 134.52, 129.57, 128.72, 128.26, 128.16, 122.09, 96.09, 91.59, 76.75, 54.97, 46.99, 38.49, 33.39, 30.37, 27.91, 25.97, 20.19 ppm; IR (ATR): v = 3367, 3311, 3060, 2936, 2862, 1728, 1636, 1615, 1546, 1519, 1457, 1424, 1387, 1223, 1200, 1159, 1052, 820, 790, 679 cm⁻¹; MS (ESI): m/z: 465.3 $[M+1]^+$; elemental analysis calcd (%) for $C_{26}H_{32}N_4O_4$ (464.56): C 67.22, H 6.94, N 12.06; found: C 67.46, H 6.69, N, 12.31.

(2E)-N-(Benzyloxy)-N'-{4-[(6-methoxyquinolin-8-yl)amino]pentyl}-

but-2-ene-diamide (4b): From the reaction of 3b (0.215 g) and after purification by column chromatography (dichloromethane/ methanol/ethyl acetate/cyclohexane 9.5:0.5:10:10) and crystallization (ether), **4b** was obtained as a pale-yellow solid (0.119 g, 43%): m.p. 157–158 °C; ¹H NMR ([D₆]DMSO): $\delta = 11.52$ (s, 1H, 5'), 8.55– 8.53 (dd, J=1.5, 4.2 Hz, 1 H, 10), 8.45 (t, J=5.5 Hz, 1 H, 1), 8.09-8.06 (dd, J = 1.4, 8.3 Hz, 1 H, 12), 7.45–7.37 (m, 1 H, 11, 8'-12'), 6.90 (d, J =15.1 Hz, 1 H, 2'), 6.62 (d, J=15.2 Hz, 1 H, 3'), 6.47 (d, J=2.4 Hz, 1 H, 16), 6.27 (d, J=2.3 Hz, 1 H, 14), 6.15 (d, J=8.7 Hz, 1 H, 7), 4.86 (s, 2H, 6'), 3.82 (s, 3H, 17), 3.68-3.58 (m, 2H, 5), 3.20-3.15 (m, 2H, 2), 1.70–1.49 (m, 4H, 3,4), 1.21 ppm (d, J=6.2 Hz, 3H, 6); ¹³C NMR ([D₆]DMSO): $\delta = 163.17$, 161.22, 158.99, 144.60, 144.22, 134.82, 134.49, 133.58, 129.58, 129.07, 122.09, 96.16, 91.64, 63.35, 54.97, 46.97, 38.95, 33.42, 25.76, 20.19 ppm; IR (ATR): $\tilde{v} =$ 3383, 3283, 3213, 3008, 2936, 2865, 1738, 1626, 1575, 1556, 1518, 1454, 1386, 1337, 1219, 1203, 1157,1051, 974, 818, 790, 738, 695, 675 cm⁻¹; MS (ESI): m/z: 463.1 $[M+1]^+$; elemental analysis calcd (%) for $C_{26}H_{30}N_4O_4$ (462.54): C 67.51, H 6.54, N 12.11; found: C 67.79, H 6.81, N 11.95.

N-(Benzyloxy)-*N*'-{4-[(6-methoxyquinolin-8-yl)amino]pentyl}pentanediamide (**4c**): From the reaction of **3c** (0.224 g) and after purification by column chromatography (dichloromethane/methanol 9.5:0.5) and crystallization (ether/petroleum ether), **4c** was obtained as a pale-yellow solid (0.144 g, 50%): m.p. 84–85 °C; ¹H NMR ([D₆]DMSO): δ = 10.95 (s, 1H, 6'), 8.54–8.53 (dd, *J*=4.1, 1.5 Hz, 1H, 10), 8.09–8.06 (dd, *J*=8.3, 1.4 Hz, 1H, 12), 7.78 (t, *J*=5.4 Hz, 1H, 1), 7.44–7.33 (m, 6H, 11, 9'-13'), 6.47 (d, *J*=2.4 Hz, 1H, 16), 6.26 (d, *J*=

ChemistryOpen 2018, 7, 624-638





2.3 Hz, 1 H, 14), 6.13 (d, J=8.8 Hz, 1 H, 7), 4.77 (s, 2 H, 7'), 3.82 (s, 3 H, 17), 3.62–3.56 (m, 1 H, 5), 3.08–3.04 (m, 2 H, 2), 2.04 (t, J=7.4 Hz, 2 H, 4'), 1.95 (t, J=7.3 Hz, 2 H, 2'), 1.74–1.43 (m, 6 H, 3', 3, 4), 1.20 ppm (d, J=6.2 Hz, 3 H, 6); ¹³C NMR ([D₆]DMSO): δ =171.32, 169.03, 159.00, 144.62, 144.23, 136.09, 134.79, 134.52, 129.57, 128.72, 128.26, 128.16, 122.09, 96.10, 91.59, 76.77, 54.97, 46.98, 38.41, 34.57, 33.43, 31.70, 25.99, 21.21, 20.19 ppm; IR (ATR): $\tilde{\nu}$ =3367, 3325, 3187, 2960, 2931, 2857, 1738, 1677, 1634, 1614, 1518, 1455, 1387, 1219, 1203, 1170, 1157, 1052, 1033, 821, 790, 740, 696, 676 cm⁻¹; MS (ESI): m/z: 479.3 [M+1]⁺; elemental analysis calcd (%) for C₂₇H₃₄N₄O₄ (478.58): C 67.76, H 7.16, N 11.71; found: C 67.59, H 6.98, N, 12.04.

N-(Benzyloxy)-*N*'-{4-[(6-methoxyquinolin-8-yl)amino]pentyl}hexanediamide (4d): From the reaction of 3d (0.232 g) and after purification by column chromatography (dichloromethane/methanol 9.5:0.5) and crystallization (ether), 4d was obtained as a paleyellow solid (0.151 g, 51%): m.p. 102-105°C; ¹H NMR ([D₆]DMSO): $\delta = 10.91$ (s, 1 H, 7'), 8.53 (dd, J = 4.0, 1.4 Hz, 1 H, 10), 8.06 (dd, J =8.2 Hz, 1 H, 12), 7.74 (s, 1 H, 1), 7.43-7.41 (m, 1 H, 11), 7.37-7.34 (m, 5 H, 10'-14'), 6.47 (d, J=2.3 Hz, 1 H, 16), 6.26 (d, J=2.2 Hz, 1 H, 14), 6.11 (d, J=8.7 Hz, 1 H, 7), 4.77 (s, 2 H, 8'), 3.82 (s, 3 H, 17), 3.65-3.59 (m, 1 H, 5), 3.08-3.02 (m, 2 H, 2), 2.02 (s, 2 H, 2'), 1.93 (s, 2 H, 5'), 1.68-1.62, 1.56-1.41 (2m, 8H, 3, 4, 3', 4'), 1.20 ppm (d, J=6.3 Hz, 3 H, 6); 13 C NMR ([D₆]DMSO): δ = 171.61, 169.19, 158.96, 144.59, 144.17, 136.03, 134.72, 134.48, 129.52, 128.66, 128.19, 129.09, 122.02, 96.06, 91.60, 76.70, 54.93, 53.57, 46.97, 38.34, 35.13, 33.42, 32.04, 25.94, 24.83, 24.62, 20.15 ppm; IR (ATR): $\tilde{\nu} =$ 3383, 3287, 2934, 2865, 1737, 1635, 1578, 1519, 1457, 1423, 1387, 1224, 1202, 1168, 1052, 820, 791, 743, 696 cm⁻¹; MS (ESI): *m/z*: 493.3 [*M*+1]⁺; elemental analysis calcd (%) for $C_{28}H_{36}N_4O_4$ (492.61): C 68.27, H 7.37, N 11.37; found: C 68.47, H 7.25, N, 11.49.

 N^{1} -(Benzyloxy)- N^{4} -{4-[(6-methoxyquinolin-8-yl)amino]pentyl}-benzene-1,4-dicarboxamide (4e): From the reaction of 3e (0.245 g) and after purification by column chromatography (dichloromethane/methanol 9.5:0.5) and crystallization (ether), 4e was obtained as a pale-yellow solid (0.188 g, 61%): m.p. 183-184°C; ¹H NMR ([D₆]DMSO): $\delta = 11.88$ (s, 1 H, 9'), 8.59 (t, J = 5.5 Hz, 1 H, 1), 8.54-8.52 (dd, J=4.2, 1.6 Hz, 1 H, 10), 8.09-8.06 (dd, J=8.3, 1.6 Hz, 1 H, 12), 7.89 (d, J=8.4 Hz, 2 H, 3', 7'), 7.80 (d, J=8.4 Hz, 2 H, 4', 6'), 7.45-7.34 (m, 6H, 11, 12'-16'), 6.47 (d, J=2.5 Hz, 1H, 16), 6.28 (d, J=2.4 Hz, 1 H, 14), 6.15 (d, J=8.6 Hz, 1 H, 7), 4.94 (s, 2 H, 10'), 3.81 (s, 3H, 17), 3.71-3.63 (m, 1H, 5), 3.33-3.27 (m, 2H, 2), 1.75-1.58 (m, 4H, 3, 4), 1.23 ppm (d, J = 6.3 Hz, 3H, 6); ¹³C NMR ([D₆]DMSO): $\delta =$ 165.35, 163.70, 158.99, 144.61, 144.22, 137.30, 135.82, 134.80, 134.52, 134.38, 129.57, 128.93, 128.32, 127.32, 127.02, 122.09, 96.13, 91.59, 77.03, 54.96, 47.02, 39.23, 33.39, 25.88, 20.19 ppm; IR (ATR): $\tilde{v} = 3384$, 3287, 2969, 2935, 1738, 1630, 1577, 1519, 1492, 1387, 1320, 1224, 1159, 1053, 863, 820, 790, 747, 718, 69 cm⁻¹; MS (ESI): 513.3 $[M+1]^+$; elemental analysis calcd (%) for $C_{30}H_{32}N_4O_4$ (512.60): C 70.29, H 6.29, N 10.93; found: C 70.55, H 6.03, N, 11.21.

General procedure for the preparation of SAHAquines **5** a–d: A suspension of *O*-benzylhydroxamic acid **4** (0.27 mmol) and 10% Pd/C (20 mg) in methanol (7 mL) was stirred at room temperature for 2–4 h under a hydrogen atmosphere. The catalyst was filtered off, and the mother liquor was concentrated under reduced pressure.

N-Hydroxy-*N*'-{4-[(6-methoxyquinolin-8-yl)amino]pentyl}butanediamide (**5a**): From the reaction of **4a** (0.125 g) and after crystallization (ether), **5a** was obtained as a pale-yellow solid (0.077 g, 76%): m.p. 109–110°C; ¹H NMR ([D₆]DMSO): δ = 10.37 (s, 1 H, 5'), 8.67 (s, 1 H, 6'), 8.54–8.53 (dd, *J*=4.2, 1.6 Hz, 1 H, 10), 8.09–8.06 (dd, *J*=8.3, 1.5 Hz, 1 H, 12), 7.85 (t, *J*=5.3 Hz, 1 H, 1), 7.45–7.41 (m, 1 H, 11), 6.47 (d, J=2.4 Hz, 1H, 16), 6.26 (d, J=2.4 Hz, 1H, 14), 6.12 (d, J=8.7 Hz, 1H, 7), 3.82 (s, 3H, 17), 3.67–3.56 (m, 1H, 5), 3.09–3.00 (m, 2H, 2), 2.29 (t, J=6.5 Hz, 2H, 2'), 2.17 (t, J=6.5 Hz, 2H, 3'), 1.72–1.56, 1.56–1.39 (2m, 4H, 3, 4), 1.20 ppm (d, J=6.3 Hz, 3H, 6); ¹³C NMR ([D₆]DMSO): δ =170.84, 168.46, 159.00, 144.63, 144.24, 134.80, 134.52, 129.58, 122.11, 96.10, 91.60, 54.98, 47.00, 38.49, 33.42, 30.64, 27.92, 25.97, 20.20 ppm; IR (ATR): $\tilde{\nu}$ =3471, 3375, 3283, 3204, 3008, 2964, 2934, 1744, 1647, 1612, 1556, 1521, 1455, 1385, 1366, 1226, 1205, 1172, 1157, 1056, 821, 789, 677 cm⁻¹; MS (ESI): m/z: 375.2 $[M+1]^+$; elemental analysis calcd (%) for C₁₉H₂₆N₄O₄ (374.43): C 60.95, H 7.00, N 14.96; found: C 60.84, H 6.81, N, 15.17.

N-Hydroxy-N'-{4-[(6-methoxyquinolin-8-yl)amino]pentyl}pentanediamide (5b): From the reaction of 4c (0.129 g) and after crystallization (ether), 5b was obtained as a pale-yellow solid (0.077 g, 73%): m.p. 99–102 °C; ¹H NMR ([D₆]DMSO): $\delta = 10.32$ (s, 1 H, 6'), 8.63 (s, 1 H, 7'), 8.55-8.52 (dd, J=4.2, 1.6 Hz, 1 H, 12), 8.09-8.05 (dd, J=8.3, 1.6 Hz, 1 H, 12), 7.76 (t, J=5.3 Hz, 1 H, 1), 7.45-7.39 (m, 1 H, 11), 6.47 (d, J=2.4 Hz, 1 H, 16), 6.26 (d, J=2.4 Hz, 1 H, 14), 6.11 (d, J=8.7 Hz, 1 H, 7), 3.82 (s, 3 H, 17), 3.68–3.56 (m, 1 H, 5), 3.05 (d, J=5.6 Hz, 2 H, 2), 2.04 (t, J=7.4 Hz, 2H, 2'), 1.94 (t, J=7.4 Hz, 2H, 4'), 1.76-1.57, 1.57-1.40 (2m, 6H, 3, 4, 3'), 1.21 ppm (d, J=6.3 Hz, 3H, 6); ¹³C NMR ([D₆]DMSO): $\delta = 171.37$, 168.74, 158.98, 144.60, 144.20, 134.75, 134.50, 129.54, 122.05, 96.08, 91.61, 54.95, 46.98, 38.39, 34.73, 33.43, 31.75, 25.95, 21.42, 20.17 ppm; IR (ATR): $\tilde{\nu} = 3468$, 3375, 3283, 3210, 3008, 2963, 2933, 1744, 1645, 1613, 1520, 1455, 1386, 1204, 1171, 1158, 1056, 821, 789, 677 cm⁻¹; MS (ESI): *m/z*: 389.2 [*M*+1]⁺; elemental analysis calcd (%) for C₂₀H₂₈N₄O₄ (388.46): C 61.84, H 7.27, N 14.42; found: C 61.69, H 7.03, N, 14.77.

N-Hydroxy-*N*'-{4-[(6-methoxyquinolin-8-yl)amino]pentyl}hexanediamide (5 c): From the reaction of 4d (0.133 g) and after crystallization (ether), 5c was obtained as a pale-yellow solid (0.082 g, 75%): m.p. 109–111 °C; ¹H NMR ([D₆]DMSO): $\delta = 10.34$ (s, 1 H, 7'), 8.66 (s, 1 H, 8'), 8.55-8.52 (dd, J=4.2, 1.5 Hz, 1 H, 10), 8.10-8.05 (dd, J=8.3, 1.4 Hz, 1 H, 12), 7.77 (t, J=5.3 Hz, 1 H, 1), 7.45-7.40 (m, 1 H, 11), 6.47 (d, J=2.4 Hz, 1 H, 16), 6.26 (d, J=2.4 Hz, 1 H, 14), 6.12 (d, J=8.8 Hz, 1 H, 7), 3.82 (s, 3 H, 17), 3.67–3.57 (m, 1 H, 5), 3.04 (d, J=5.5 Hz, 2 H, 2), 2.03 (s, 2H, 2'), 1.92 (s, 2H, 5'), 1.72-1.58, 1.58-1.41 (2m, 8H, 3, 4, 3', 4'), 1.20 ppm (d, J=6.3 Hz, 3 H, 6); 13 C NMR ([D₆]DMSO): $\delta =$ 171.70, 168.95, 159.00, 144.62, 144.24, 134.80, 134.52, 129.57, 122.10, 96.10, 91.60, 54.98, 46.98, 38.38, 35.20, 33.44, 32.11, 26.01, 24.97, 24.87, 20.20 ppm; IR (ATR): $\tilde{\nu} = 3428$, 3283, 3191, 3088, 3044, 2943, 2923, 2856, 1738, 1621, 1578,1556, 1523, 1385, 1367, 1204, 1170, 1160, 954, 822, 788, 676 cm⁻¹; MS (ESI): *m/z*: 403.3 [*M*+1]⁺; elemental analysis calcd (%) for C₂₁H₃₀N₄O₄ (402.49): C 62.67, H 7.51, N 13.92; found: C 62.98, H 7.36, N, 14.09.

*N*¹-Hydroxy-*N*⁴-{4-[(6-methoxyquinolin-8-yl)amino]pentyl}benzene-

1,4-dicarboxamide (**5 d**): From the reaction of **4e** (0.138 g) and after purification by column chromatography (dichloromethane/ methanol 9:1) and crystallization (ether), **5 d** was obtained as a pale-yellow solid (0.048 g, 42%): m.p. 156–158°C; ¹H NMR ([D₆]DMSO): δ = 11.32 (s, 1H, 9'), 9.12 (s, 1H, 10'), 8.57 (t, *J*=5.5 Hz, 1H, 1), 8.54–8.53 (dd, *J*=4.2, 1.6 Hz, 1H, 10), 8.09–8.06 (d, *J*=8.3, 1.6 Hz, 1H, 12), 7.88 (d, *J*=8.3 Hz, 2H, 3', 7'), 7.80 (d, *J*=8.3 Hz, 2H, 4', 6'), 7.44–7.40 (m, 1H, 11), 6.47 (d, *J*=2.2 Hz, 1H, 16), 6.28 (d, *J*=2.1 Hz, 1H, 14), 6.15 (d, *J*=8.7 Hz, 1H, 7), 3.81 (s, 3H, 17), 3.71–3.62 (m, 1H, 5), 3.33–3.27 (m, 2H, 2), 1.75–1.58 (m, 4H, 3, 4), 1.23 ppm (d, *J*=6.2 Hz, 3H, 6); ¹³C NMR ([D₆]DMSO): δ =165.44, 163.49, 159.00, 144.62, 144.20, 136.91, 134.86, 134.79, 134.53, 129.57, 127.17, 126.78, 122.09, 96.12, 91.60, 54.96, 47.02, 39.23, 33.41, 25.89, 20.19 ppm; IR (ATR): \tilde{v} =3427, 3303, 3199, 2969, 2938, 1738, 1672, 1636, 1616, 1521, 1458, 1388, 1223, 1203, 1172, 1015, 898,





858, 821, 789, 677 cm⁻¹; MS (ESI): m/z: 423.1 $[M+1]^+$; elemental analysis calcd (%) for C₂₃H₂₆N₄O₄ (422.48): C 65.39, H 6.20, N 13.26; found: C 65.47, H 6.12, N, 12.99.

General procedure for the preparation of *O*-methylhydroxamic acids **6a–e**: A solution of corresponding acid **3** (0.2 mmol), DIEA (0.052 g, 0.4 mmol), and HATU (0.076 g, 0.2 mmol) in dichloromethane (5 mL) was stirred at room temperature. After 10 min, *O*-methylhydroxylamine hydrochloride (0.020 g, 0.24 mmol) and Et₃N (0.024 g, 0.24 mmol) were added. The mixture was stirred at room temperature for 2 h and was then concentrated under reduced pressure. The residue was dissolved in ethyl acetate (8 mL) and extracted with water (3 ×). The organic layer was dried with sodium sulfate, filtered, and concentrated under reduced pressure.

N-Methoxy-N'-{4-[(6-methoxyquinolin-8-yl)amino]pentyl}butanediamide (6a): From the reaction of 3a (0.072 g) and after purification by column chromatography (dichloromethane/methanol 9.5:0.5) and crystallization (ether/petroleum ether), 6a was obtained as a pale-yellow solid (0.055 g, 71 %): m.p. 121-122 °C; ¹H NMR ([D₆]DMSO): $\delta = 10.93$ (s, 1 H, 5'), 8.54–8.53 (dd, J = 3.0 Hz, 1 H, 10), 8.06 (d, J=8.2 Hz, 1 H, 12), 7.82 (s, 1 H, 1), 7.43-7.41 (m, 1 H, 11), 6.47 (d, J=2.2 Hz, 1 H, 16), 6.26 (d, J=2.1 Hz, 1 H, 14), 6.11 (d, J= 8.7 Hz, 1 H, 7), 3.83 (s, 3 H, 17), 3.63-3.58 (m, 1 H, 5), 3.54 (s, 3 H, 6'), 3.08-3.02 (m, 2H, 2), 2.29 (t, J=7.4 Hz, 2H, 2'), 2.16 (t, J=7.0 Hz, 2H, 3'), 1.67–1.63, 1.55–1.45 (2m, 4H, 3, 4), 1.21 ppm (d, J=6.3 Hz, 3H, 6); $^{13}{\rm C}$ NMR ([D₆]DMSO): $\delta\!=\!170.64$, 168.44, 158.96, 144.59, 144.17, 134.71, 134.48, 129.52, 122.01, 96.04, 91.60, 62.97, 54.92, 46.97, 38.44, 33.37, 30.27, 27.84, 25.90, 20.14 ppm; IR (ATR): $\tilde{\nu} =$ 3302, 3234, 3001, 2969, 2933, 1739, 1658, 1634, 1556, 1519, 1457, 1424, 1388, 1336, 1224, 1204, 1168, 1053, 820, 791, 679 cm⁻¹; MS (ESI): m/z: 389.2 $[M+1]^+$; elemental analysis calcd (%) for $C_{20}H_{28}N_4O_4$ (388.46): C 61.84, H 7.27, N 14.42; found: C 62.05, H 7.01, N 14.65.

(2E)-N-Methoxy-N'-{4-[(6-methoxyquinolin-8-yl)amino]pent-yl}but-2enediamide (6b): From the reaction of 3b (0.072 g) and after purification by column chromatography (cyclohexane/ethyl acetate/ methanol 3:1:0.5) and crystallization (ether), 6b was obtained as a pale-yellow solid (0.038 g, 49%): m.p. 177-179°C; ¹H NMR ([D₆]DMSO): $\delta = 11.53$ (s, 1 H, 5'), 8.55–8.53 (dd, J = 1.6, 4.2 Hz, 1 H, 10), 8.44 (t, J=5.5 Hz, 1 H, 1), 8.09-8.06 (dd, J=1.5, 8.3 Hz, 1 H, 12), 7.45–7.41 (m, 6H, 11), 6.89 (d, J = 15.1 Hz, 1H, 2'), 6.60 (d, J =15.1 Hz, 1 H, 3'), 6.47 (d, J=2.2 Hz, 1 H, 16), 6.27 (d, J=2.2 Hz, 1 H, 14), 6.14 (d, J=8.7 Hz, 1 H, 7), 3.82 (s, 3 H, 17), 3.65 (s, 4 H, 5, 6'), 3.20–3.14 (m, 2H, 2), 1.70–1.49 (m, 4H, 3, 4), 1.21 ppm (d, J =6.2 Hz, 3 H, 6); ¹³C NMR ([D₆]DMSO): $\delta = 163.16$, 160.97, 159.00, 144.60, 144.22, 135.72, 134.82, 134.49, 133.58, 129.58, 129.12, $128.85,\ 129.34,\ 122.09,\ 96.16,\ 91.63,\ 76.97,\ 54.97,\ 46.97,\ 38.77,$ 33.42, 25.77, 20.19 ppm; IR (ATR): $\tilde{\nu} =$ 3285, 3223, 3097, 3007, 2969,2935, 1738, 1630, 1570, 1519, 1457, 1388, 1338, 1224, 1168, 1066, 1053, 991, 821, 792, 659 cm⁻¹; MS (ESI): *m/z*: 387.1 [*M*+1]⁺; elemental analysis calcd (%) for $C_{20}H_{26}N_4O_4$ (386.44): C 62.16, H 6.78, N 14.50; found: C 61.94, H 6.55, N 14.87.

N-Methoxy-*N*'-{4-[(6-methoxyquinolin-8-yl)amino]pentyl}pentanediamide (**6c**): From the reaction of **3c** (0.075 g) and after crystallization (ether), **6c** was obtained as a pale-yellow solid (0.045 g, 56%): m.p. 112–113 °C; ¹H NMR ([D₆]DMSO): δ =10.94 (s, 1H, 6'), 8.55–8.53 (dd, J=4.2, 1.5 Hz, 1H, 10), 8.09–8.06 (dd, J=8.3, 1.4 Hz, 1H, 12), 7.89 (t, J=5.3 Hz, 1H, 1), 7.45–7.41 (m, 1H, 11), 6.48 (d, J=2.4 Hz, 1H, 16), 6.26 (d, J=2.3 Hz, 1H, 14), 6.12 (d, J=8.7 Hz, 1H, 7), 3.82 (s, 3H, 17), 3.62 (m, 1H, 5), 3.56 (s, 3H, 7'), 3.06–3.04 (m, 2H, 2), 2.04 (t, J=7.3 Hz, 2H, 2'), 1.93 (t, J=7.3 Hz, 2H, 4'), 1.74–1.40 (m, 6H, 3', 3,4'), 1.20 ppm (d, J=6.3 Hz, 3H, 6); ¹³C NMR *N*-Methoxy-*N*'-{4-[(6-methoxyquinolin-8-yl)amino]pentyl}hexanediamide (6d): From the reaction of 3d (0.075 g) and after purification by column chromatography (dichloromethane/methanol 9.5:0.5) and crystallization (ether), 6d was obtained as a pale-yellow solid (0.042 g, 51 %): m.p. 108–109 $^{\circ}\text{C};~^{1}\text{H}$ NMR ([D_6]DMSO): $\delta\!=\!10.91$ (s, 1 H, 7'), 8.55-8.52 (dd, J=4.1, 1.6 Hz, 1 H, 10), 8.09-8.06 (dd, J=8.3, 1.5 Hz, 1 H, 12), 7.75 (t, J=5.4 Hz, 1 H, 1), 7.44-7.41 (m, 1 H, 11), 6.47 (d, J=2.4 Hz, 1 H, 16), 6.26 (d, J=2.4 Hz, 1 H, 14), 6.11 (d, J=8.7 Hz, 1 H, 7), 3.82 (s, 3 H, 17), 3.65-3.55 (m, 4 H, 5, 8'), 3.08-3.00 (m, 2 H, 2), 2.03 (s, 2 H, 2'), 1.92 (s, 2 H, 5'), 1.55–1.49, 1.49–1.40 (m, 8 H, 3, 4, 3', 4'), 1.20 ppm (d, J=6.3 Hz, 3 H, 6); $^{13}{\rm C}$ NMR ([D_6]DMSO): $\delta\!=$ 171.64, 168.91, 158.98, 144.61, 144.20, 134.76, 134.50, 129.55, 122.06, 96.08, 91.60, 63.05, 54.95, 46.97, 38.36, 35.15, 33.43, 32.07, 25.97, 24.86, 24.60, 20.17 ppm; IR (ATR): $\tilde{v} =$ 3395, 3287, 3166, 3090, 2927, 2861, 1739, 1674, 1631, 1577, 1563, 1521, 1457, 1423, 1388, 1227, 1206, 1170, 1161, 1053, 821, 790, 679 cm⁻¹; MS (ESI): *m/z*: 417.3 $[M+1]^+$; elemental analysis calcd (%) for C₂₂H₃₂N₄O₄ (416.51): C 63.44, H 7.74, N 13.45; found: C 63.26, H 7.53, N 13.71.

*N*¹-Methoxy-*N*⁴-{4-[(6-methoxyquinolin-8-yl)amino]pentyl}benzene-1,4-dicarboxamide (6e): From the reaction of 3e (0.082 g) and after crystallization (ether), 6e was obtained as a pale-yellow solid (0.040 g, 45%): m.p. 142–145 °C; ¹H NMR ([D₆]DMSO): $\delta = 11.84$ (s, 1 H, 9'), 8.58 (t, J=5.56 Hz, 1 H, 1), 8.54–8.53 (dd, J=1.6, 4.2 Hz, 1 H, 10), 8.08–8.06 (dd, J=1.6, 8.3 Hz, 1 H, 12), 7.89 (d, J=8.4 Hz, 2 H, 3', 7'), 7.80 (d, J = 8.4 Hz, 2 H, 4', 6'), 7.43–7.41 (m, 1 H, 11), 6.47 (d, J =2.5 Hz, 1 H, 16), 6.28 (d, J=2.4 Hz, 1 H, 14), 6.15 (d, J=8.8 Hz, 1 H, 7), 3.82 (s, 3H, 17), 3.72 (s, 2H, 10'), 3.69-3.65 (m, 1H, 5), 3.31-3.28 (m, 2 H, 2), 1.76–1.66, 1.65–1.56 (m, 4 H, 3, 4), 1.23 ppm (d, J =6.3 Hz, 3 H, 6); ¹³C NMR ([D₆]DMSO): $\delta = 165.32$, 163.36, 158.98, 144.61, 144.20, 137.28, 134.76, 134.52, 134.31, 129.55, 127.21, 126.93), 122.06, 96.10, 91.60, 63.26, 54.94, 47.01, 39.23, 33.41, 25.85, 20.17 ppm; IR (ATR): $\tilde{\nu} = 3394$, 3309, 3176, 3060, 2970, 2937, 1738, 1670, 1624, 1557, 1520, 1456, 1389, 1221, 1205, 1161, 1039, 867, 821, 791, 678 cm⁻¹; MS (ESI): *m/z*: 437.3 [*M*+1]⁺; elemental analysis calcd (%) for $C_{24}H_{28}N_4O_4$ (436.50): C 66.04, H 6.47, N 12.84; found: C 66.33, H 6.31, N 13.04.

Biological Evaluation

Cell viability: All synthesized compounds were first screened for their inhibition of mitochondrial metabolic activity by using the MTT assay.^[65] Mitochondrial metabolic activity correlates with cell viability. The following cell lines were tested: U2OS, HepG2, MCF-7, and Hek293 cells. The compounds were dissolved in DMSO (3×10^{-3} M) and stored at -20 °C. The cells were seeded at a density of 3000 cells per 200 µL in a 96-well plate (Sarsted) in triplicate and grown in Dulbecco's modified Eagle medium with 4500 mg L⁻¹ glucose (DMEM-high glucose) (Lonza) containing 10% fetal bovine serum (FBS) (Gibco), 100 UmL⁻¹ penicillin, and 100 µg mL⁻¹ streptomycin (Sigma–Aldrich). The next day, the medium was aspirated and cells were treated for 72 h. Only the compounds that led to more than a 40% reduction in mitochondrial metabolic activity at a concentration of 5×10^{-5} M were selected for further analysis. The following concentrations of selected compounds were used: $5 \times$

ChemistryOpen 2018, 7, 624-638



10⁻⁵, 1×10⁻⁵, 1×10⁻⁶, 1×10⁻⁷, and 1×10⁻⁸ M. All working concentrations were freshly prepared in DMEM on the day of the testing. A fresh growth medium was added to untreated control cells, which were defined as 100% viable. DMSO (1.67, 0.33, 0.03, 0.3×10⁻², and 0.3×10⁻³%) in DMEM was considered as a negative control. Cisplatin, PQ, and SAHA were used as positive controls. After 72 h of incubation in the presence of selected compounds, media were aspirated and MTT (0.5 mg mL⁻¹; 40 µL per well) was added, and the cells were incubated for 4 h 37 °C. Media were aspirated and formazan crystals were dissolved in DMSO (170 µL per well). The absorbance was measured on a microplate reader (Promega) at λ = 560 nm. The IC₅₀ values (concentration required to decrease viability by 50%) were calculated by using nonlinear regression on the sigmoidal dose–response plots and are expressed as mean ± SD of two independent experiments.

MCF-7 cell labeling with Hoechst 33342: Fluorescent dye Hoechst 33342 was used to determine the total number of cells. MCF-7 cells were seeded at a density of 3000 cells per 100 μ L (72 h treatment) or 6000 cells per 100 μ L (24 h treatment) in a 96-well plate (Corning Inc.). Cells were grown in triplicate in DMEM containing 10% FBS (Wisent), 100 UmL⁻¹ penicillin, and 100 μ g mL⁻¹ streptomycin (Thermo Fischer). The next day, cells were treated with SA-HAquines **4e** and **5b**, SAHA, PQ, and cisplatin in ten different concentrations ranging from 5×10⁻⁵ to 1×10⁻⁹ M. After 72 h, cells were stained with Hoechst 33342 (10 μ M, 10 min). Cells were then imaged by using a fluorescence microscope (Leica, DMI4000B). Results were calculated by using nonlinear regression on the sigmoidal dose–response plot and are expressed as mean ± SD of two independent experiments.

In vitro drug sensitivity assay against erythrocytic stages of *P. falciparum*: The antiplasmodial activity of compounds **2–6** was tested in a drug-sensitivity assay against two laboratory *P. falciparum* strains (3D7—chloroquine sensitive, Dd2—chloroquine-resistant) as described before by using the histidine-rich protein 2 (HRP2) assay.^[66,67] In brief, 96-well plates were precoated with the tested compounds in a threefold dilution before ring-stage parasites were added in complete culture medium at a hematocrit of 1.5% and a parasitaemia of 0.05%. After 3 days of incubation at 37 °C, 5% CO₂ and 5% oxygen, plates were frozen until analyzed by HRP2-ELISA. All compounds were evaluated in duplicate in at least two independent experiments. The IC₅₀ was determined by analyzing the nonlinear regression of log concentration–response curves using the drc-package v0.9.0 of R v2.6.1.^[68]

In vitro activity against P. berghei hepatic stages: Compound activity on P. berghei infection of a human hepatoma cell line (HuH7) was assessed employing the luminescence-based method, as previously described.^[69] Briefly, hepatic infection was determined by measuring the luminescence intensity of lysates of HuH7 cells infected with a firefly luciferase-expressing P. berghei line. HuH7 cells $(1.0 \times 10^4 \text{ per well})$ were seeded in 96-well plates the day before infection. One hour prior to infection, the medium was replaced by medium containing the appropriate drug concentrations. The addition of 1.0×10^4 sporozoites was followed by centrifugation at $1800 \times q$ for 5 min and parasite infection load was measured 48 h after parasite addition by a bioluminescence assay (Biotium, USA) using a multiplate reader Infinite M200 (Tecan, Switzerland). The effect of the different treatments on the viability of HuH7 cells was assessed by the CellTiter-Blue assay (Promega, USA) according to the manufacturer's protocol. Nonlinear regression analysis was employed to fit the normalized results of the dose-response curves, and IC₅₀ values were determined using GraphPad Prism V5.0.



Immunocytochemistry for histone acetylation: MCF-7 cells were seeded at a density of 5000 cells per coverslip in DMEM containing 10% FBS, 100 U mL $^{-1}$ penicillin, and 100 $\mu g\,mL^{-1}$ streptomycin and were left for 24 h to adhere. Cells were treated with 4e, 5b, SAHA, or PQ (at 1 µm concentration) for 24 h. Following treatment, cells were fixed with 4% paraformaldehyde (10 min), then permeabilized using 0.1% Triton X-100 (10 min). Blocking was performed using 10% goat serum in phosphate buffer saline (PBS) (1 h), followed by incubation with primary antibody (rabbit anti-acetyl-Histone-H3 polyclonal antibody 1/500; Millipore 06-559) for 24 h in a humidified chamber at 4 $^{\circ}$ C. Samples were washed with PBS (3 \times 5 min). Samples were then incubated with secondary antibody (Alexa Fluor 647 goat anti-rabbit IgG -2 mg mL^{-1} 1/500; Life Technologies A21244) for 1 h in the dark, after which they were washed with PBS (3×5 min). Nuclei were labeled with Hoechst 33342 (10 µм, 10 min). Samples were mounted on microscope slides using Poly Aqua Mount (PolySciences) and were dried overnight before imaging with a fluorescence microscope (Leica). Fluorescence intensity was guantified using ImageJ software, and the results are presented as means of fluorescence per cell \pm SD (as fold increase in the untreated control = 1).

Acknowledgements

Support for this study by Croatian Science Foundation (project number IP-09–2014-1501), Canadian Institute of Health Research (CIHR; project number MOP-119425), and National Sciences and Engineering Research Council of Canada (NSERC; project number RGPIN 04994-15) is acknowledged.

Conflict of Interest

The authors declare no conflict of interest.

Keywords: acetylation \cdot anticancer agents \cdot antiplasmodial activity \cdot cytostatic activity \cdot drug design

- [1] A. Remesh, Int. J. Basic Clin. Pharmacol. 2012, 1, 2–12.
- [2] D. B. Longley, P. G. Johnston, J. Pathol. 2005, 205, 275-292.
- [3] N. J. Curtin, Nat. Rev. 2012, 12, 801-817.
- [4] C. Azad, M. Saxena, A. J. Siddiqui, J. Bhardwaj, S. K. Puri, G. P. Dutta, N. Anand, A. K. Saxena, *Chem. Biol. Drug Des.* 2017, 90, 254–261.
- [5] C. Teixeira, N. Vale, B. Pérez, A. Gomes, J. R. Gomes, P. Gomes, Chem. Rev. 2014, 114, 11164–11220.
- [6] D. Shahinas, A. Folefoc, D. R. Pillai, Pathogens 2013, 2, 33-54.
- [7] D. Agarwal, R. D. Gupta, S. K. Awasthi, Antimicrob. Agents Chemother. 2017, 61, e00249-17.
- [8] K. Takasu, T. Shomogama, C. Saiin, H.-S. Kim, Y. Wataya, R. Brun, M. Ihara, Chem. Pharm. Bull. 2005, 53, 653–661.
- [9] A. Gomes, M. Machado, L. Lobo, F. Nogueira, M. Prudencio, C. Teixeira, P. Gomes, *ChemMedChem* 2015, 10, 1344–1349.
- [10] S. Higginbotham, W. R. Wong, R. G. Linington, C. Spadafora, L. Iturrado, A. E. Arnold, *PLoS One* 2014, 9, e84549.
- [11] W. T. Johnston, N. Mutalima, D. Sun, B. Emmanuel, K. Bhatia, P. Aka, X. Wu, E. Borgstein, G. N. Liomba, S. Kamiza, N. Mkandawire, M. Batumba, L. M. Carpenter, H. Jaffe, E. M. Molyneux, J. J. Goedert, D. Soppet, R. Newton, S. M. Mbulaiteye, *Sci. Rep.* **2014**, *4*, 3741.
- [12] K. H. Khan, Germs 2013, 3, 26-35.
- [13] D. A. Fedosov, M. Dao, G. E. Karniadakis, S. Suresh, Ann. Biomed. Eng. 2014, 42, 368-387.
- [14] R. Hooft van Huijsduijnen, R. K. Guy, K. Chibale, R. K. Haynes, I. Peitz, G. Kelter, M. A. Phillips, J. L. Vennerstrom, Y. Yuthavong, T. N. Wells, *PLoS One* **2013**, *8*, e82962.

^{© 2018} The Authors. Published by Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim





- [15] K.-C. Keum, N.-C. Yoo, W.-M. Yoo, K. K. Chang, Y. N. Choon, Y. W. Min, WO 2002013826 A1, 2002.
- [16] F. Liu, Y. Shang, S-Z. Chen, Acta Pharmacol. Sin. 2014, 35, 645-652.
- [17] A. Kamal, A. Aziz, S. Shouman, E. El-Demerdash, M. Elgendy, A. B. Abdel-Naim, Sci. Proc. 2014, 1, e384.
- [18] A. Ganguli, D. Choudhury, S. Datta, S. Bhattacharya, G. Chakrabarti, *Bio-chimie* 2014, 107, 338–349.
- [19] G. W. Soo, J. H. Law, E. Kan, S. Y. Tan, W. Y. Lim, G. Chay, N. I. Bukhari, I. Segarra, Anticancer Drugs 2010, 21, 695–703.
- [20] Y. K. Wong, C. Xu, K. A. Kalesh, Y. He, Q. Lin, W. S. F. Wong, H. M. Shen, J. Wang, *Med. Res. Rev.* 2017, *37*, 1492–1517.
- [21] I. Nakase, H. Lai, N. P. Singh, T. Sasaki, Int. J. Pharm. 2008, 354, 28-33.
- [22] C. Cheng, T. Wang, Z. Song, L. Peng, M. Gao, O. Hermine, S. Rousseaux,
 S. Khochbin, J. Q. Mi, J. Wang, *Cancer Med.* **2018**, *7*, 380–396.
- [23] T. Shi, X. X. Yu, L. J. Yan, H. T. Xiao, Cancer Chemother. Pharmacol. 2017, 79, 287–294.
- [24] G. Džimbeg, B. Zorc, M. Kralj, K. Ester, K. Pavelić, J. Balzarini, E. De Clercq, M. Mintas, *Eur. J. Med. Chem.* 2008, 43, 1180–1187.
- [25] M. Šimunović, I. Perković, B. Zorc, K. Ester, M. Kralj, D. Hadjipavlou-Litina, E. Pontiki, *Bioorg. Med. Chem.* 2009, 17, 5605 – 5613.
- [26] I. Perković, S. Tršinar, J. Žanetić, M. Kralj, I. Martin-Kleiner, J. Balzarini, D. Hadjipavlou-Litina, A. M. Katsori, B. Zorc, J. Enzyme Inhib. Med. Chem. 2013, 28, 601–610.
- [27] K. Pavić, I. Perković, M. Cindrić, M. Pranjić, I. Martin-Kleiner, M. Kralj, D. Schols, D. Hadjipavlou-Litina, A.-M. Katsori, B. Zorc, *Eur. J. Med. Chem.* 2014, 86, 502–514.
- [28] I. Perković, M. Antunović, I. Marijanović, K. Pavić, K. Ester, M. Kralj, J. Vlainić, I. Kosalec, D. Schols, D. Hadjipavlou-Litina, E. Pontiki, B. Zorc, *Eur. J. Med. Chem.* **2016**, *124*, 622–636.
- [29] K. Pavić, I. Perković, P. Gilja, F. Kozlina, K. Ester, M. Kralj, D. Schols, D. Hadjipavlou-Litina, E. Pontiki, B. Zorc, *Molecules* 2016, *21*, 1629–1653.
- [30] K. Pavić, I. Perković, Š. Pospíšilová, M. Machado, D. Fontinha, M. Prudêncio, J. Jampilek, A. Coffey, L. Endersen, H. Rimac, B. Zorc, *Eur. J. Med. Chem.* 2018, 143, 769–779.
- [31] A. Gomes, I. Fernandes, C. Teixeira, N. Mateus, M. J. Sottomayor, P. Gomes, ChemMedChem 2016, 11, 2703–2712.
- [32] C. Pérez, I. Fernandes, N. Mateus, C. Teixeira, P. Gomes, *Bioorg. Med. Chem. Lett.* 2013, 23, 6769–6772.
- [33] S. Slezakova, J. Ruda-Kucerova, Anticancer Res. 2017, 37, 5995-6003.
- [34] Q. Sun, J. Wang, Y. Li, J. Zhuang, Q. Zhang, X. Sun, D. Sun, Chem. Biol. Drug Des. 2017, 90, 1019–1028.
- [35] B. S. Pybus, S. R. Marcsisin, X. Jin, G. Deye, J. C. Sousa, Q. Li, D. Caridha, Q. Zeng, G. A. Reichard, C. Ockenhouse, J. Bennett, L. A. Walker, C. Ohrt, V. Melendez, *Malar. J.* 2013, *12*, e212.
- [36] N. Vale, R. Moreira, P. Gomes, *Eur. J. Med. Chem.* **2009**, 44, 937–953.
- [37] S. O. Tcherniuk, O. Chesnokova, I. V. Oleinikov, A. I. Potopalsky, A. V. Oleinikov, *Malar. J.* 2015, 14, 425.
- [38] A. Wildbolz, Ther. Umsch. 1973, 30, 218-222.
- [39] B. Pouvelle, P. J. Farley, C. A. Long, T. F. Taraschi, J. Clin. Invest. 1994, 94, 413–417.
- [40] E. A. Usanga, FEBS Lett. **1986**, 209, 23–27.
- [41] L. Nair, V. K. Bhasin, Jpn. J. Med. Sci. Biol. 1994, 47, 241-252.
- [42] A. Kreidenweiss, P.G. Kremsner, B. Mordmüller, Malar. J. 2008, 7, 187.
- [43] S. Grant, C. Easley, P. Kirkpatrick, Nat. Rev. Drug. Discov. 2007, 6, 21-22.
- [44] M. J. Chua, M. S. J. Arnold, W. Xu, J. Lancelot, S. Lamotte, G. F. Späth, E. Prina, R. J. Pierce, D. P. Fairlie, T. S. Skinner-Adams, K. T. Andrews, *Int. J. Parasitol.* 2017, 7, 42–50.

- [45] K. T. Andrews, A. Haque, M. K. Jones, Immunol. Cell Biol. 2012, 90, 66– 77.
- [46] B. K. Chaal, A. P. Gupta, B. D. Wastuwidyaningtyas, Y. H. Luah, Z. Bozdech, PLoS Pathog. 2010, 6, e1000737.
- [47] H. Hu, A. Cabrera, M. Kono, S. Mok, B. K. Chaal, S. Haase, K. Engelberg, S. Cheemadan, T. Spielmann, P. R. Preiser, T.-W. Gillberger, Z. Bozdech, *Nat. Biotechnol.* 2010, 28, 91–98.
- [48] K. T. Andrews, T. N. Tran, N. C. Wheatley, D. P. Fairlie, Curr. Top. Med. Chem. 2009, 9, 292–308.
- [49] J. A. Engel, A. J. Jones, V. M. Avery, S. D. Sumanadasa, S. S. Ng, D. P. Fairlie, T. Skinner-Adams, K. T. Andrews, *Int. J. Parasitol.* **2015**, *5*, 117–126.
- [50] K. Trenholme, L. Marek, S. Duffy, G. Pradel, G. Fisher, F. K. Hansen, T. S. Skinner-Adams, A. Butterworth, C. J. Ngwa, J. Moecking, C. D. Goodman, G. I. McFadden, S. D. M. Sumanadasa, D. P. Fairlie, V. M. Avery, T. Kurz, K. T. Andrews, *Antimicrob. Agents Chemother.* **2014**, *58*, 3666–3678.
- [51] F. W. Muregi, A. Ishih, Drug Dev. Res. 2010, 71, 20-32.
- [52] R. Oliveira, D. Miranda, J. Magalhaes, R. Capela, M. J. Perry, P. M. O'Neill, R. Moreira, F. Lopes, *Bioorg. Med. Chem.* 2015, *23*, 5120–5130.
- [53] S. Shaveta, S. Mishra, P. Singh, *Eur. J. Med. Chem.* 2016, *124*, 500-536.
 [54] M. Mishra, V. K. Mishra, V. Kashaw, A. K. Iyer, S. K. Kashaw, *Eur. J. Med. Chem.* 2017, *125*, 1300-1320.
- [55] X. Zhang, J. Zhang, L. Tong, Y. Luo, S. Mingbo, Y. Zang, J. Li, W. Lu, Y. Chen, *Bioorg. Med. Chem.* 2013, 21, 3240–3244.
- [56] J. E. Payne, C. Bonnefous, C. A. Hassig, K. T. Symons, X. Guo, P. M. Nguyen, T. Annable, P. L. Wash, T. Z. Hoffman, T. S. Rao, A. K. Shiau, J. W. Malecha, S. A. Noble, J. F. Hager, N. D. Smith, *Bioorg. Med. Chem. Lett.* 2008, 18, 6093–6096.
- [57] A. S. Madsen, H. M. Kristensen, G. Lanz, C. A. Olsen, ChemMedChem 2014, 9, 614–626.
- [58] B. Zorc, Z. Rajić Džolić, I. Butula, Croat. Chem. Acta 2012, 85, 595-602.
- [59] Chemicalize, 2017, ChemAxon Ltd. Available from: http://www.chemicalize.org.
- [60] E. L. Luzina, A. V. Popov, Eur. J. Med. Chem. 2012, 53, 364-373.
- [61] B. E. Lauffer, R. Mintzer, R. Fong, S. Mukund, C. Tam, I. Zilberleyb, B. Flicke, A. Ritscher, G. Fedorowicz, R. Vallero, D. F. Ortwine, J. Gunzner, Z. Modrusan, L. Neumann, C. M. Koth, P. J. Lupardus, J. S. Kaminker, C. E. Heise, P. Steiner, J. Biol. Chem. 2013, 288, 26926–26943.
- [62] R. Duffy, C. Wade, R. Chang, Drug Discovery Today 2012, 17, 942-953.
- [63] J. Held, A. Kreidenweiss, B. Mordmüller, Expert Opin. Drug Discovery 2013, 8, 1325 – 1337.
- [64] D. Barbaras, M. Kaiser, R. Brun, K. Gademann, *Bioorg. Med. Chem. Lett.* 2008, 18, 4413–4415.
- [65] M. R. Boyd, K. D. Paull, Drug Dev. Res. 1995, 34, 91-109.
- [66] J. Held, T. Gebru, M. Kalesse, R. Jansen, K. Gerth, R. Müller, B. Mordmüller, Antimicrob. Agents Chemother. 2014, 58, 6378–6384.
- [67] H. Noedl, J. Bronnert, K. Yingyuen, B. Attlmayr, H. Kollaritsch, M. Fukuda, Antimicrob. Agents Chemother. 2005, 49, 3575–3577.
- [68] R Core Team (2015). R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing, Vienna, Austria. https://www.R-project.org/.
- [69] A. M. Mendes, I. S. Albuquerque, M. Machado, J. Pissarra, P. Meireles, M. Prudêncio, Antimicrob. Agents Chemother. 2017, 61, e02005-16.

Received: June 21, 2018