This article was downloaded by: [Lulea University of Technology] On: 14 September 2013, At: 03:46 Publisher: Taylor & Francis Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Organic Preparations and Procedures International: The New Journal for Organic Synthesis

Publication details, including instructions for authors and subscription information: http://www.tandfonline.com/loi/uopp20

A NOVEL ROUTE TO THYMOL FROM m-CRESOL

K. J. Divakar $^{\rm a}$, V. V. Dhekne $^{\rm a}$, B. D. Kulkami $^{\rm a}$, P. L. Joshi $^{\rm a}$ & A. S. Rao $^{\rm a}$

^a Division of Organic Chemistry, (Synthesis) National Chemical Laboratory, Pune, 411008, INDIA Published online: 18 Feb 2009.

To cite this article: K. J. Divakar , V. V. Dhekne , B. D. Kulkami , P. L. Joshi & A. S. Rao (2000) A NOVEL ROUTE TO THYMOL FROM m-CRESOL, Organic Preparations and Procedures International: The New Journal for Organic Synthesis, 32:1, 92-94, DOI: <u>10.1080/00304940009356753</u>

To link to this article: <u>http://dx.doi.org/10.1080/00304940009356753</u>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <u>http://www.tandfonline.com/page/terms-and-conditions</u>

A NOVEL ROUTE TO THYMOL FROM m-CRESOL[†]

Submitted by K. J. Divakar, V.V. Dhekne, B. D. Kulkarni, P. L. Joshi, and A. S. Rao^{*} (10/04/99)

Division of Organic Chemistry (Synthesis) National Chemical Laboratory, Pune 411008, INDIA

Thymol 2, a naturally occurring phenolic terpene used as an antiseptic,¹ has been prepared by alkylation of *m*-cresol 1, with propylene in the presence of alumina (method A)² or aluminium 3-



methylphenoxide (method B).³ With alumina, **2** is formed in 63% yield (7 % of 4-isopropyl-3methylphenol as by-product) while method B gives **2** in 75% yield (13% of 2,6-diisopropyl-3methylphenol as by-product); however, both procedures require a purification step.



i) ethylacetoacetate, acidify, heat; *ii*) NaOH, heat / acidify; *iii*) heat; *iv*) H_2 , Raney Ni; *v*) NaOH, ethanediol, heat / acidify

We now report an improved procedure which passes through compound **5** which itself may be valuable for the preparation of naturally occurring compounds related to thymol.^{4,5}

4,7-Dimethylcoumarin (3) is a crystalline compound which may be obtained in good yield (71%) and high purity (>99%) from the Pechmann reaction with m-cresol. Fries and Fickewirth⁶ had prepared the unsaturated acid 4 by heating 3 with alkali followed by acidification and showed that on heating, acid 4 readily loses CO_2 to give phenol 5. It has been observed during the present study that heating the coumarin 3 with alkali at a high temperature (~190°) provides the phenol 5 (after acidification); evidently, the acid 4 underwent decarboxylation in alkaline medium. The hydroxyl group, situated *ortho* to the acid side-chain, is essential to facilitate decarboxylation since cinnamic acid itself and other substituted cinnamic acids do not undergo decarboxylation when heated with alkali at 190°.⁷ Hydrogenation of the phenol 5 furnished thymol 2 which is free from regioisomers; thus, the overall yield of 2, free from regioisomers, is 51% starting from 1.

EXPERIMENTAL SECTION

All mps and bps are uncorrected. IR spectra were recorded as liquid film or Nujol mull on Perkin-Elmer model 137B or 599B spectrophotometer. ¹H-NMR. spectra were recorded on Varian T-60 or WH-90 spectrometers. Microanalysis was carried out in the microanalytical section of our laboratory. GC was determined on Carlo-Erba: Fractovap No. 2450, Integrator: 3390A Hewlett Packard and Detector: F. I. D.

2-(2-Hydroxy-4-methylphenyl)-1-propene (5).- A mixture of 4,7-dimethylcoumarin **3** (87.0g, 0.5 mmol),^{6,8} NaOH (80.0g, 2.0 mmol) and 1,2-ethanediol (500mL) was heated under reflux under N₂ for 2 hrs, cooled, diluted with water (1.0L) and covered with ether (300mL). Dilute HCl (0.5N) was added with stirring till the pH was 2. The organic layer was separated and aqueous layer extracted with ether (200mL ×2). The combined ethereal extracts were washed with water and dried (Na₂SO₄). After removal of solvent, the residue was distilled *in vacuo* to yield 55g (74%) of **5** as a colorless liquid, bp. 130-135°/ 40mm; NMR (CCl₄): δ , 6.87-6.52 (m, 3H, ArH), 5.38 (s, 1H, OH), 5.27 (m, 1H, vinyl H), 5.05 (m, 1H, vinyl H), 2.27 (s, 3H, ArCH₃), 2.08 (m,3H, CH₃-C=C).

Anal. Calcd for C10H12O: C, 81.04; H, 8.16. Found: C, 80.86; H, 8.28

Thymol (2).- A mixture of phenol 5 (50.0g,), ethanol (500mL) and Raney nickel (10g) was stirred under hydrogen in an autoclave at 3-5 atmospheres at room temperature (30°) till hydrogen absorption ceased. The reaction mixture was filtered and the filtrate was evaporated *in vacuo*. Distillation of the residue furnished 50.0g. (98%) of 2 as a colorless liquid, bp. 135-140°/50mm; the IR and ¹H-NMR spectra were identical with those of an authentic sample. 1H-NMR and GLC studies show that thymol thus prepared contains less than detectable amounts (1%) of regioisomers of thymol. The distillate solidified on standing at room temperature. A sample recrystallized from ether/pet ether mixture showed mp. 50-51°, lit.⁶ mp. 51°.

REFERENCES

- [†] This work is covered by Indian Patent No. 146327; Chem. Abstr., 92, P58436n (1980).
- The Merck Index, p. 1604, 12th Edition, 1996; S. Budavari, Ed., Merck Research Laboratories, White House Station, NJ.
- 2. M. Nitta, K. Romura and K. Yamaguchi, Bull. Chem. Soc. Jpn, 47, 2360 (1974).
- 3. R. Stroh, R. Seydel and W. Hahn, Angew. Chem., 69, 699 (1957).
- 4. K. J. Divakar, B. D. Kulkarni and A. S. Rao, Indian J. Chem., 15B, 322 (1977).
- 5. K. J. Divakar, B. D. Kulkarni and A. S. Rao, *ibid.*, 15B, 849 (1977).
- 6. K. Fries and G. Fickewirth, Ber., 41, 367 (1908).

- 7. V. V. Dhekne, B. D. Kulkarni and A. S. Rao, Indian J. Chem., 15B, 755 (1977).
- 8. S. Sethna and R.Phadke, Org. React., 7, 1 (1953).

SYNTHESIS OF 4,6-DINITROINDOLE

 Submitted by (09/14/99)
 Vladimir V. Rozhkov*, Alexander M. Kuvshinov and Svyatoslav A. Shevelev

 N. D. Zelinsky Institute of Organic Chemistry Russian Academy of Sciences Leninsky prospekt 47, 117913, Moscow, RUSSIA

Nitroindoles represent an attractive synthetic target for the preparation of physiologically active compounds.^{1,2} To our surprise, 4,6-dinitroindole is not a known compound. The present paper describes the synthesis of 1 from 2,4,6-trinitrotoluene (TNT) according to the following scheme.



Although compounds 2, 3 and 4 have been described previously,^{6,7} they were not characterized by NMR spectroscopy. The assignment of protons H_a , H_b and H_c in 2,4,6-trinitrostyrene (4) has been made on the basis of the following NMR criteria: ${}^{3}J_{trans} > {}^{3}J_{cis} > {}^{2}J_{gem}$.⁸ The ortho-nitro group is substituted selectively upon interaction of styrene 4 with sodium azide in DMF at 20°. 2-Azido-4,6dinitrostyrene (5) is a labile compound which is converted to 1 even during crystallization from PrⁱOH. The structure of 1 was unambiguously determined by IR, ¹H and ¹³ C NMR spectroscopy, mass spectrometry and microanalysis.