

PII: S0957-4166(97)00144-4

## Enantioselective conjugate addition of diethylzinc to chalcone catalyzed by Co(acac)<sub>2</sub> and chiral amino alcohols

André H. M. de Vries and Ben L. Feringa\*

Department of Organic and Molecular Inorganic Chemistry, Groningen Center for Catalysis and Synthesis, University of Groningen, Nijenborgh 4, 9747 AG Groningen, The Netherlands

Abstract:  $Co(acac)_2$  in the presence of chiral ligands has been employed as catalyst for the enantioselective conjugate addition of diethylzinc to chalcone. With chiral amino alcohols derived from (+)-camphor, enantioselectivities up to 83% were achieved. © 1997 Elsevier Science Ltd

Conjugate addition reactions of carbon nucleophiles to  $\alpha,\beta$ -unsaturated compounds are among the most widely used methods for carbon–carbon bond formation in organic synthesis.<sup>1</sup> Asymmetric conjugate addition reactions, however, still lack the 'ideal chiral catalyst' which promotes enantioselective conjugate addition of readily available organometallic reagents to a wide variety of substrates.<sup>2</sup> Considerable progress has been made with the copper catalyzed conjugate addition of Grignard reagents<sup>3</sup> and the nickel catalyzed conjugate addition of diethylzinc.<sup>4</sup> Also a copper catalyzed 1,4addition of diethylzinc to cyclohexenone has been reported.<sup>5</sup> Recently, we and others have found the first catalytic system capable of the enantioselective conjugate addition of dialkylzinc reagents to *both* cyclic and acyclic substrates.<sup>5</sup>

Although transmetallation reactions of organozinc reagents to several metal salts have been proposed,<sup>6</sup> Knochel and co-workers recently reported the first synthetically useful transmetallations of organozinc reagents to *cobalt* in acylation and allylation reactions.<sup>7</sup> The use of organocobalt reagents in conjugate addition reactions has been limited.<sup>8</sup> Herein we describe catalytic enantioselective conjugate addition reactions of diethylzinc to chalcone using Co(acac)<sub>2</sub> and chiral amino alcohols (Equation 1).



When Co(acac)<sub>2</sub> (7 mol%) and chiral amino alcohol **3** (16 mol%) were employed in the model reaction the 1,4-product **2** was isolated in 61% yield (70% conversion according to GC analysis, after 16 h at  $-30^{\circ}$ C).<sup>9</sup> No 1,2-addition was observed and chiral HPLC analysis<sup>9</sup> revealed an enantioselectivity of 67%.

Three other chiral amino alcohol ligands 4–6 were also examined in the model reaction. With ligand 4 product 2 was isolated in 73% yield with an e.e. of 83%. Ligands  $5^{10}$  and 6 gave substantial lower conversions to the 1,4-product (~50%) and moderate e.e.'s (see Figure 1).

Using chiral amino alcohols the cobalt catalyzed conjugate addition is relatively slow compared to the nickel catalyzed reaction (>95% conversion to the 1,4-product 2 after 2 h at  $-30^{\circ}C)^{4c}$  and a considerable amount (5% with ligands 3 and 4, 20% with ligands 5 and 6) of the reduced by-product 1,3-diphenylpropan-1-one has been detected.<sup>11</sup> However, relatively high enantioselectivities were found in this reaction and moreover transmetallation processes of organozinc reagents to cobalt seem to have a greater applicability in (enantioselective) organic reactions<sup>7</sup> than transmetallation to

<sup>\*</sup> Corresponding author. E-mail: Feringa@chem.rug.nl



Figure 1. Chiral amino alcohols used in the  $Co(acac)_2$  catalyzed addition of  $Et_2Zn$  to 1. In brackets the e.e. of product 2 achieved with the given ligand.

other metal salts.<sup>12</sup> Therefore, further studies involving enantioselective cobalt mediated reactions of organozinc reagents are in progress in our laboratories.

## Acknowledgements

We are grateful to Prof. J. Martens, University of Oldenburg, for a gift of chiral ligand 5.

## References

- 1. Perlmutter, P. Conjugate Addition Reactions in Organic Synthesis Tetrahedron Organic Chemistry Series, No. 9, Pergamon, Oxford, 1992.
- a) Rossiter, B. E.; Swingle, N. M. Chem. Rev. 1992, 92, 771. b) Feringa, B. L.; de Vries, A. H. M. in Advances in Catalytic Processes, Vol 1: Asymmetric Chemical Transformations Doyle, M. P., Ed., JAI Press, Connecticut, 1995, p. 151.
- a) Zhou, Q.-L.; Pfaltz, A. *Tetrahedron* 1994, 50, 4467. b) van Klaveren, M.; Lambert, F.; Eijkelkamp, D. J. F. M.; Grove, D. M.; van Koten, G. *Tetrahedron Lett.* 1994, 35, 6135. c) Spescha, M.; Rihs, G. *Helv. Chim. Acta* 1993, 76, 1219. d) Kanai, M.; Tomioka, K. *Tetrahedron Lett.* 1995, 36, 4275.
- 4. a) Soai, K.; Hayasaka, T.; Ugajin, S. J. Chem. Soc., Chem. Commun. 1989, 516. b) Bolm, C.; Ewald, M.; Felder, M. Chem. Ber. 1992, 125, 1205. c) de Vries, A. H. M.; Jansen, J. F. G. A.; Feringa, B. L. Tetrahedron 1994, 50, 4479. d) Corma, A.; Iglesias, M.; Martín, V.; Rubio, J.; Sánchez, F. Tetrahedron: Asymmetry 1992, 3, 845.
- 5. a) Alexakis, A.; Frutos, J.; Mangeney, P. *Tetrahedron: Asymmetry* **1993**, *4*, 2427–2430. b) de Vries, A. H. M.; Meetsma, A.; Feringa, B. L. Angew. Chem., Int. Ed. Engl. **1996**, *35*, 2374.
- 6. a) Boersma, J. In Comprehensive Organometallic Chemistry Wilkinson, G.; Stone, F. G. A.; Abel, E. W., Eds., Pergamon, Oxford, 1982, Vol. 2, Chapter 16. b) Knochel, P.; Singer, R. D. Chem. Rev. 1993, 93, 2117.
- 7. a) Devasagayaraj, A.; Knochel, P. Tetrahedron Lett. 1995, 36, 8411. b) Reddy, C. K.; Knochel, P. Angew. Chem., Int. Ed. Engl. 1996, 35, 1700.
- a) Kauffmann, T.; Hopp, G.; Laarmann, B.; Stegemann, D.; Wingbermühle, D. Tetrahedron Lett. 1990, 31, 511. b) For a recent review on organocobalt reagents derived from alkyllithium or Grignard reagents, see: Kauffmann, T. Angew. Chem., Int. Ed. Engl. 1996, 35, 386.
- 9. For a typical experimental procedure, see: de Vries, A. H. M.; Imbos, R.; Feringa, B. L. Tetrahedron: Asymmetry 1997, 8, 1467.
- 10. Kindly provided by Prof. J. Martens, see: Behnen, W.; Mehler, T.; Martens, J. Tetrahedron: Asymmetry 1993, 4, 1413.
- Confirmed by GCMS analysis. Nickel catalyzed conjugate reduction of α,β-unsaturated ketones is a known process, see: Caporusso, A. M.; Giacomelli, G.; Lardicci, L. J. Org. Chem. 1982, 47, 4640.
- 12. With Pd(OAc)<sub>2</sub>, PdCl<sub>2</sub>, and Fe(acac)<sub>3</sub> as catalyst in the model reaction only traces of the 1,4product were determined.

(Received in UK 19 February 1997; accepted 25 March 1997)