

ChemComm

Accepted Manuscript



This article can be cited before page numbers have been issued, to do this please use: R. Menon, S. Undeela, S. Thadkapally, J. B. Nanubolu and K. K. Singarapu, *Chem. Commun.*, 2015, DOI: 10.1039/C5CC04871K.



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.



Journal Name

COMMUNICATION

Catalyst-controlled divergence in cycloisomerisation reactions of *N*-propargyl-*N*-vinyl sulfonamides: Gold-catalysed synthesis of 2-sulfonylmethyl pyrroles and dihydropyridines

Received 00th January 20xx,
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

www.rsc.org/


Sridhar Undeela,^a Srinivas Thadkapally,^a Jagadeesh Babu Nanubolu,^b Kiran Kumar Singarapu^c and Rajeev S. Menon^{*a}

Gold-catalysed, divergent synthesis of 2-sulfonylmethyl pyrroles and dihydropyridines from *N*-propargyl-*N*-vinyl sulfonamides has been achieved. Echavarren's gold(I) catalyst promoted the formation of pyrrole derivatives whereas the combination of PPh₃AuCl/AgSbF₆ afforded dihydropyridines. The aza-enyne precursors for the cycloisomerisation reaction were prepared by a base-mediated formal vinylic substitution reaction of 2-bromoallyl sulfones.

The development of suitable catalysts and conditions for promoting a desired reaction pathway of multifunctional substrates is one of the fundamental challenges of catalysis.¹ An additional level of selectivity may be contemplated wherein a common substrate can be transformed into more than one products by careful choice of catalysts/conditions. Such endeavors, commonly known as *divergent catalysis*, are highly attractive for the discovery of drugs and functional materials (Scheme 1a).² Although the benefits of divergent catalytic processes are obvious, their development is extremely challenging.³

Catalytic cycloisomerisation of enynes constitute a powerful, modern method for the construction of a variety of carbocycles and heterocycles.⁴ Among the various catalysts that are employed for enyne cycloisomerisation, gold complexes stand out owing to their ability to selectively activate alkynes in complex molecular settings.⁵ As a result, tremendous progress has been made in the recent years in this area.^{4,5a-b,6} Often, two or more cyclisation pathways are available for substituted enynes and a complex interplay of inherent substrate bias and catalysts dictate the outcome of the cycloisomerisation.⁷ While it is highly desirable, control over the available divergent pathways is realised rather rarely.⁸ Here we report gold-catalysed divergent cycloisomerisation reactions of *N*-

propargyl-*N*-vinyl sulfonamides **1** to afford 2-sulfonylmethyl pyrroles **2** or 2-sulfonylmethyl-1,2-dihydropyridines **3** (Scheme 1b). Remarkably, complete selectivity can be achieved by the choice of catalysts under otherwise identical conditions. Additionally, the cyclisation precursors were assembled via a novel, base-mediated formal vinylic substitution reaction of propargyl sulfonamides **4** and 2-bromoallyl sulfones **5** (Scheme 1b).



Scheme 1. a) Conceptual framework of divergent catalysis. b) Synthesis of aza-enynes and their divergent cycloisomerisation. [Au] = [JohnPhosAu(CH₃CN)]SbF₆. [Au-Ag] = Ph₃PAuCl/AgSbF₆.

Discovery of the above-described vinylic substitution has its origins in our recent investigations on the cyclocondensation reactions of unsaturated sulfones.⁹ We were interested in allenyl sulfones as potential building blocks for sulfone-containing cyclic systems. In view of the rich and versatile chemistry of allenates,¹⁰ allenyl sulfones appear to hold significant but largely untapped synthetic potential. The relative scarcity of studies involving allenyl sulfones may be attributed to their base-sensitivity and propensity to undergo nucleophile-triggered oligomerisation reactions.¹¹ Our efforts to circumvent this problem culminated in the development of a base-mediated formal vinylic substitution reaction of the 2-bromoallyl sulfones **5a-b**¹² and various sulfonamide nucleophiles **4a-h** to afford the *N*-propargyl-*N*-vinyl sulfonamide **1aa-hb** (Table 1). The intermediacy of 1-(phenylsulfonyl)allene was confirmed by further experiments (See supporting information for details).

^a Medicinal Chemistry and Pharmacology Division, CSIR-Indian Institute of Chemical Technology, Hyderabad, 500 007, India.

^b Centre for X-ray Crystallography, CSIR-Indian Institute of Chemical Technology, Hyderabad, 500 007, India.

^c Centre for Nuclear Magnetic Resonance, CSIR-Indian Institute of Chemical Technology, Hyderabad, 500 007, India.

Electronic Supplementary Information (ESI) available: [General experimental procedures, NMR data, and single crystal X-ray data for CCDC 1063242.] For ESI and crystallographic data, See DOI: 10.1039/x0xx00000x

COMMUNICATION

Journal Name

The *N*-propargyl-*N*-vinyl sulfonamides **1aa-hb** constitute an assortment of 3-aza-1,5-enynes with structural features that make them attractive substrates for catalytic processes for the synthesis of nitrogen heterocycles.¹³ The activation of the alkyne subunit by π -acids such as gold and silver catalysts was investigated with a view to promote cyclisation reactions. Screening of a handful of gold and silver catalysts revealed that two cyclisation pathways are operative leading to the formation of the pyrrole derivative **2aa** and the dihydropyridine product **3aa** (Table 2).

Table 1. Cs₂CO₃-mediated formal vinylic substitution of 2-bromoallyl sulfones **5a-b** for the synthesis of *N*-propargyl-*N*-vinyl sulfonamides **1aa-hb**.^a

Entry	Sulfonamides 4a-h	Bromoallyl sulfones 5a-b	Product	Yield ^a
1	4a , R ¹ = R ² = H, R ³ = Me	5a , R ⁴ = Ph	1aa	94%
2	4a	5b , R ⁴ = <i>p</i> -tolyl	1ab	89%
3	4b , R ¹ = R ² = H, R ³ = Ph	5a	1ba	96%
4	4b	5b	1bb	92%
5	4c , R ¹ = R ² = H, R ³ = <i>p</i> -tolyl	5a	1ca	93%
6	4c	5b	1cb	90%
7	4d , R ¹ = Ph, R ² = H, R ³ = Me	5a	1da	95%
8	4d	5b	1db	90%
9	4e , R ¹ = thiophene-2-yl, R ² = H, R ³ = Me	5a	1ea	82%
10	4e	5b	1eb	94%
11	4f , R ¹ = 2-bromophenyl, R ² = H, R ³ = Me	5a	1fa	93%
12	4f	5b	1fb	89%
13	4g , R ¹ = 4-methoxyphenyl, R ² = H, R ³ = Me	5a	1ga	88%
14	4g	5b	1gb	84%
15	4h , R ¹ = H, R ² = Ph, R ³ = Me	5b	1hb	55%

^aReaction conditions: **4a-g** (1 mmol), **5a-b** (1.2 mmol), Cs₂CO₃ (2.2 mmol), acetonitrile (5 mL), 25 °C, 4 h. ^bYields of products isolated after column chromatography. ^cReaction conditions **4h** (1 mmol), **5b** (1.2 mmol), NaH (2.2 mmol), THF (5 mL), 0 °C, 0.5 h and **1hb** obtained as a mixture of isomers (see supporting information for details).

Table 2. Optimisation of reaction conditions for cycloisomerisation of *N*-propargyl-*N*-vinyl sulfonamide **1aa**.^a

Entry	Catalysts (mol%)	Time (h)	Yield of 2aa ^b	Yield of 3aa ^b
1	AuCl ₃ (5)	3	78%	-
2	(JohnPhos)Au(CH ₃ CN)SbF ₆ (2)	1	89%	-
3	Ph ₃ PAuCl (5)	2	-	58%
4	AuCl ₃ (5) & AgSbF ₆ (5)	2	Traces	46%
5	(JohnPhos)Au(CH ₃ CN)SbF ₆ (2) & AgSbF ₆ (5)	1	-	48%
6	Ph ₃ PAuCl (2) & AgSbF ₆ (5)	1	-	92%
7	AgSbF ₆ (10)	4	-	66%
8 ^c	AgOTf (10)	2	-	-
9 ^d	No catalyst	24	-	-

^aReaction conditions: **1aa** (0.5 mmol), catalysts, CH₂Cl₂ (2 mL). ^bIsolated yield after chromatography. ^cHydrolysis of **1aa** was observed. ^d**1aa** was unchanged [Ms = methanesulfonyl, JohnPhos = (2-biphenyl)di-*tert*-butylphosphine].

Gold(III) chloride (entry 1) and Echavarren's gold(I) catalyst [JohnPhosAu(CH₃CN)]SbF₆ (entry 2) promoted the formation of the pyrrole derivative **2aa** in 78% and 89% yields respectively, whereas Ph₃PAuCl catalysed the formation of the dihydropyridine **3aa** in 58% yield (entry 3). Interestingly, dihydropyridine **3aa** was selectively formed when a silver co-catalyst AgSbF₆ was used with any of these gold catalysts (entries 4-6). Selective formation of the dihydropyridine **3aa** in a reasonably good yield (66%) was also observed when AgSbF₆ alone was employed as a catalyst (entry 7). Silver triflate, however, failed to promote the cyclisation reaction altogether (entry 8). The combination of 2 mol% Ph₃PAuCl and 5 mol% AgSbF₆ afforded the best yield of dihydropyridine **3aa**, while Echavarren's catalyst (2 mol%) furnished the highest yield of the pyrrole **2aa**. It may be noted that, in almost all cases, the cyclisation afforded either the dihydropyridine or the pyrrole exclusively. The position of substituents on the pyrrole ring of **2aa** indicated that **1aa** has undergone a gold-catalysed propargyl-Claisen rearrangement¹⁴ prior to the final cyclisation, whereas the skeletal reorganisation was not apparent in the structure of dihydropyridine **3aa**. The intriguing catalyst-controlled divergence in the cyclisation pathways called for further investigations. Thus, the *N*-propargyl-*N*-vinyl sulfonamides **1aa-hb** (table 1), prepared via the formal vinylic substitution method, were separately subjected to the optimised conditions for both the cycloisomerisation reactions. A variety of pyrrole derivatives **2aa-hb** and dihydropyridines **3aa-hb** were formed in these reactions and they are listed in Table 3. It was evident that the cycloisomerisation reactions and the catalyst-

controlled divergence were general. The sulfonyl groups on the nitrogen (from alkyne component) as well as carbon (allene component) can be varied. Cycloisomerisations of aza-enynes with arenes or a heteroarene (2-thienyl) as the alkyne substituent (R^1) afforded 3-(hetero)arylpyrroles (**2da-fb**) and 4-(hetero)aryldihydropyridines (**3da-fb**) in excellent yields. Single crystal X-ray analysis of the representative pyrrole **2da** confirmed the assigned structure.¹⁵

Interestingly, dihydropyridine formation was not observed when the aza-enyne **1hb** carrying a propargylic phenyl substituent (R^2 = Ph) was subjected to either of the conditions for cyclisation. In both instances, 2-benzyl-5-sulfonylmethyl pyrrole **2hb** was formed as the sole product. Presumably, presence of the propargylic phenyl substituent imparts a substrate bias for pyrrole formation that overrides the catalyst-control of cycloisomerisation pathways.

It is important to note that 3-arylpyrroles are much-preferred intermediates for the synthesis of a number of natural products such as rhazinilam, lamellarins and pyrrolnitrin.¹⁶ Similarly, 1,2-dihydropyridines are useful precursors for the preparation of polysubstituted 6-membered nitrogen heterocycles and isoquinuclidines.¹⁷ Therefore, the quick and efficient assembly reported here, of pyrroles and 1,2-dihydropyridines armed with readily malleable substituents at 2- and 4-positions, has the potential to find applications in targeted synthesis.


A tentative mechanistic rationalisation for the formation and cycloisomerisation of the representative aza-enyne **1da** is depicted in Scheme 2. The control experiments indicated that the allenyl sulfone **6** is a plausible intermediate in the formation of the aza-enyne **1**. The latter undergoes a metal-mediated propargyl-Claisen rearrangement¹⁴ to generate the β -allenyl imine **7A**. It may be noted that the aza-enyne **1aa** did not undergo any rearrangement in the absence of catalysts (table 2, entry 9). Gold-mediated 5-exo-dig cyclisation of **7A** and subsequent aromatisation via hydrogen shift affords the pyrrole derivative **2da**. The dihydropyridine **3da** is presumably formed *via* the tautomerisation¹⁸ of **7A** to the azatriene **7B** and subsequent 6π -aza-electrocyclisation.

The transformations listed in Scheme 3 provide a glimpse of the synthetic potential embedded in the heterocycles generated in this study. A regioselective bromination of the pyrrole **2aa** to afford 3-bromopyrrole **8** was achieved by treatment with *N*-bromosuccinimide in THF. Similarly, site-selective catalytic hydrogenation of the dihydropyridine **3ab** furnished the tetrahydropyridine **9**. The dihydropyridine **3ab** was also converted smoothly into the corresponding pyridine derivative **10**¹⁹ via base-promoted elimination of methanesulfonyl group. The phenylsulfonyl group has been termed 'arguably the most versatile functional group' by Fuchs²⁰ and it is conceivable that its potential may be exploited in further synthetic elaboration at the methylenesulfonyl unit of the products.

Table 3. Gold and silver-catalysed divergent cycloisomerisation of aza-enynes **1aa-hb**.^a


2aa , R^3 = Me, R^4 = Ph, 89% 2ab , R^3 = Me, R^4 = <i>p</i> -tolyl, 96% 2ba , R^3 = Ph, R^4 = Ph, 98% 2bb , R^3 = Ph, R^4 = <i>p</i> -tolyl, 97% 2ca , R^3 = <i>p</i> -tolyl, R^4 = Ph, 97% 2cb , <i>p</i> -tolyl, R^4 = <i>p</i> -tolyl, 99%	2da , R^4 = Ph, 96% 2db , R^4 = <i>p</i> -tolyl, 92% 2ea , R^4 = Ph, 95% 2eb , R^4 = <i>p</i> -tolyl, 97% 2fa , R^4 = Ph, 87% 2fb , R^4 = <i>p</i> -tolyl, 89% 2ga , R^4 = Ph, 91% 2gb , R^4 = <i>p</i> -tolyl, 98% 2ha , R^4 = Ph, 78% 2hb , R^4 = <i>p</i> -tolyl, 87% 2ib , R^4 = <i>p</i> -tolyl, 96% 2ja , R^4 = Ph, 98% 2jb , R^4 = <i>p</i> -tolyl, 97%

^aReaction conditions: **1aa-hb** (0.5mmol), catalysts, CH₂Cl₂ (2 mL), 25 °C, 1 h. Isolated yields. [Au] = 2 mol% (JohnPhos)Au(CH₃CN)SbF₆; [Au-Ag] = 2 mol% Ph₃PAuCl & 5 mol% AgSbF₆.



Scheme 2. Mechanistic rationalisation for the formation and cycloisomerisation of aza-enyne **1da**.

COMMUNICATION

Scheme 3. Synthetic applications of the pyrrole **2aa** and dihydropyridine **3ab**.

In summary, a catalyst-controlled divergent cycloisomerisation reaction of *N*-propargyl-*N*-vinyl sulfonamides to selectively afford either 2-sulfonylmethyl pyrroles or dihydropyridines was developed. The *N*-propargyl-*N*-vinyl sulfonamides were in turn prepared via a novel, base-mediated formal vinylic substitution reaction of 2-bromoallyl sulfones. A variety of 2,3,4-substituted pyrroles and 2,4-substituted dihydropyridines were assembled in two steps from propargyl sulfonamides and 2-bromoallyl sulfones. It is presumable that the rich chemistry of the methylenesulfonyl moiety may be exploited for further elaboration of the heterocycles into important natural products and analogues. Efforts along this direction and a detailed investigation of the cesium carbonate mediated vinylic substitution reaction are currently underway.

The Department of Science and Technology (DST), India is acknowledged for the award of Ramanujan fellowships to both RSM and KKS. DST is thanked for the award of a Start-up Research Grant (young scientists) for RSM. Financial support in part from the XIIth five year plan project "Affordable Cancer Therapeutics (ACT)" and Research Fellowship for SU from the Council of Scientific and Industrial Research (CSIR), India is also acknowledged.

Notes and references

- (a) Sharpless's deliberation, in his Nobel-prize lecture, on the regio- and stereoselective oxidation of the olefinic units of geraniol may be recalled here: K. B. Sharpless, *Angew. Chem. Int. Ed.*, 2002, **41**, 2024; Also see; (b) C. A. Lewis, S. J. Miller, *Angew. Chem. Int. Ed.*, 2006, **45**, 5616; (c) C. A. Lewis, K. E. Longcore, S. J. Miller, P. A. Wender, *J. Nat. Prod.*, 2009, **72**, 1864.
- J. Mahatthananchai, A. M. Dumas, J. W. Bode, *Angew. Chem. Int. Ed.* 2012, **51**, 10954 and references cited therein.
- For selected recent examples of catalyst-controlled divergent reactions, see: (a) A. García-Rubia, J. A. Romero-Revilla, P. Mauleón, R. G. Arrayás, J. C. Carretero, *J. Am. Chem. Soc.*, 2015, **137**, 6857; (b) R. Sakae, K. Hirano, M. Miura, *J. Am. Chem. Soc.*, 2014, **137**, 6460; (c) Y. Jiang, C. -M. Park, *Chem. Sci.*, 2014, **5**, 2347; (d) J. H. Kim, J. Bouffard, S.-g. Lee, *Angew. Chem. Int. Ed.*, 2014, **53**, 6435; (e) B. Li, Y. Park, S. Chang, *J. Am. Chem. Soc.*, 2014, **136**, 1125; (f) Y. Tani, T. Fujihara, J. Terao, Y. Tsuji, *J. Am. Chem. Soc.*, 2014, **136**, 17706; (g) G. -Q. Chen, X. -N. Zhang, Y. Wei, X. -Y. Tang, M. Shi, *Angew. Chem. Int. Ed.*, 2014, **53**, 8492; (h) W. Muramatsu, Y. Takemoto, *J. Org. Chem.*, 2013, **78**, 2336.
- For a review, see: (a) V. Michelet, P. Y. Toullec, J. -P. Genêt, *Angew. Chem. Int. Ed.*, 2008, **47**, 4268
- For selected reviews, see: (a) R. Dorel, A. M. Echavarren, *Chem. Rev.*, 2015, DOI:10.1021/cr500691k; (b) M. E. Muratore, A. Homs, C. Obradors, A. M. Echavarren, *Chem. Asian J.*, 2014, **9**, 3066; (c) M. Rudolph, A. S. K. Hashmi,

- Chem. Soc. Rev.*, 2012, **41**, 2448; (d) H. C. Shen, *Tetrahedron*, 2008, **64**, 3885; (e) Z. Li, C. Brouwer, C. He, *Chem. Rev.* 2008, **108**, 3239; (f) A. S. K. Hashmi, *Chem. Rev.*, 2007, **107**, 3180.
- For selected reviews, see: (a) L. Fensterbank, M. Malacria, *Acc. Chem. Res.*, 2014, **47**, 953; (b) C. Obradors, A. M. Echavarren, *Acc. Chem. Res.*, 2014, **47**, 902; (c) A. Fürstner, *Acc. Chem. Res.*, 2014, **47**, 925.
- (a) E. Jiménez-Núñez, A. M. Echavarren, *Chem. Rev.*, 2008, **108**, 3326; (b) D. J. Gorin, F. D. Toste, *Nature*, 2007, **446**, 395; (c) A. Gimeno, A. B. Cuenca, S. Suárez-Pantiga, C. R. de Arellano, M. Medio-Simón, G. Asenio, *Chem. -Eur. J.*, 2014, **20**, 683.
- (a) W. Rao, W. Susanti, B. J. Ayers, P. W. H. Chan, *J. Am. Chem. Soc.*, 2015, **137**, 6350; (b) C. Wang, X. Xie, J. Liu, Y. Liu, Y. Li, *J. Sun, Angew. Chem. Int. Ed.*, 2015, **54**, 883; (d) R. K. Shiroodi, O. Koleda, V. Gevorgyan, *J. Am. Chem. Soc.*, 2014, **136**, 13146; (e) Q. Zeng, L. Zhang, J. Yang, B. Xu, Y. Xiao, J. Zhang, *Chem. Commun.*, 2014, **50**, 4203; (f) W. Rao, M. J. Koh, P. W. H. Chan, *J. Org. Chem.*, 2013, **78**, 3183; (g) P. Mauleón, R. M. Zeldin, A. Z. González, F. D. Toste, *J. Am. Chem. Soc.*, 2009, **131**, 6348.
- (a) P. R. Joshi, S. Undeela, D. D. Reddy, K. K. Singarapu, R. S. Menon, *Org. Lett.*, 2015, **17**, 1449; (b) S. Reddy, S. Thadkapally, M. Mamidyalu, J. B. Nanubolu, R. S. Menon, *RSC Adv.*, 2015, **5**, 8199.
- (a) A. Jose, A. J. Jayakrishnan, B. Vedhanarayana, R. S. Menon, S. Varughese, E. Suresh, V. Nair, *Chem. Commun.*, 2014, **50**, 4616; (b) E. Li and Y. Huang, *Chem. Commun.*, 2014, **50**, 948; (c) S. Yu and S. Ma, *Angew. Chem. Int. Ed.*, 2012, **51**, 3074; (d) A. Jose, K. C. S. Lakshmi, E. Suresh, V. Nair, *Org. Lett.*, 2013, **15**, 1858.
- (a) A. Núñez Jr., M. R. Martín, A. Fraile, J. L. G. Ruano, *Chem. -Eur. J.*, 2010, **16**, 5443; (b) C. Lu, X. Lu, *Tetrahedron*, 2004, **60**, 6575.
- S. Ma, Q. Wei, *J. Org. Chem.*, 1999, **64**, 1026.
- For relevant examples, see: (a) A. Saito, T. Konishi, Y. Hanzawa, *Org. Lett.*, 2010, **12**, 372; (b) T. Harschneck, S. F. Kirsch, *J. Org. Chem.*, 2011, **76**, 2145.
- B. D. Sherry, F. D. Toste, *J. Am. Chem. Soc.*, 2004, **126**, 15978.
- CCDC 1063242 (**2da**) contains the supplementary crystallographic data for this compound. This data can be obtained free of charge from The Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif.
- (a) For a synthesis of rhazinilam family of alkaloids, see: L. McMurray, E. M. Beck, M. J. Gaunt, *Angew. Chem. Int. Ed.*, 2012, **51**, 9288; (b) For the synthesis of a lamellarin alkaloid, see: K. Hasse, A. C. Willis, M. G. Banwell, *Eur. J. Org. Chem.*, 2011, 88; (c) For pyrrolnitrin synthesis, see: M. D. Morrison, J. J. Janthorn, D. A. Pratt, *Org. Lett.*, 2009, **11**, 1051.
- (a) E. M. P. Silva, P. A. M. M. Varandas, A. M. S. Silva, *Synthesis*, 2013, **45**, 3053; (b) B. M. Trost, B. Biannic, *Org. Lett.*, 2015, **17**, 1433; (c) H. Mizoguchi, R. Watanabe, S. Minami, H. Oikawa, H. Oguri, *Org. Biomol. Chem.*, 2015, **13**, 5955; (d) K. Ishihara, H. Yamada, M. Akakura, *Chem. Commun.*, 2014, **50**, 6357 and referenced cited therein.
- H. Menz, S. F. Kirsch, *Org. Lett.*, 2006, **8**, 4795.
- H. Richter, S. Beckendorf, O. G. Mancheño, *Adv. Synth. Catal.*, 2011, **353**, 295.
- A. El-Awa, M. N. Noshi, X. M. du Jourdin, P. L. Fuchs, *Chem. Rev.*, 2009, **109**, 2315.