

Green Chemistry

Accepted Manuscript

This article can be cited before page numbers have been issued, to do this please use: A. R. K. Tiwari and B. M. Bhanage, *Green Chem.*, 2015, DOI: 10.1039/C5GC01884F.



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about *Accepted Manuscripts* in the **Information for Authors**.

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard <u>Terms & Conditions</u> and the <u>Ethical guidelines</u> still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.



www.rsc.org/greenchem



Journal Name

COMMUNICATION

Received 00th January 20xx, Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

www.rsc.org/

Polythene glycol (PEG) as a Reusable Solvent System for The Synthesis of 1,3,5-triazines *via* Aerobic Oxidative Tandem Cyclization of Benzylamines and *N*-substituted Benzylamines with Amidines under Transition Metal-Free Conditions

Abhishek R. Tiwari and Bhalchandra M. Bhanage*

A green and highly efficient protocol for the syntheis of 1,3,5triazines from benzylamines and N-substituted benzylamines amidines in PEG-600 has been developed firstly. This protocol is transition-metal free, phosphine ligand free and uses inexpensive, easily available molecular oxygen (O_2) as an oxidant. A series of 1,3,5-triazines derivatives were synthesized in good to excellent yields in shorter reaction time. The ease of the products separation and resuability of PEG-600 makes it more environmentally benign and economically affordable for the gramscale synthesis.

Green Chemistry emphasized on the development of economically feasible, synthetic procedures that avoid the use toxic transition-metals and the utilization of of environmentally benign substances and non-toxic solvents.¹ Recently, transition-metal free reactions have gained prominence in organic synthesis.² Not only does it avoids the use of toxic transition-metal, but also reduces the cost of the developed procedure. Oxidation reactions are very important transformations in organic chemistry. Molecular oxygen (O₂) is an ideal oxidant as it is inexpensive and easily available.³ In addition, no toxic by-product makes it a highly attractive reagent from the viewpoint of green sustainable chemistry.⁴ The most of organic reactions are affected by solvent as it plays an important role in mixing the ingredients to make the system homogeneous and allow molecular interactions to be more efficient.⁵ Thus, selection of solvent is very crucial. The solvent should be inexpensive, non-toxic, non-volatile and easy recyclability. In this context, environmentally benign solvents such as water⁶, ionic-liquids⁷ has been employed in various organic reactions. However, most of organic moieties have very low solubility in water which results into low yield of products. Further, toxicity and environmental burden data for the most of the ionic-liquids are still unknown. Over the past

available, recyclable, immiscibility with various organic solvents and non-toxic media for phase transfer catalysts.⁹ Further, PEG is an inexpensive, completely non-halogenated and complete toxicity profiles are available for a range of polyethylene glycol (PEG) molecular weights, many are approved for internal utilization by the US FDA.¹⁰ 1,3,5-triazines chemistry is of great importance due to their wide applications in biological and medicinal activities such as antimicrobial,^{11a} antimalarial,^{11b} antitumor agents^{11c} antituberculosis^{11d} and inhibition of photosynthetic electron transport (PET) and binding.^{11e} In addition, they are used as

decade, polythene glycol (PEG) has been used in several

organic reactions.⁸ PEG and its monomethyl ethers posses

unique properties such as thermally stable, commercially

chelating ligands for the preparation of organometallic materials,¹² transition-metal catalysts,¹³ liquid crystals,¹⁴ fluorescent brighteners.¹⁵ Although, 1,3,5-trianzines possess extensive functions, but only a few methods for the synthesis of 1,3,5-triazines have been reported. Traditionally, they were synthesized from halogenated 1,3,5-triazines in the presence



Scheme 1 Synthesis of 1,3,5-triazines

of transition metal-catalysts,¹⁶ and from the cyclotrimerization reaction of nitriles.¹⁷ However, the former require transitionmetal Palladium (Pd), less-environmentally benign halogenated substrates, and produce stoichiometric amounts

Department of Chemistry, Institute of Chemical Technology (ICT), Mumbai, India bm.bhanage@gmail.com, bm.bhanage@ictmumbai.edu.in,

Fax: +91 2233611020; Tel.: +91 2233612601

<code>†Electronic Supplementary Information (ESI)</code> available: [¹H NMR, ¹³C NMR, HRMS and GCMS]. See DOI: 10.1039/x0xx00000x

DOI: 10.1039/C5GC01884F Journal Name

COMMUNICATION

of undesirable waste, and the later usually needs excess of amines as the co-catalysts. Alternatively, these compounds were also obtained by cyclization of aromatic aldehydes with amidines.¹⁸ However, the use of aldehydes have several disadvantages such as aldehydes could undergo a decarbonylation reaction under harsh reaction conditions,¹⁹ oxidation of active aldehyde groups, leading to formation of unwanted by-products, hence require inert conditions.²⁰ Recently, two methods for synthesis of 1,3,5-triazines from of benzyl alcohol and amidines have been reported (Scheme 1). Nevertheless, these methods require costly rutheniumcomplex²¹ or less environmentally begin Cu(OAc)₂²² as a catalyst. Moreover, difficult preparation step of Ru-phoshine complex, toxic solvents such as DMSO and toluene, longer reaction time, reflux conditions and aqueous work up has imposed limitations on the applicability of these methods.

In continuation with our ongoing work on transition-metal free approach in organic synthesis.^{2h} Herein, we report a transition-metal free method for synthesis of 1,3,5-triazines. Notably, this reaction proceeds *via* aerobic oxidative tandem cyclization of benzylamines with amidines at 130 °C for 3h.

Table 1 Optimization of reaction condition



^a Reaction conditions: benzylamine (**1a**, 0.5 mmol), benzamidine hydrochloride (**2a**, 1.0 mmol), Cs₂CO₃ (1.0 mmol) and solvent (2.5 mL). ^b Isolated Yield. ^c Na₂CO₃ was used instead of Cs₂CO₃. ^d K₂CO₃ was used instead of Cs₂CO₃. ^e Cs₂CO₃ (0.5 mmol).

Initially, the direct reaction between benzylamine (1a) and benzamidine hydrochloride (2a) was selected as a model reaction to evaluate the feasibility of our system in environment benign solvent at 100 °C for 24 h. When the reaction was carried out in H_2O , ethanol and glycerol, no

formation of the desired product was noted (table 1, entries 1-3). Surprisingly, changing the solvent system to PEG-600 resulted into formation of the desired product 3a in 70% yield (table 2, entry 4). This surprise result encouraged us to choose this PEG-600 as solvent. To our delight, increasing the temperature has a significant effect on the yield of 3a, provided 95% yield at 130 °C (table 1, entry 5-7). Increasing the reaction temperature above 130 °C didn't have significant effect on the yield of 3a (table 1, entry 8). Reaction time could be reduced to 3 h from 24 h (table 1, entries 9-11). Decreasing the reaction time beyond 3 h leads to significant decrease in the yield of desired product 3a (table 1, entry 12). Before heating the reaction mixture was transparent but after heating for 3h, it turns into yellow in colour. The addition of DMSO resulted into decrease in the yield of product 3a (table 2, entry 13). When reaction was carried out in the absence of base, no formation of desired product was observed, even after running the reaction for 24 h (table 1, entry 14). Bases such as Na₂CO₃ and K₂CO₃ did not give good results for the formation of desired product 3a (table 1, entry 16). Further, decreasing the amount of base from 1.0 mmol to 0.5 mmol resulted into the formation product 3a only in 51% (table 1, entry 16). Thus, optimized reaction conditions are: benzylamine (1a, 0.5 mmol) amidine hydrochloride (2a, 1.0 mmol), Cs₂CO₃ (1.0 mmol), PEG-600 (2.5 mL) for 3 h under O₂.

Encouraged by these results, we used various benzylamines and amidines to establish the scope and limitations of this protocol. A series of 1,3,5-triazines were synthesized in good to excellent yields under the optimized reaction conditions and representative results are listed in table 2. Firstly, the effect of electron donating group and electron withdrawing on benzylamine with benzamidine was studied (table 2, entries 1-8). It was found that electron donating substituents such as -Me, -OMe, proceeds smoothly and furnished desired products (3b-3c) in excellent isolated yields (table 2, entries 2 and 3). Subsequently, the reaction of benzylamines bearing strong electron withdrawing groups such as -CN and -NO2 at different position provided corresponding products (3d, 3e and 3f) in good yields (table 2, entries 5-6). Interestingly, halogen substituents such as -Cl and -F could also be transformed in efficient manner, providing respective products in very good yields (table 2, entries 7-8). Next, heteroatom containing benzylamines such as pyridine-2yl-2-methanamine, pyridin-3-ylmethanamine (1i and 1j) and furan-2-ylmethanamine (1k) were tested. Delightfully, all the reactions progressed efficiently affording products 3i-3k in very good yields (table 2, entries 9-11). These obtained heteroaryl substituted 1,3,5-triazines have potential to be used as C^N or C^N^C ligands in pincer complexes.²³ Next, an apparent substituents effect on amidine was also explored (table 2, entries 12-15). eaction of 1a, 1c and 1m with paramethylbenzamidine (2b) progressed very well affording 1,3,5triazines triazines (3I-3n) in excellent yields (table 2, entries 12-14). Also, reaction of 1c with para-bromobenzamidine resulted into the formation of desired product (30) in good yield (table 2, entry 15). Unfortunately, no formation of corresponding products could be observed when the reaction was carried out

Journal Name

$\ensuremath{\textbf{Table 2}}$ Aerobic oxidative cyclization of benzylamines with amidine hydrochlorides $\ensuremath{^{\circ}}$









 a Reaction conditions: benzylamine (1, 0.5 mmol), amidine hydrochlorides (2, 1.0 mmol), Cs₂CO₃ (1.0 mmol), solvent (2.5 mL) for 3 h. b Isolated Yield.

with both aliphatic amines and aliphatic amidines (results are not shown).

Efforts were also made to expand the scope of the method to *N*-mono and *N*,*N*-di substituted benzylamines and results are summarized in table 3. The reaction proceeded very well for *N*-methylbenzylamines (**1aa**) and *N*,*N*-dimethylbenzylamine

COMMUNICATION

Published on 01 September 2015, Downloaded by University of California - San Diego on 01/09/2015 17:33:24

Table 3 Aerobic oxidative cyclization of *N*-substituted amines with amidine hydrochlorides^a



Entry	Amine (1)	Amidine (2)	Product (3)	Yield ^b (%)
1	1aa	2a	3a	94
2	1ca	2a	3c	95
3	d 1ma	2a		89
4	1aa	2b	3m	83
5	1ca	2b	3n	84
6	1ma	2b	30	86
7	1ab	2a	3a	90
8	1aaa	2a	3a	78,82 c
9	1aaa	2b	3m	81
10	1ab'	2a	3a	86
11	1abb	2a	3a	70 ^c

^a Reaction conditions: benzylamine (**1**, 0.5 mmol), amidine hydrochlorides (**2**, 1.0 mmol), Cs_2CO_3 (1.0 mmol), solvent (2.5 mL) for 3 h. ^b Isolated Yield. ^C Reaction was carried out for 8 h.

(1aaa) providing corresponding products in very good yield (table 3, entries 1-6 and 8). Also, the reactions of N-ethylbenzylamine (1ab) and N-ethanolbenzylamine (1ab') were found to be effective (table 3, entry 7 and 10). However, reaction of *N*,*N*-diethylbenzylamine (1abb) gave relatively low yield of 3a than 1aaa (table 3, entries 8 and 11).

To show the synthetic utility of this protocol, gram-scale reaction were carried out by using substrates **1a** (2 g, 18.87 mmol) with **2a** (5.89 g, 37.74 mmol) and **1c** (2.0 g, 16.53 mmol) with **2a** (5.16 g, 33.06 mmol) under the optimized reaction condition. As per our expectation, reaction preceded well by providing **3c** and **3g** in 91% and 85% isolated yields respectively.

In order to understand the mechanism of these reactions, some control experiments were carried out (Scheme 2). When reaction was carried out in the absence of amidines and Cs₂CO₃ formation of imine (5a) was noted by self coupling of benzylamine. On the other hand, N-methyl and N,Ndimethylbenzylamine remained unchanged. This is an agreement with the previous report on the synthesis of imines.²⁴ However, in the presence of para-methoxyaniline (4a) formation of imine (5aa) was observed form 1a, 1aa and 1aaa. The yields of these imines are very low even after running the reaction for 16 h. This is possibly because 4a is very less basic than amidines. The formation of these imines (5a and 5aa) was confirmed by GCMS. This proves that the presence of another amine influence the reaction and leads to the formation of aldehyde via oxidative cleavage of benzylic carbon and nitrogen bond. Based on the observations of control experiments, a plausible reaction mechanism has been illustrated (Scheme 3). Reaction proceeds with the in situ



Scheme 2 Control experiments



generation of aldehyde (**A**). Meanwhile, amidine salt (**2**) is neutralized by Cs_2CO_3 from its hydrochloride salt. Consequentially, reacting with **A** to give 1,3,5-tirazine (**3a**) *via* dehydrogenative aromatisation of **C**. This proposed mechanism is consistent with the previous report.^{21,22} To the best of our knowledge, there is no report on the synthesis of imines form N-mono substituted benzylamine under transition-metal free condition. This is for the first time we have shown the formation of imine from N-mono and disubstituted benzylamine under transition-metal free conditions. Further, unlike the previous report,^{24,25} self coupling reaction of benzylamine does not involve the use of any activating agents such as catalyst or acid.

DOI: 10.1039/C5GC01884F

Journal Name

Journal Name

At last, we attempted to reuse the PEG-600. After completion of the reaction, PEG was recovered and subjected to another run, affording the product in almost same yield. This process was repeated three more times, affording the product in excellent yields (Table 4). It is important to note that weight loss ~10% of PEG was observed for every run due to handling loss. The simple experimental and ease of product separation combined with the easy recovery and reuse of PEG is expected to contribute to the development of a green methodology for the synthesis of 1,3,5-triazines.

Table 4 Recyclability study of PEG

Run	1	2	3
Yield ^b (%)	95	95	94

 a Reaction conditions: benzylamine (1, 0.5 mmol), amidine hydrochlorides (2, 1.0 mmol), Cs₂CO₃ (1.0 mmol), PEG (2.5 mL) for 3 h. b Isolated Yield.

Conclusions

In conclusion, a highly efficient, transition-metal free synthesis of 1,3,5-triazines has been developed by employing molecular oxygen as a green oxidant in PEG-600. The use of molecular oxygen and non toxic as solvent PEG-600 has made this protocol economical, environmentally benign and potentially viable for commercial and academic applications. Various 1,3,5-triazines were synthesized in good to excellent yields. The developed methodology is suitable for the gram-scale synthesis, owing to its simple non-aqueous work up, shorter reaction time and higher yield. This protocol offers an excellent chance to avoid the toxic solvents, catalysts and use of natural resources than previous reports. Noteworthy, for the first time we have demonstrated the synthesis of imines form N-mono and N,N-di substituted benzylamines. Moreover, the after extraction of with ether, PEG-600 was recovered and could be reused up to three consecutive cycles.

Acknowledgment

The author (ART) is greatly thankful to UGC (University Grant Commission) India, New Delhi for providing BSR fellowship.

Notes and references

- 1 R. A. Sheldon, I. Arends, and U. Hanefeld, Green Chemistry and Catalysis, WILEY-VCH Verlag GmbH & Co. KGaA, Weinheim, 2007.
- 2 (a) Y. Yuan, I. Thomé, S. H. Kim, D. Chen, A. Beyer, J. Bonnamour, E. Zuidema, S. Chang and C. Bolm, Adv. Synth. Catal., 2010, 352, 2892; (b) R. Cano, D. J. Ramón and M. Yus, J. Org. Chem., 2011, 76, 654; (c) Y. Fang, Y. Zheng and Z. Wang, Eur. J. Org. Chem., 2012, 1495; (d) L. H. Zou, J. Reball, J. Mottweiler, and C. Bolm, Chem. Commun., 2012, 48, 11307; (e) F. Diness and D. P. Fairlie, Angew. Chem., Int. Ed., 2012, 51, 8012; (f) M. C. Pérez-Aguilar, and C. Valdés, Angew. Chem., Int. Ed., 2012, 51, 5953; (g) N. Jalalian, T. B. Petersen, and B. Olofsson, Chem. Eur. J., 2012, 18, 14140; (h) A. R. Tiwari and B. M. Bhanage, RSC Adv., 2015, 5, 57235.

Green Chemistry Accepted Manuscrip

- 3 A. N. Campbell and S. S. Stahl, Acc. Chem. Res., 2012, 45, 851.
- 4 (a) B. M. Trost, *Science*, 1991, 254, 1471; (b) R. A. Sheldon, I. W. C. E. Arends, G. J. Ten Brink and A. Dijksman, *Acc. Chem. Res.*, 2002, 35, 774; (c) A. S. K. Hashmi and G. J. Hutchings, *Angew. Chem., Int. Ed.*, 2006, 45, 7896; (d) A. Corma and H. Garcia, *Chem. Soc. Rev.*, 2008, 37, 2096; (e) Y. Chen, D. M. Ho and C. Lee, *J. Am. Chem. Soc.*, 2005, 127, 12184; (f) L. Ackermann and L. T. Kaspar, *J. Org. Chem.*, 2007, 72, 6149; (g) For a review, see : Z. Shi, C. Zhang, C. Tanga and N. Jiao, *Chem. Soc. Rev.*, 2012, 41, 3381.
- 5 C. Reichardt and T. Welton, Solvents and Solvent Effects in Organic Chemistry, WILEY-VCH Verlag GmbH & Co. KGaA, Weinheim, 2011.
- 6 (a) Organic Synthesis in Water, ed. P. A. Grieco, Blackie Academic and Professional, London, 1998; (b) Organic Reactions in Aqueous Media, ed. C. J. Li, T. H. Chan, John Wiley and Sons, New York, 1997; (c) R. Breslow, Acc. Chem. Res., 1991, 24, 159; (d) E. Rangel Rangel, E. M. Maya, F. Sánchez, J. G. de la Campaa and M. Iglesias, Green Chem., 2015, 17, 466; (e) P. Puthiaraja and K. Pitchumani, Green Chem., 2014, 16, 4223; (f) For a review, see : M. B. Gawande, V. D. B. Bonifa'cio, R. Luque, P. S. Brancoa and R. S. Varma, Chem. Soc. Rev., 2013, 42, 5522.
- 7 (a) R. A. Sheldon, *Chem. Commun.*, 2001, 2399; (b) C. L. Hussey, *Pure Appl. Chem.*, 1988, **60**, 1763; (c) M. J. Earle and K. R. Seddon, *Pure Appl. Chem.*, 2000, **72**, 1391; (d) T. Welton, *Chem. Rev.*, 1999, **99**, 2071; (e) P. Wasserscheid and W. Keim, *Angew. Chem.*, *Int. Ed.*, 2000, **39**, 3772; (e) K.V. Wagh, B.M. Bhanage, *Green Chem.*, 2015, **17**, 4446; (f) K.V. Wagh, B.M. Bhanage, *RSC Adv.*, 2014, **4**, 22763;
- 8 (a) For a review, see: J. Chen, S. K. Spear, J. G. Huddleston and R. D. Rogers, *Green. Chem.*, 2005, **7**, 64; (b) R. Kumar, P. Chaudhary, S. Nimesh and R. Chandra, *Green Chem.*, 2006, **8**, 356; (c) G. P. Lu, L. Y. Zeng and C. Cai, *Green Chem.*, 2011, **13**, 998; (d) K. S. Feu, A. F. de la Torre, S. Silva, M. A. F de M. Junior, A. G. Corrêa and M. W. Paixão, *Green Chem.*, 2014, **16**, 3169.
- 9 (a) Poly(ethylene Glycol) Chemistry, Biotechnological and Biomedical Applications, ed. J. M. Harris, Plenum Press, New York, 1992; (b) Polyethylene Glycol: Chemistry and Biological Application, ed. J. M. Harris, S. Zalipsky, American Chemical Society, Washington, DC, 1997.
- (a) N. Akiya and P. E. Savage, *Chem. Rev.*, 2002, **102**, 2725;
 (b) U. M. Lindström, *Chem. Rev.*, 2002, **102**, 2751.
- (a) M. Saleh, S. Abbott, V. Perron, C. Lauzon, C. Penney, B. Zacharie, *Bioorg. Med. Chem. Lett.*, 2010, **20**, 945; (b) S. Melato, D. Prosperi, P. Coghi, B. Basilico, D. Monti, *Chem. Med. Chem.*, 2008, **3**, 873 and references cited therein; (c) W. Zhu, Y. Liu, Y. Zhao, H. Wang, L. Tan, W. Fan and P. Gong, *Arch. Pharm. Chem. Life Sci.*, 2012, **345**, 812; (d) R. V. Patel, P. Kumari, D. P. Rajani, K. H. Chikhalia, *Eur. J. of Med. Chem.*, 2011, **46**, 4354; (e) A. Ohki, N. Kuboyama, K. Koizumi, A. Tanaka, Y. Sato, H. Kohno, Peter Böger, and K. Wakabayashi, *J. Agric. Food Chem.*, 1999, **47**, 4398.
- 12 (a) S. Naik, M. Kumaravel, J. T. Mague, M. S. Balakrishna, *Inorg. Chem.*, 2014, **53**, 1370; (b) C. Xiao, Y. Li, H. Lun, C. Cui, Y. Xu, J. Solid State Chem., 2013, **208**, 127.
- (a) M. H. Juàrez, M. Vaquero, E. Álvarez, V. Salazar, A. Suàrez, Dalton Trans., 2013, 42, 351; (b) P. K. Santra, P. Sagar, J. Mol. Catal. A: Chem., 2003, 197, 37.
- 14 (a) S. Kotha, D. Kashinath, S. Kumar, *Tetrahedron Lett.* 2008, 49, 5419; (b) C. H. Lee, T. Yamamoto, *Bull. Chem. Soc. Jpn.* 2002, 75, 615.
- 15 A. García, B. Insuasty, M. Herranz, R. M. Álvarez and N. Martín, *Org. Lett.*, 2009, **13**, 5398.
- 16 (a) D. Janietz and M. Bauer, Synthesis, 1993, 1, 33; (b) A. L. Isfahani, I. M. Baltork, V. Mirkhani, A. R. Khosropour, M.

Page 5 of 6

Published on 01 September 2015. Downloaded by University of California - San Diego on 01/09/2015 17:33:24.

Moghadam, S. Tangestaninejad and R. Kia, *Adv. Synth. Catal.*, 2013, **355**, 957.

- 17 (a) A. Diaz-Ortiz, A. de la Hoz, A. Moreno, A. S. Migallon and G. Valiente, *Green Chem.*, 2002, 4, 339; (b) F. Xu, J. H. Sun, H. B. Yan and Q. Shen, *Synth. Commun.*, 2000, 30, 1017; (c) R. D. Spencer and B. H. Beggs, *Anal. Chem.*, 1963, 35, 1633.
- 18 S. Biswas and S. Batra, Eur. J. Org. Chem., 2012, 18, 3492.
- (a) A. Modak, A. Deb, T. Patra, S. Rana, S. Maity and D. Maiti, Chem. Commun., 2012, **48**, 4253. (b) T. Iwai, T. Fujihara and Y. Tsuji, *Chem. Commun.*, 2008, 6215.
- 20 G. V. R. Sharma and A. Robert, *Res. Chem. Intermed.*, 2013, **39**, 3251.
- 21 F. Xie, M. Chen, X. Wang, H. Jiangb and M. Zhang, Org. Biomol. Chem., 2014, 12, 2761.
- 22 Q. You, Fei Wang, C. Wu, T. Shi, D. Min, H. Chen and W. Zhang, Org. Biomol. Chem., 2015, 13, 6723.
- (a) W. Yang, H. Fu, Q. Song, M. Zhang and Y. Ding, *Organometallics*, 2010, **30**, 77; (b) W. Wei, Y. Qin, M. Luo, P. Xia and M. S. Wong, *Organometallics*, 2008, **27**, 2268; (c) D. A. Smith, D. A. Roşca and M. Bochmann, *Organometallics*, 2012, **31**, 5998.
- 24 T. B. Nguyen, L. Ermolenko and A. Al Mourabit, *Green Chem.*, 2013, **15**, 2713.
- 25 (a) Z. Hu and F. M. Kerton, Org. Biomol. Chem., 2012, 10, 1618; (b) R. D. Patil and S. Adimurthy, Adv. Synth. Catal., 2011, 353, 1695; (c) L. Aschwanden, T. Mallat, F. Krumeich and A. Baiker, J. Mol. Catal. A: Chem., 2009, 309, 57; (d) B. Zhu, M. Lazar, B. G. Trewyna and R. J. Angelici, J. Catal. 2008, 260; (e) H. Guo, M. Kemell, A. Al Hunaiti, S. Rautiainen, M. Leskelä and T. Repo, Catal. Commun., 2011, 12, 1260; (f) S. Naya, K. Kimura and H. Tada, ACS Catal., 2013, 3, 10; (g) T. Hirao, M. Higuchi, I. Ikeda and Y. Ohshiro, J. Chem. Soc. Chem. Commun., 1993, 194; (h) K. Shimizu, K. Shimura, K. Ohshima, M. Tamura and A. Satsuma, Green Chem., 2011, 13, 3096; (i) S. Furukawa, Y. Ohno, T. Shishido, K. Teramura and T. Tanaka, ACS Catal., 2011, 1, 1150; (j) A. E. Wendlandt and S. S. Stahl, Org. Lett., 2012, 14, 2850; (k) C. Su, M. Acik, K. Takai, J. Lu, S. Hao, Y. Zheng, P. Wu, Q. Bao, T. Enoki, Y. J. Chabal and K. P. Loh, Nature Commun., 2012, 3, 1298; (I) H. Huang, J. Huang, Y. Liu, H. He, Y. Cao and K. Fan, Green Chem., 2012, 14, 930; (m) X. Lang, H. Ji, C. Chen, W. Ma and J. Zhao, Angew. Chem. Int. Ed., 2011, 50, 3934; (n) E. Zhang, H. Tian, S. Xu, X. Yu and Q. Xu, Org. Lett., 2013, 15, 2704; (o) F. Su, M. Antonietti, X. Wang, S. C. Mathew, L. Moehlmann and S. Blechert, Angew. Chem. Int. Ed. 2011, 50, 657; (p) H. Yuan, W. Yoo, H. Miyamura and S. Kobayashi, J. Am. Chem. Soc., 2012, 134, 13970; (q) M. Largeron and M. B. Fleury, Angew. Chem. Int. Ed., 2012, 51, 5409; (r) M. Largeron and M. B. Fleury, Science, 2013, 339, 43.