

## Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/gpss20>

### A Facile One-Pot Synthesis of Some New Spiro-thiazolidin-4-ones and Benzimidazoles of Biological Interest

Nadia Hanafy Metwally<sup>a</sup>

<sup>a</sup> Department of Chemistry, Faculty of Science ,  
University of Cairo , Giza, Egypt

Published online: 20 Feb 2008.

To cite this article: Nadia Hanafy Metwally (2007) A Facile One-Pot Synthesis of Some New Spiro-thiazolidin-4-ones and Benzimidazoles of Biological Interest, Phosphorus, Sulfur, and Silicon and the Related Elements, 183:1, 34-43, DOI: [10.1080/10426500701555702](https://doi.org/10.1080/10426500701555702)

To link to this article: <http://dx.doi.org/10.1080/10426500701555702>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or

indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

## A Facile One-Pot Synthesis of Some New Spiro-thiazolidin-4-ones and Benzimidazoles of Biological Interest

Nadia Hanafy Metwally

Department of Chemistry, Faculty of Science, University of Cairo,  
Giza, Egypt

*In a one-pot procedure, aromatic amines 1a–c, 9-fluorenone (2) and 2-mercaptoacetic acid (3) were converted into fluorenespiro-thiazolidinone derivatives 4a–c, which undergo condensation, oxidation and thiation to afford 6a–i, 7a–c, and 4-thioxo 8a–c, respectively. A new fluorenespiro-thiazolo-benzimidazole 10 was also obtained in one step via cyclocondensation of 1,2-phenylenediamine, 9-fluorenone and 2-mercaptoacetic acid. The obtained products seem to be interesting from the biological point of view.*

**Keywords** 2-Mercaptoacetic acid; 9-fluorenone; fluorenespiro[9.2']-thiazolo[3',4'-a]benzimidazole; fluorenespiro[9.2']-3'-aryl-thiazolidin-4-one derivatives

## INTRODUCTION


2-Substituted-thiazolidin-4-one derivatives exhibit unusual high activity in vitro against Mycobacterium Tuberculosis (TB), as drugs for anti-HIV and anticancer activity.<sup>1–4</sup> Recently, a number of 4-thiazolidinones having an 2-aryl substituted have been synthesized and found to exhibit potent selective anti-plateret activating factor both in vitro and in vivo.<sup>5</sup> Thiazolo[3,4-a]benzimidazoles and their analogues were reported to be potent anti-HIV agents.<sup>6,7</sup> Spiroheterocyclic compounds including thiazolidine moiety have also been shown to exhibit a variety of interesting biological activities.<sup>8–11</sup> In continuation to our search of new compounds with anticipated biological activity from accessible starting materials,<sup>12–17</sup> we report here a simple one-pot synthesis of some new fluorenespiro-thiazolidin-4-one and fluorenespiro-thiazolo-benzimidazole derivatives.

Received 5 May 2007; accepted 5 June 2007.

Address correspondence to Nadia Hanafy Metwally, Department of Chemistry, Faculty of Science, University of Cairo, Giza, Egypt. E-mail: nhmmohamed@yahoo.com


## RESULTS AND DISCUSSION

Aromatic amines **1a–c**, 9-fluorenone (**2**) and 2-mercaptocetic acid (**3**) react as a one pot reaction to afford white solids. The structures assigned for these solid compounds were proved based on elemental and spectral analysis and structures **4a–c** were assigned for them. For example, the IR spectrum of the isolated product **4a** showed absorption band near  $1684\text{ cm}^{-1}$  corresponding to (CO) group. The  $^1\text{H}$  NMR spectrum of **4a** revealed a singlet signal at  $\delta = 4.36$  ppm attributed to methylene protons and a multiplet at  $\delta = 6.60\text{--}7.93$  ppm corresponding to aromatic protons. Furthermore, the  $^{13}\text{C}$  NMR spectrum of **4a** revealed signals at  $\delta = 32.45$ ,  $75.13$ , and  $171.39$  ppm attributable to the methylene carbon, C-2' and thiazolidine carbonyl carbon, besides the signals corresponding to aromatic carbons (cf. Experimental Section and Scheme 1). Elemental analysis and mass spectral data agree with the proposed structures **4a–c**.



SCHEME 1

The methylene group in position-5 of thiazolidin-4ones was reported to be active.<sup>18,19</sup> Thus, compounds **4a–c** undergo condensation with the aldehydes **5a–c** in refluxing ethanol under basic catalysis to yield the 5-arylidene derivatives **6a–i**. The IR spectra of the isolated products showed absorption bands near  $1684\text{ cm}^{-1}$  due to CO group. The  $^1\text{H}$  NMR spectra of these products revealed the disappearance of methylene signals at  $\delta = 4.36\text{ ppm}$ , and showed a multiplet at  $\delta \approx 6.65\text{ ppm}$  attributable to olefinic and aromatic protons (cf. Experimental Section and Scheme 2).




**SCHEME 2**

Oxidation of compounds **4a–c** with a solution of  $\text{KMnO}_4$  under acidic conditions at room temperature furnished the products **7a–c**. The IR of these products showed absorption bands near  $1695\text{ cm}^{-1}$  corresponding to CO group;  $1341$  and  $1070\text{ cm}^{-1}$  corresponding to  $\text{SO}_2$  (asymmetrical and symmetrical str.) group. The  $^1\text{H}$  NMR spectra of these isolated products revealed a singlet at  $\sim\delta = 5.0\text{ ppm}$  attributed to the active methylene protons. Elemental analysis and mass spectra are in agreement with the sulfone structures **7a–c** (cf. Experimental Section and Scheme 2).

It has been reported that the biological activity thiazolidinone derivatives was much improved upon transformed to the thioxo-analogues.<sup>10,20</sup> Therefore, we have converted compounds **4a–c** into the corresponding thioxo-derivatives **8a–c** upon reaction with Lawesson's reagent under reflux in dry toluene for two hours (cf. Experimental Section and Scheme 2).

It is also interesting to extend our work to synthesize a novel spiro-thiazolo-benzimidazole via one-pot reaction of a mixture of 1,2-phenylenediamine, 9-fluorenone and 2-mercaptoacetic acid in refluxing toluene. From this reaction we could obtain a product of m.p. 168°C. Mass spectrum of this product showed  $m/e$  326. The IR spectrum showed the disappearance of band at  $\sim 1684\text{ cm}^{-1}$  corresponding to the carbonyl group. The  $^1\text{H}$  NMR spectrum revealed a singlet (2H) at  $\delta = 4.99\text{ ppm}$  attributed to methylene protons, beside the expected signals of aromatic protons. Furthermore, the  $^{13}\text{C}$  NMR showed a signal at  $\delta = 45.28\text{ ppm}$  corresponding to methylene carbon and no signal that can be attributed to a carbonyl carbon was shown at  $\sim \delta = 170\text{ ppm}$  and appearance of signal at  $\delta = 143.90\text{ ppm}$  corresponding to imine carbon ( $\text{C}=\text{N}$ ), besides the signals corresponding to aromatic carbons (cf. Experimental Section and Scheme 3). Elemental analysis



SCHEME 3

and mass spectra agree with the IR,  $^1\text{H}$  NMR and  $^{13}\text{C}$  NMR spectra data. Based on these collective data the fluorenespiro[9.2']thiazolo[3',4'- $\alpha$ ]benzimidazole structure **10** was assigned to this product. It is assumed that the reaction proceeds through the formation of non-isolable Schiff's base, followed by addition of the thiol to the C=N bond to afford the non-isolable **9**, which in role, undergoes ring closure in situ with the elimination of water to afford **10** (cf. Experimental Section and Scheme 3).

## EXPERIMENTAL

Melting points were determined on an electrothermal (9100) apparatus and are uncorrected. The IR spectra were recorded as KBr pellets on a Perkin Elmer 1430 spectrophotometer.  $^1\text{H}$  NMR and  $^{13}\text{C}$  NMR spectra were recorded in deuterated dimethylsulfoxide at 300 MHz on a Varian Gemini NMR spectrometer using tetramethylsilane as internal reference, and results are expressed as  $\delta$  values. Mass spectra were taken on a Shimadzu GCMS-QP 1000 Ex mass spectrometer at 70 eV. Elemental analyses were carried out at the Microanalysis Center of Cairo University, Giza, Egypt.

### Fluorenespiro[9.2']-3'-aryl-thiazolidin-4-ones **3a-c**—General Procedure

A mixture of the appropriate amines **1a-c** (20 mmol), 9-fluorenone (1.80 g, 10 mmol) and 2-mercaptoacetic acid (2.76 g, 30 mmol) in 25 ml dry toluene was refluxed for 8 h, and the solvent was evaporated till dryness under reduced pressure and the residue was triturated with ether. The ether layer was washed with sodium bicarbonate solution (1–2%) and finally with  $\text{H}_2\text{O}$ . The organic layer was dried over sodium sulfate and evaporated to dryness at reduced pressure. The solid so formed was collected by filtration and recrystallized from ethanol to give the title compounds, respectively (cf. Tables I and II).

### 5'-Arylidene-fluorenespiro[9.2']-3'-aryl-thiazolidin-4-ones **6a-i**—General Procedure

A mixture of compounds **4a-c** (10 mmol) and aldehydes **5a-c** (10 mmol) in 25 ml sodium ethoxide solution was refluxed for 1 h, left at room temperature, and the solid so formed was filtered off and recrystallized from ethyl alcohol/dioxin (cf. Tables I and II).

TABLE I Elemental Analyses of the Newly Synthesized Compounds

Compound no.	Solvent of Cry. yield %	Color M.P. °C	Mol. Formula Mol. wt	Analysis calcd./found %				
				C	H	N	S	Cl
<b>4a</b>	Ethanol 70	White 228	C <sub>21</sub> H <sub>15</sub> NOS 329	76.57 76.74	4.59 4.74	4.25 4.43	9.73 9.57	—
<b>4b</b>	Ethanol 74	White 198	C <sub>22</sub> H <sub>17</sub> NO <sub>2</sub> S 359	73.51 73.68	4.77 4.58	3.90 3.72	8.92 8.78	—
<b>4c</b>	Ethanol 72	White 208	C <sub>21</sub> H <sub>14</sub> ClNOS 363	69.32 69.51	3.88 4.06	3.85 4.04	8.81 8.98	9.74 9.92
<b>6a</b>	Ethanol/dioxane 86	Whitish brown 245	C <sub>28</sub> H <sub>19</sub> NOS 417	80.55 80.37	4.59 4.78	3.35 3.55	7.68 7.50	—
<b>6b</b>	Ethanol/dioxane 78	Whitish yellow 195	C <sub>29</sub> H <sub>21</sub> NO <sub>2</sub> S 447	77.83 77.70	4.73 4.58	3.13 3.30	7.16 7.30	—
<b>6c</b>	Ethanol/dioxane 82	Yellow 250	C <sub>28</sub> H <sub>18</sub> ClNOS 451	74.41 74.32	4.01 4.18	3.10 3.31	7.09 7.27	7.84 7.65
<b>6d</b>	Ethanol/dioxane 72	Yellow 228	C <sub>29</sub> H <sub>21</sub> NO <sub>2</sub> S 447	77.83 78.0	4.73 4.58	3.13 3.32	7.16 7.0	—
<b>6e</b>	Ethanol/dioxane 75	Yellow 236	C <sub>30</sub> H <sub>23</sub> NO <sub>3</sub> S 477	75.45 75.58	4.85 4.68	2.93 2.74	6.71 6.55	—
<b>6f</b>	Ethanol/dioxane 79	Yellow 250	C <sub>29</sub> H <sub>20</sub> ClNO <sub>2</sub> S 481	72.26 72.45	4.18 4.37	2.91 3.08	6.65 6.49	7.36 7.53
<b>6g</b>	Ethanol/dioxane 77	Yellow 202	C <sub>28</sub> H <sub>18</sub> ClNOS 451	74.41 74.58	4.01 4.19	3.10 3.27	7.09 7.25	7.84 7.67



TABLE I Elemental Analyses of the Newly Synthesized Compounds

Compound no.	Solvent of <i>Cry.</i> yield %	Color M.P. °C	Mol. Formula Mol. wt	Analysis calcd./found %				
				C	H	N	S	Cl
<b>6h</b>	Ethanol/dioxane 81	Yellow 248	C <sub>29</sub> H <sub>20</sub> ClNO <sub>2</sub> S 481	72.26 72.08	4.18 4.34	2.91 3.10	6.65 6.48	7.36 7.53
<b>6i</b>	Ethanol/Dioxane 75	Yellow 297	C <sub>28</sub> H <sub>17</sub> Cl <sub>2</sub> NOS 485	69.14 69.0	3.52 3.35	2.88 3.05	6.59 6.75	14.58 14.75
<b>7a</b>	Ethanol 74	White 248	C <sub>21</sub> H <sub>15</sub> NO <sub>3</sub> S 361	69.79 69.63	4.18 4.35	3.88 3.70	8.87 8.71	—
<b>7b</b>	Ethanol 72	White 238	C <sub>22</sub> H <sub>17</sub> NO <sub>4</sub> S 391	67.50 67.64	4.38 4.55	3.58 3.76	8.19 8.35	—
<b>7c</b>	Ethanol 78	White 252	C <sub>21</sub> H <sub>14</sub> ClNO <sub>3</sub> S 395	63.72 63.88	3.56 3.39	3.54 3.72	8.0 7.82	8.98 9.15
<b>8a</b>	Ethanol 66	Yellow 206	C <sub>21</sub> H <sub>15</sub> NS <sub>2</sub> 345	73.01 73.13	4.38 4.53	4.05 4.23	18.56 18.40	—
<b>8b</b>	Ethanol 69	Yellow 168	C <sub>22</sub> H <sub>17</sub> NOS <sub>2</sub> 375	70.37 70.53	4.56 4.72	3.73 3.55	17.08 17.24	—
<b>8c</b>	Ethanol 69	Yellow 198	C <sub>21</sub> H <sub>14</sub> CINS <sub>2</sub> 379	66.39 66.23	3.71 3.89	3.69 3.86	16.88 17.05	9.33 9.16
<b>10</b>	Ethanol 75	White 168	C <sub>21</sub> H <sub>14</sub> N <sub>2</sub> S 326	77.27 77.11	4.32 4.48	8.58 8.40	9.82 9.98	—

**TABLE II Spectral Data of the Newly Synthesized Compound**

Compound No.	Spectral data, IR (cm <sup>-1</sup> ), <sup>1</sup> H NMR, and <sup>13</sup> C NMR
<b>4a</b>	IR: 1684 (C=O); <sup>1</sup> H NMR: (DMSO-d <sub>6</sub> ) δ 4.36 (s, 2H, CH <sub>2</sub> ) and 6.60–9.93 (m, 13H, Ar-H); <sup>13</sup> C NMR: (DMSO-d <sub>6</sub> ) δ 32.45, 75.13, 122.44, 125.33, 127.69, 128.32, 128.47, 129.87, 136.28, 138.31, 145.62, and 171.39
<b>4b</b>	IR: 1682 (C=O); <sup>1</sup> H NMR: (DMSO-d <sub>6</sub> ) δ 3.57 (s, 3H, OCH <sub>3</sub> ), 4.38 (s, 2H, CH <sub>2</sub> ), and 6.61–7.77 (m, 12H, Ar-H)
<b>4c</b>	IR: 1683 (C=O); <sup>1</sup> H NMR: (DMSO-d <sub>6</sub> ) δ 4.38 (s, 2H, CH <sub>2</sub> ) and 6.99–7.88 (m, 12H, Ar-H)
<b>6a</b>	IR: 1683 (CO); <sup>1</sup> H NMR: (DMSO-d <sub>6</sub> ) δ 7.03–7.81 (m, 19H, CH and Ar-H)
<b>6b</b>	IR: 1684 (C=O); <sup>1</sup> H NMR: (DMSO-d <sub>6</sub> ) δ 3.57 (s, 3H, OCH <sub>3</sub> ) and 6.61–7.77 (m, 18H, CH and Ar-H)
<b>6c</b>	IR: 1684 (C=O); <sup>1</sup> H NMR: (DMSO-d <sub>6</sub> ) δ 6.70–7.80 (m, 18H, CH, and Ar-H)
<b>6d</b>	IR: 1684 (C=O); <sup>1</sup> H NMR: (DMSO-d <sub>6</sub> ) δ 3.57 (s, 3H, OCH <sub>3</sub> ) and 6.75–7.86 (m, 18H, CH, and Ar-H)
<b>6e</b>	IR: 1685 (C=O); <sup>1</sup> H NMR: (DMSO-d <sub>6</sub> ) δ 3.56, 3.57 (s, 6H, OCH <sub>3</sub> ) and 6.71–7.90 (m, 17H, CH and Ar-H)
<b>6f</b>	IR: 1685 (C=O); <sup>1</sup> H NMR (DMSO-d <sub>6</sub> ) δ 3.57 (s, 3H, OCH <sub>3</sub> ) and 6.74–7.93 (m, 17H, Ar-H)
<b>6g</b>	IR: 1684 (C=O); <sup>1</sup> H NMR: (DMSO-d <sub>6</sub> ) δ 7.11–7.86 (m, 18H, CH, and Ar-H)
<b>6h</b>	IR: 1685 (C=O); <sup>1</sup> H NMR: (DMSO-d <sub>6</sub> ) δ 3.57 (s, 3H, OCH <sub>3</sub> ) and 6.71–7.88 (m, 17H, CH, and Ar-H)
<b>6i</b>	IR: 1684 (C=O); <sup>1</sup> H NMR: (DMSO-d <sub>6</sub> ) δ 6.71–7.91 (m, 17H, Ar-H)
<b>7a</b>	IR: 1695 (C=O), 1340, 1070 (SO <sub>2</sub> ); <sup>1</sup> H NMR: (DMSO-d <sub>6</sub> ) δ 5.07 (s, 2H, CH <sub>2</sub> ) and 7.06–7.86 (m, 13H, Ar-H)
<b>7b</b>	IR: 1696 (C=O), 1341, 1072 (SO <sub>2</sub> ); <sup>1</sup> H NMR: (DMSO-d <sub>6</sub> ) δ 3.56 (s, 3H, OCH <sub>3</sub> ), 5.10 (s, 2H, CH <sub>2</sub> ), and 6.80–7.88 (m, 12H, Ar-H)
<b>7c</b>	IR: 1696 (C=O), 1341, 1072 (SO <sub>2</sub> ); <sup>1</sup> H NMR: (DMSO-d <sub>6</sub> ) δ 5.12 (s, 2H, CH <sub>2</sub> ) and 6.95–7.90 (m, 12H, Ar-H)
<b>8a</b>	<sup>1</sup> H NMR: (DMSO-d <sub>6</sub> ) δ 4.95 (s, 2H, CH <sub>2</sub> ) and 7.02–7.91 (m, 13H, Ar-H)
<b>8b</b>	<sup>1</sup> H NMR: (DMSO-d <sub>6</sub> ) δ 3.56 (s, 3H, OCH <sub>3</sub> ), 4.96 (s, 2H, CH <sub>2</sub> ), and 7.10–7.85 (m, 12H, Ar-H)
<b>8c</b>	<sup>1</sup> H NMR: (DMSO-d <sub>6</sub> ) δ 5.0 (s, 2H, CH <sub>2</sub> ) and 7.12–7.90 (m, 12H, Ar-H)
<b>10</b>	<sup>1</sup> H NMR: (DMSO-d <sub>6</sub> ) δ 4.99 (s, 2H, CH <sub>2</sub> ) and 7.33–7.94 (m, 12H, Ar-H); <sup>13</sup> C NMR: (DMSO-d <sub>6</sub> ) 45.28, 48.40, 118.30, 124.49, 127.75, 128.50, 128.70, 130.20, 133.93, 137.18, 138.30, 143.90, and 145.70

### Fluorenespiro[9.2']-3'-aryl-thiazolidin-4-one-1',1'-dioxides 7a-c—General Procedure

To a solution of **4a-c** (10 mmol) in 10 ml of glacial acetic acid was added a saturated solution of potassium permanganate (5 ml) in ice bath for 10 min; and, a saturated solution of sodium hydrogen sulfite was added till the solution was faded. The precipitate so formed was filtered off, washed well with water, and crystallized from the appropriate solvent (cf. Tables I and II).

### Fluorenespiro[9.2']-3'-aryl-thiazolidin-4-thiones 8a-c—General Procedure

A mixture of **4a-c** (10 mmol) and Lawesson's reagent (0.44 g, 10 mmol) in dry toluene (15 mL) was heated at reflux for 2 h. The reaction mixture was filtered off, and the solid so formed was filtered off and recrystallized from methanol (cf. Tables I and II).

### Fluorenespiro[9.2']thiazolo[3',4'-a]benzimidazole (10)—General Procedure

A mixture of 1,2-phenylenediamine (2.16 g, 20 mmol), 9-fluorenone (1.80 g, 10 mmol) (**2**), and 2-mercaptoacetic acid (**3**) (2.76 g, 30 mmol) in 25 ml dry toluene was refluxed for 48 h, and then concentrated till dryness under reduced pressure and the residue was taken-up in ether. The ether layer was washed with sodium bicarbonate solution (1–2%) and finally with H<sub>2</sub>O. The organic layer was dried over sodium sulfate and evaporated to dryness at reduced pressure to afford **10** (cf. Tables I and II).

## REFERENCES

- [1] J. P. Narain, S. P. Tripathy, and E. Pontali, In *Tuberculosis Epidemiology and Control*, J. P. Narain; Ed. WHO Document # SEATB/248, (WHO: 2002) chapter **6**, 83.
- [2] K. Babaoglu, M. A. Page, V. C. Jones, M. R. McNeil, C. Dong, J. H. Naismith, and R. E. Lee, *Bioorg. Med. Chem. Lett.*, **13**, 3227 (2003).
- [3] A. Rao, J. Balzarini, A. Carbone, A. Chimirri, E. De Clercq, A. M. Monforte, P. Monforte, C. Pannecouque, and M. Zappala, *Farmaco*, **59**(1), 33–39 (2004).
- [4] A. Rao, J. Balzarini, A. Carbone, A. Chimirri, E. De Clercq, A. M. Monforte, P. Monforte, C. Pannecouque, and M. Zappala, *Antiviral Res.*, **63**, 97–84 (2004).
- [5] T. Srivastava, W. Haq, and S. B. Katti, *Tetrahedron*, **58**, 7619 (2002).
- [6] A. Chimirri, P. Monforte, L. Musumeci, A. Rao, M. Zappala, and A. M. Monforte, *Archiv der Pharm. Med. Chem.*, **334**, 203–208 (2001).
- [7] A. Chimirri, S. Grasso, P. Monforte, A. Rao, M. Zappala, A. M. Monforte, C. Pannecouque, M. Witvrouw, J. Balzarini, and E. De Clercq, *Antiviral Chem. Chemother.*, **10**, 211 (1999).

- [8] L. N. Lavrova, A. M. Vasil'ev, V. S. Korytnyi, M. K. Indulen, G. M. Ryazantseva, V. Yu. Kovtun, and V. G. Yashunskii, *Pharmaceutical Chem. J.*, **27**, 211–214 (1993).
- [9] S. C. Jain, J. Sinha, S. Bhagat, W. Errington, and C. E. Olsen, *Synthetic Communications*, **33**, 563–577 (2003).
- [10] A. Dandia, R. Singh, and K. Arya, *Phosphorus, Sulfur and Silicon and the Related Elements*, **179**, 551–564 (2004).
- [11] T. Srivastava, A. K. Gaikwad, N. Haq, S. Sinha, and S. B. Katti, *Arkivoc*, 120–130 (2005).
- [12] S. O. Abdalh, H. A. Ead, N. A. Kassab, and N. H. Metwally, *Heterocycles*, **20**, 637 (1983).
- [13] H. A. Ead, S. O. Abdalh, N. A. Kassab, N. H. Metwally, and Y. E. Saleh, *Arch. Pharm. (Weinheim)*, **320**, 1227 (1987).
- [14] H. A. Ead, S. O. Abdalh, N. A. Kassab, and N. H. Metwally, *Sulfur Letters*, **9**, 23 (1989).
- [15] H. A. Ead and N. A. Metwally, *Egypt J. Chem.*, **32**, 691 (1989).
- [16] H. A. Ead, N. H. Metwally, and N. Morsy, *Arch. Pharm. (Weinheim)* **13**, 5 (1990).
- [17] N. H. Metwally, *Sulfur Chemistry*, **28**, 1–11 (2007).
- [18] S. N. Baranov, *Zhur. Obshch. Khim.*, **2**, 512 (1961).
- [19] A. R. Katritzky and A. F. Pozhorskii, *Handbook of Heterocyclic Chemistry* (Pergamon, 2000), 2nd ed., p. 392 (2000).
- [20] A. Dandia, V. Kaur, and P. Singh, *Indian J. of Pharmaceutical Sciences*, **55**, 129–136 (1993).