

Dipyrrolyl Precursors to Bisalkoxide Molybdenum Olefin Metathesis Catalysts

A. S. Hock,[†] Richard R. Schrock,^{*,†} and Amir H. Hoveyda[‡]

Contribution from the Department of Chemistry 6-331, Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, and Department of Chemistry, Merkert Chemistry Center, Boston College, Chestnut Hill, Massachusetts 02467


Received September 12, 2006; E-mail: rrs@mit.edu

Abstract: Addition of 2 equiv of lithium pyrrolide to Mo(NR)(CHCMe₂R')(OTf)₂(DME) (OTf = OSO₂CF₃; R = 2,6-*i*-Pr₂C₆H₃, 1-adamantyl, or 2,6-Br₂-4-MeC₆H₂; R' = Me or Ph) produces Mo(NR)(CHCMe₂R')(NC₄H₄)₂ complexes in good yield. All compounds can be recrystallized readily from toluene or mixtures of pentane and ether and are sensitive to air and moisture. An X-ray structure of a 2,6-diisopropylphenylimido species shows it to be an unsymmetric dimer, {Mo(NAr)(*syn*-CHCMe₂Ph)(η^5 -NC₄H₄)}{Mo(NAr)(*syn*-CHCMe₂Ph)(η^1 -NC₄H₄)}, in which the nitrogen in the η^5 -pyrrolyl bound to one Mo behaves as a donor to the other Mo. All complexes are fluxional on the NMR time scale at room temperature, with one symmetric species being observed on the NMR time scale at 50 °C in toluene-*d*₆. The dimers react with PMe₃ (at Mo) or B(C₆F₅)₃ (at a η^5 -NC₄H₄ nitrogen) to give monomeric products in high yield. They also react rapidly with 2 equiv of monoalcohols (e.g., Me₃COH or (CF₃)₂MeCOH) or 1 equiv of a biphenol or binaphthol to give 2 equiv of pyrrole and bisalkoxide or diolate complexes in ~100% yield.

We have been searching for methods of synthesizing Mo(NR)(CHCMe₂R')(OR'')₂ (R' = Me or Ph) species (or species that contain enantiomerically pure biphenolate or binaphtholate ligands¹) in situ by treating an appropriate Mo(NR)(CHCMe₂R')-X₂ species with a monoalcohol or diol. The main reason is that an increasing number of applications (e.g., asymmetric olefin metathesis¹) require that many catalysts having different combinations of imido and alkoxide ligands be evaluated for a given metathesis transformation and, therefore, that many catalysts be synthesized, isolated, stored, and manipulated. In the long run the synthesis and isolation of many catalysts will be impractical. Of course the synthesis of Mo(NR)(CHCMe₂R')-(OR'')₂ species from Mo(NR)(CHCMe₂R')X₂ species requires that both X groups be replaced readily with OR, that the HX product of this reaction not interfere to any significant degree with subsequent reactions that involve Mo(NR)(CHCMe₂R')-(OR'')₂, and that the HX product not react with any organic species in the reaction. We found that when X = CH₂CMe₃ only 1 equiv of alcohol reacts readily to yield Mo(NAr)(CH-*t*-Bu)(CH₂-*t*-Bu)(OR) or Mo(NAr)(CH₂-*t*-Bu)₃(OR) species.^{2,3} A second approach in which X = NPh₂ allows both X groups to be replaced, but often slowly and incompletely and not at all when NR = NAr (Ar = 2,6-diisopropylphenyl) and the diol is the bulky H₂[Biphen] (H₂[Biphen] = 3,3'-di-*tert*-butyl-5,5',6,6'-

tetramethyl-1,1'-biphenyl-2,2'-diol).⁴ Syntheses of Mo(NR)-(CHCMe₂R')(NPh₂)₂ species from Mo(NR)(CHCMe₂R')(OTf)₂-(dimethoxyethane) species⁵ are also plagued by poor yields as a consequence of competitive deprotonation of the alkylidene. We have now found that a variety of dipyrrolyl complexes, Mo(NR)(CHCMe₂R')(NC₄H₄)₂, can be prepared in good yield from bistriflate precursors and that they react rapidly, even with H₂-[Biphen] when NR = NAr, to yield 2 equiv of pyrrole and bisalkoxide or biphenolate or binaphtholate species.

Addition of 2 equiv of lithium pyrrolide to a stirred diethyl ether suspension of Mo(NR)(CHCMe₂R')(OTf)₂(DME) (OTf = OSO₂CF₃; R = 2,6-*i*-Pr₂C₆H₃ or 1-adamantyl) produces yellow to orange Mo(NR)(CHCMe₂R')(NC₄H₄)₂ complexes in ~75% yield (eq 1). An analogous reaction when R = 2,6-Br₂-4-



MeC₆H₂ is successful when the solvent is a mixture of diethyl ether and dichloromethane. Little or no competitive deprotonation of the alkylidene to give an alkylidyne complex^{6,7} has been observed in any case. All compounds are sensitive to air and moisture and can be recrystallized readily from toluene or mixtures of pentane and ether.

[†] Massachusetts Institute of Technology.

[‡] Boston College.

- (1) Schrock, R. R.; Hoveyda, A. H. *Angew. Chem., Int. Ed.* **2003**, *42*, 4592.
- (2) Sinha, A.; Lopez, L. P. H.; Schrock, R. R.; Hock, A. S.; Müller, P. *Organometallics* **2006**, *25*, 1412.
- (3) Blanc, F.; Copéret, C.; Thivolle-Cazat, J.; Basset, J.-M.; Lesage, A.; Emsley, L.; Sinha, A.; Schrock, R. R. *Angew. Chem., Int. Ed.* **2006**, *45*, 1216.

(4) Sinha, A.; Schrock, R. R.; Müller, P.; Hoveyda, A. H. *Organometallics* **2006**, *25*, 4621–4626.

(5) Schrock, R. R.; Murdzek, J. S.; Bazan, G. C.; Robbins, J.; DiMare, M.; O'Regan, M. *J. Am. Chem. Soc.* **1990**, *112*, 3875.

(6) Schrock, R. R.; Jamieson, J. Y.; Araujo, J. P.; Bonitatebus, P. J. J.; Sinha, A.; Lopez, L. P. H. *J. Organomet. Chem.* **2003**, *684*, 56.

(7) Tonzetich, Z. J.; Schrock, R. R.; Müller, P. *Organometallics* **2006**, *25*, 4301.




Figure 1. Variable temperature proton NMR spectrum of Mo(NAr)(CHCMe₂Ph)(NC₄H₄)₂ in toluene-*d*₈.

All dipyrrolyl complexes are fluxional on the proton NMR time scale. At 22 °C the spectra contain broad resonances, as shown, for example, for Mo(NAr)(CHCMe₂Ph)(NC₄H₄)₂ in toluene-*d*₈ (at 500 MHz) in Figure 1. At high temperature one alkylidene resonance at ~13.3 ppm and two pyrrolyl resonances at ~6.1 and ~6.3 ppm are observed. At low temperatures two alkylidene resonances at ~13.2 and ~13.6 ppm are observed in a 1:1 ratio, and the pyrrolyl proton resonances are resolved into an obscured set of resonances downfield of 6.3 ppm, along with a pattern of four sharp resonances near 5 ppm.⁸ No fluoride resonance is observed in the ¹⁹F NMR spectrum, and no solvent resonances are observed in the ¹H NMR spectrum upon addition of trimethylphosphine, which yields a base adduct (vide infra). A ¹³C NMR spectrum of Mo(NAr)(CHCMe₂Ph)(NC₄H₄)₂ at -50 °C in methylene chloride-*d*₂ reveals resonances at 313.9 ppm (*J*_{CH} = 122.8 Hz) and 293.9 ppm (*J*_{CH} = 121.3 Hz) characteristic of *syn* alkylidene species.⁹

An X-ray structural study of Mo(N-2,6-*i*-Pr₂C₆H₃)(CHCMe₂Ph)(NC₄H₄)₂ shows it to be an unsymmetric dimer, {Mo(NAr)(*syn*-CHCMe₂Ph)(η^5 -NC₄H₄)(η^1 -NC₄H₄)}{Mo(NAr)(*syn*-CHCMe₂Ph)(η^1 -NC₄H₄)₂}, in which the nitrogen in the η^5 -pyrrolyl behaves as a donor to the other Mo (Figure 2). The electron count in the Mo(NAr)(*syn*-CHCMe₂Ph)(η^5 -NC₄H₄)(η^1 -NC₄H₄) half is 18, and in the Mo(NAr)(*syn*-CHCMe₂Ph)(η^1 -




Figure 2. Structure of Mo(NAr)(CHCMe₂Ph)(NC₄H₄)₂. Hydrogen atoms and cocrystallized solvent have been removed for clarity. Thermal ellipsoids are at 50%. Selected bond distances (Å): Mo(1)–C(1A), 1.859(5); Mo(2)–C(1B), 1.912(5); Mo(1)–N(2A), 2.082(4); Mo–N(3A), 2.097(4); Mo(2)–N(2B), 2.060(4); Mo(2)–N(3B), 2.479(4); Mo(1)···Mo(2), 4.53. Selected bond angles (deg): N(1A)–Mo(1)–C(1A), 99.5(2); N(1A)–Mo(1)–N(3B), 155.16(16); N(2A)–Mo(1)–N(3A), 150.98(16); pyrrolyl centroid–Mo(2)–N(1B), 157.3; N(1B)–Mo(2)–C(1B), 100.5(2); N(1B)–Mo(2)–N(2B), 101.80(17).

NC₄H₄)₂(donor) half it is 16. The Mo(NAr)(*syn*-CHCMe₂Ph)(η^1 -NC₄H₄)₂(donor) fragment is approximately a square pyramid with the alkylidene in the apical position. Bond distances and angles are unexceptional. (See figure caption for selected values.) This dimeric structure is consistent with the NMR spectra at low temperature; i.e., one half (containing Mo(2)) has no symmetry, while the second (containing Mo(1)) effectively is *C_s* symmetric. (The asymmetry that is present at Mo(2) apparently cannot be detected at Mo(1), at least under the NMR conditions employed so far.) The four sharp resonances near 5 ppm are assigned to the four protons in the η^5 -NC₄H₄ that is bound to a chiral metal center. η^5 -Pyrrolyl complexes (most of them di- or tetrasubstituted pyrroles¹⁰) have been prepared and studied for many years, the main driving force being the analogy between η^5 -NC₄H₄ and η^5 -C₅H₅.¹¹ To the best of our knowledge, only one other molybdenum pyrrolyl complex, Mo(Tp*)(NO)(η^1 -NC₄H₄)₂ (Tp* = HB(3,5-Me₂C₃N₂H)₃⁻), has been structurally characterized.¹²

(8) In structurally characterized η^5 -pyrrolyl complexes in which the pyrrolyl is disubstituted an upfield shift is observed compared to η^1 -pyrrolyl proton resonances. For example the pyrrole hydrogens in Ti(η^5 -2,5-Me₂C₄H₃N)(NMe₂)₂-Cl are found at 5.93 ppm; Duarte, M. T.; Ferreira, A.; Dias, A. R.; Salema, M. M.; da Silva, J. F. *Acta Crystallogr., Sect. C* **2005**, C61, m104.

(9) (a) Schrock, R. R. *Chem. Rev.* **2002**, 102, 145–180. (b) Feldman, J.; Schrock, R. R. *Prog. Inorg. Chem.* **1991**, 39, 1.


(10) (a) Dias, A. R.; Galvão, A. M.; Galvão, A. C.; Salema, M. S. *J. Chem. Soc., Dalton Trans.* **1997**, 1055. (b) Ascenso, J. R.; Dias, A. R.; Ferreira, A. P.; Galvão, A. C.; Salema, M. S.; Veiros, L. F. *Inorg. Chim. Acta* **2003**, 356, 249.

(11) Kershner, D. L.; Basolo, F. *Coord. Chem. Rev.* **1987**, 79, 279.


(12) Al Obaidi, N.; Brown, K. P.; Edwards, A. J.; Hollins, S. A.; Jones, C. J.; McCleverty, J. A.; Neaves, B. D. *J. Chem. Soc., Chem. Commun.* **1984**, 690.

The NMR spectra at high temperatures are consistent with a C_s symmetric $\text{Mo}(\text{NR})(\text{CHCMe}_2\text{R}')(\eta^1\text{-NC}_4\text{H}_4)_2$ species on the NMR time scale in which the pyrrolyl ligands are η^1 (on average) and rotate rapidly about the Mo–N bonds. Variable temperature spectra are identical at different concentrations, a result that does not reveal whether a small fraction of the dimer breaks up into monomers in which interconversion of $\eta^1\text{-NC}_4\text{H}_4$ and $\eta^5\text{-NC}_4\text{H}_4$ ligands is facile, or whether the equilibration process takes place entirely within the dimer. We favor the former in view of the high reactivity of the $\{\text{Mo}(\text{NR})(\text{CHCMe}_2\text{R}')(\text{NC}_4\text{H}_4)_2\}_2$ species toward alcohols and a Lewis acid or base (vide infra).

Addition of 1 equiv of trimethylphosphine to $\text{Mo}(\text{NAd})(\text{CHCMe}_2\text{Ph})(\text{NC}_4\text{H}_4)_2$ results in immediate formation of *syn*- $\text{Mo}(\text{NAd})(\text{CHCMe}_2\text{Ph})(\eta^1\text{-NC}_4\text{H}_4)_2(\text{PMe}_3)$, in which the alkylidene proton resonance is found at 12.49 ppm with $J_{\text{HP}} = 5$ Hz. An X-ray structural study¹³ shows that trimethylphosphine binds to one of the $\text{CN}_{\text{imido}}\text{N}_{\text{pyrrolyl}}$ faces of the pseudotetrahedral species, which is the face analogous to the CNO face where trimethylphosphine is observed to bind in bisalkoxide species.⁹ The Lewis acid $\text{B}(\text{C}_6\text{F}_5)_3$ also reacts immediately with $\{\text{Mo}(\text{NAr})(\text{CHCMe}_2\text{Ph})(\text{NC}_4\text{H}_4)_2\}_2$ to yield a mixture of what we propose are *syn* and *anti* alkylidenes of the adduct shown in eq 2. The four η^5 -pyrrolyl protons in the major (*syn*) isomer are found at 7.7, 7.2, 5.7, and 5.4 ppm in benzene- d_6 .



Addition of 2 equiv of monoalcohols (e.g., Me_3COH or $(\text{CF}_3)_2\text{MeCOH}$) or 1 equiv of a biphenol or binaphthol to ~ 10 mM solutions of the $\text{Mo}(\text{NR})(\text{CHCMe}_2\text{R}')(\text{NC}_4\text{H}_4)_2$ ($\text{NR} = \text{NAd}$ or NAr) species described above results in rapid formation of 2 equiv of pyrrole and previously characterized bisalkoxide or diolate complexes. The reaction is rapid and gives an $\sim 100\%$ yield in all combinations screened thus far, including the combination of what we consider to be sterically the most challenging, a 2,6-diisopropylphenylimido precursor reacting with $\text{H}_2[\text{Biphen}]$ ($\text{H}_2[\text{Biphen}] = 3,3'$ -di-*tert*-butyl-5,5',6,6'-tetramethyl-1,1'-biphenyl-2,2'-diol (eq 3)). In the case of 3,3'-bis(2,4,6-triisopropylphenyl)-2,2'-binaphthol¹ the resulting bi-



naphtholate appears to bind 1 equiv of pyrrole weakly, but the known THF adduct is generated immediately upon addition of one or more equivalents of THF. Catalysts that have been isolated only as THF adducts, or that have proven to be too unstable to isolate, are likely to be preparable from dipyrrolyl complexes. One example is $\text{Mo}(\text{N-2,6-Br}_2\text{-4-MeC}_6\text{H}_2)(\text{CHCMe}_3)\text{-}[\text{Biphen}]$. Previous attempts to prepare this species through addition of $\text{K}_2[\text{Biphen}]$ to $\text{Mo}(\text{N-2,6-Br}_2\text{-4-MeC}_6\text{H}_2)(\text{CHCMe}_3)(\text{OTf})_2(\text{DME})$ failed to produce the desired species in pure form and in a practical yield.¹⁴ We find that $\text{Mo}(\text{N-2,6-Br}_2\text{-4-MeC}_6\text{H}_2)(\text{CHCMe}_3)(\text{NC}_4\text{H}_4)_2$ reacts with *rac*- $\text{H}_2[\text{Biphen}]$ in benzene rapidly to yield the previously unknown $\text{Mo}(\text{N-2,6-Br}_2\text{-4-MeC}_6\text{H}_2)(\text{CHCMe}_3)[\text{rac-Biphen}]$ species in high yield. The alkylidene proton in $\text{Mo}(\text{N-2,6-Br}_2\text{-4-MeC}_6\text{H}_2)(\text{CHCMe}_3)\text{-}[\text{rac-Biphen}]$ is found at 11.3 ppm with a J_{CH} coupling constant of 132.6 Hz, consistent with a *syn* alkylidene isomer. The catalytic activity of in situ prepared $\text{Mo}(\text{N-2,6-Br}_2\text{-4-MeC}_6\text{H}_2)(\text{CHCMe}_3)[\text{rac-Biphen}]$ was confirmed through the ring-closing metathesis of ~ 80 equiv of diallyl ether to dihydrofuran in 15 min at room temperature in C_6D_6 .

In conclusion we have found that dimeric dipyrrolyl complexes, $\{\text{Mo}(\text{NR})(\text{CHCMe}_2\text{R}')(\text{NC}_4\text{H}_4)_2\}_2$, can be prepared readily and in good yield from $\text{Mo}(\text{NR})(\text{CHCMe}_2\text{R}')(\text{OTf})_2(\text{DME})$ species. All $\{\text{Mo}(\text{NR})(\text{CHCMe}_2\text{R}')(\text{NC}_4\text{H}_4)_2\}_2$ species react rapidly and completely with monoalcohols and diols to yield known and, in one case, an unknown catalyst, even those that contain sterically the most challenging combination of imido, neopentylidene or neophylidene, and diolate ligands. On the basis of these results we expect to be able to prepare catalysts in situ and use them for a wide variety of reactions. We expect that in some cases we can generate relatively unstable catalysts that could not be isolated but that still may be useful for catalytic purposes. We believe the possibilities for rapid screening of known and new catalysts to be significant. We also are exploring the fundamental organometallic chemistry of dipyrrolyl alkylidene complexes and their derivatives.

Acknowledgment. This research was funded by the NIH (GM-59426 to A.H.H. and R.R.S.) and the National Science Foundation (CHE-0138495 and CHE-0554734 to R.R.S.). We thank Dr. Peter Müller for assistance with the X-ray structural solution.

Supporting Information Available: Experimental details for the synthesis of all compounds. Crystal data and structure refinement, atomic coordinates and equivalent isotropic displacement parameters, bond lengths and angles, anisotropic displacement parameters, hydrogen coordinates, and isotropic displacement parameters for $\{\text{Mo}(\text{NAd})(\text{CHCMe}_2\text{Ph})(\eta^1\text{-NC}_4\text{H}_4)_2\}_2$. Data for the structure (06172) are available to the public at <http://www.reciprocalnet.org/>. This material is available free of charge via the Internet at <http://pubs.acs.org>.

JA0665904

(13) Details will be provided in a full paper in due course.

(14) Jamieson, J. Y. Ph.D. thesis, Massachusetts Institute of Technology, 2002.