

Synthesis and Antitumor Activity of Analogues of Ifosfamide Modified in the N-(2-Chloroethyl) Group

Konrad Misiura,[†] Ryszard W. Kinas,[†] Wojciech J. Stec,^{*,†} Halina Kusnierczyk,[‡] Czesław Radzikowski,[‡] and Akio Sonoda[§]

Department of Bioorganic Chemistry, Centre of Molecular and Macromolecular Studies, Polish Academy of Sciences, 90-362 Łódź, Boczna 5, Poland, Institute of Immunology and Experimental Therapy, Polish Academy of Sciences, 53-114 Wrocław, Czerska 12, Poland, and Otsuka Pharmaceutical Co., Ltd., 31, 2-chome, Otedori, Higashi-ku, Osaka, Japan.
Received March 11, 1987

A series of 3-(2-chloroethyl)-N-(2-X-ethyl)tetrahydro-2H-1,3,2-oxazaphosphorin-2-amine 2-oxides with various X substituents have been prepared by cyclization of racemic ifosfamide or its enantiomers with sodium hydride and subsequent treatment of intermediary products with hydrobromic acid, diethyl hydrogen phosphate, dibenzyl hydrogen phosphate, *p*-toluenesulfonic acid, and acetic acid. All of these compounds were tested *in vivo* against L 1210 lymphoid leukemia in mice. Only bromo analogue 13 and its enantiomers were effective, exceeding the activity of racemic ifosfamide and cyclophosphamide. The therapeutic index of the racemic 13 and its levorotatory enantiomer was about 1.7 times higher than that for ifosfamide and about 2.7 times higher than that for cyclophosphamide.

Ifosfamide [IF, 1, N,3-bis(2-chloroethyl)tetrahydro-2H-1,3,2-oxazaphosphorin-2-amine 2-oxide], like isomeric cyclophosphamide [CP, 2, N,N-bis(2-chloroethyl)tetrahydro-2H-1,3,2-oxazaphosphorin-2-amine 2-oxide], has a wide range of experimental and clinical antitumor activity.¹⁻⁴ It is generally accepted that 1 and 2 require metabolic activation,^{5,6} presumably by cytochrome P-450 and mainly in the liver, to exert their cytostatic activity. The principal feature of the metabolic profile^{4,5,7} of 1 (Scheme I) involves the initial formation of 4-hydroxyifosfamide (3), which equilibrates with the acyclic tautomer, aldoifosfamide (4).⁸ The resulting 4 undergoes spontaneous decomposition, or enzyme-assisted hydrolysis,^{9,10} with release of phosphorodiamidic acid 5, which is a lethal alkylating agent.¹¹ Compounds 3 and 4 can undergo further enzymatically mediated oxidation-dehydrogenation to give inactive metabolites 4-ketoifosfamide (6) and carboxyifosfamide (7).^{12,13} In competition with the metabolic reaction 1 → 3, side-chain hydroxylation occurs, which followed by β-elimination of chloroacetic aldehyde yields inactive N-dechloroethylated metabolites 8 and 9.¹⁴

The action mechanism of 5 is thought to resemble that for phosphoramidate mustard.¹⁵ The high alkylating reactivity of 5 may be rationalized in terms of an electronically enhanced rate of intramolecular cyclization of conjugate base 5a to form aziridinium ion 10.¹⁶ The reaction of 10 with nucleophilic groups of biomolecules (HB) gives a monosubstituted intermediate 11, which can, in turn, form a second aziridinium ion and, subsequently, afford a disubstituted product 12.

If the cytotoxic activity of 5 results from alkylation of DNA (biomolecules), any modification of the parent molecule 1, which does not have a significant effect on the metabolic activation process, but which does change the alkylating activity of 5, should affect the antitumor activity of ifosfamide congeners. In this context, we have investigated the effect of replacement of the chlorine atom in the N-(2-chloroethyl) group of IF by an atom or group undergoing nucleophilic substitution on the antitumor activity of the new analogues.

We now report the synthesis and preliminary biological characteristics of 3-(2-chloroethyl)-N-(2-X-ethyl)tetrahydro-2H-1,3,2-oxazaphosphorin-2-amine 2-oxide, where X is Br (bromofosfamide, 13), (C₂H₅O)₂(O)PO (14), (C₆H₅CH₂O)₂(O)PO (15), *p*-CH₃C₆H₄SO₃ (16), CH₃C(O)O (17), and F (18).

H₅CH₂O)₂(O)PO (15), *p*-CH₃C₆H₄SO₃ (16), CH₃C(O)O (17), and F (18).

Results and Discussion

Synthesis of Ifosfamide Analogues. Ifosfamide analogues 13, 14, 15, 16, and 17 have been prepared in two steps (Scheme II) involving intramolecular cyclization of 1 under treatment with sodium hydride,¹⁷ isolation of 2-aziridino-3-(2-chloroethyl)tetrahydro-2H-1,3,2-oxazaphosphorin 2-oxide (19), and subsequent reaction of this last compound with hydrobromic acid, diethyl hydrogen phosphate, dibenzyl hydrogen phosphate, *p*-toluenesulfonic acid, and acetic acid, respectively. Reactions of 19 with the above-mentioned compounds are highly chemoselective and lead to opening of the aziridinyl ring without any observed P-N bond cleavage.^{18,19}


- (1) Schnitker, J.; Brock, N.; Burkert, H.; Fichtner, E. *Arzneim.-Forsch.* **1976**, *26*, 1783.
- (2) Burkert, H.; Voight, H. C. *Proceedings of the International Holoxan Symposium*, Düsseldorf, March 21-23, 1977; Asta-Werke: Bielefeld, F.R.G.
- (3) Hilgard, P.; Herdrich, K.; Brade, W. *Cancer Treat. Rev.* **1983**, *10* (Suppl. A), 183.
- (4) Brade, W. P.; Herdrich, K.; Varini, M. *Cancer Treat. Rev.* **1985**, *12*, 1.
- (5) Connors, T. A.; Cox, P. J.; Farmer, P. B.; Foster, A. B.; Jarman, M. *Biochem. Pharmacol.* **1974**, *23*, 115.
- (6) Brock, N.; Hohorst, H.-J. *Arzneim.-Forsch.* **1963**, *13*, 1021.
- (7) Misiura, K.; Okruszek, A.; Pankiewicz, K.; Stec, W. J.; Czownicki, Z.; Utracka, P. *J. Med. Chem.* **1983**, *26*, 674.
- (8) Ludeman, S. M.; Boyd, V. L.; Regan, J. B.; Gallo, K. A.; Zon, G.; Ishii, K. *J. Med. Chem.* **1986**, *29*, 716.
- (9) Voelcker, G.; Bielicki, L.; Hohorst, H.-J. *J. Cancer Res. Clin. Oncol.* **1981**, *99A*, 58.
- (10) Bielicki, L.; Voelcker, G.; Hohorst, H.-J.; *J. Cancer Res. Clin. Oncol.* **1984**, *107*, 195.
- (11) Bryant, B. M.; Jarman, M.; Baker, M. H.; Smith, I. E.; Smith, J. E. *Cancer Res.* **1980**, *40*, 4734.
- (12) Hill, D. L.; Laster, W. R., Jr.; Kirk, M. C.; ElDareer, S.; Struck, R. F. *Cancer Res.* **1973**, *33*, 1016.
- (13) Takamizawa, A.; Iwata, T.; Matsumoto, S.; *Chem. Pharm. Bull.* **1977**, *25*, 2900.
- (14) Norpoth, K.; Müller, G.; Raidt, H. *Arzneim.-Forsch.* **1976**, *26*, 1376.
- (15) Engle, T. W.; Zon, G.; Egan, W. *J. Med. Chem.* **1982**, *25*, 1347.
- (16) Friedman, O. M. *Cancer Chemother. Rep.* **1967**, *51*, 374.
- (17) Pankiewicz, K.; Kinas, R. W.; Stec, W. J.; Foster, A. B.; Jarman, M.; Van Maanen, J. M. S. *J. Am. Chem. Soc.* **1979**, *101*, 7712.

[†]Centre of Molecular and Macromolecular Studies, Polish Academy of Sciences.


[‡]Institute of Immunology and Experimental Therapy, Polish Academy of Sciences.

[§]Otsuka Pharmaceutical Co., Ltd.

Scheme I



Scheme II



Attempts to open the aziridiny ring in 19 by means of hydroiodic acid or an ethereal solution of hydriodide, although successful, led to highly unstable 3-(2-chloroethyl)-*N*-(2-iodoethyl)tetrahydro-2*H*-1,3,2-oxazaphosphorin-2-amine 2-oxide (20) [³¹P NMR (ether) δ 11.0; TLC (chloroform-ethanol, 9:1) *R_f* 0.42]. Concentration of an ether or chloroform solution of 20 by evaporation of solvent under reduced pressure left dark resinous material. Similar instability of iodo derivatives was observed by Takamizawa²⁰ for 3-(2-chloroethyl)-4-hydroperoxy-*N*-(2-iodoethyl)tetrahydro-2*H*-1,3,2-oxazaphosphorin-2-amine 2-oxide. Treatment of 19 with poly(hydrogen fluoride)-pyridine complex in THF as a solvent did not afford the desired 3-(2-chloroethyl)-*N*-(2-fluoroethyl)tetrahydro-2*H*-1,3,2-oxazaphosphorin-2-amine 2-oxide (18). Instead of 18, 3-(2-chloroethyl)-2-fluorotetrahydro-2*H*-1,3,2-oxazaphosphorin 2-oxide (21) was obtained as the sole product. However, the synthesis of 18 has been achieved by the condensation of phosphorus oxychloride with 2-fluoroethylamine hydrochloride (22) and subsequent condensa-


tion of the resulting *N*-(2-fluoroethyl)phosphoramidic dichloride (23) with 3-aziridinopropanol.

Since enantiomeric forms of ifosfamide (1) became available¹⁷ due to our long-term interest in stereodifferentiated metabolism and antitumor activity of P-stereogenic oxazaphosphorins,^{21,22} we have also prepared the enantiomeric forms of bromofosfamide (13). Cyclization of (+)-1 with sodium hydride and the reaction of a chloroform solution of the resulting 19 with an equimolar amount of 6% aqueous HBr afforded (+)-13. In an analogous way, (-)-1 was converted to (-)-13. Bromofosfamides (+)-13 ($[\alpha]^{25}_D = +41.8^\circ$) and (-)-13 ($[\alpha]^{25}_D = -41.7^\circ$) were enantiomerically pure. The ³¹P NMR spectra of (+)-13 and (-)-13 run in a solution of optically active Eu(tfc)₃ in CCl₄ (compound:Eu(tfc)₃, 1:1.36, molar ratio) indicated the presence of only one enantiomer with induced chemical shifts of δ_+ -65.35 and δ_- -66.90, respectively, whereas the ³¹P NMR spectrum of the mixture of (+)-13 and racemic 13 (85.8:14.2, w/w) indicated the presence of both enantiomers in the ratio 92:8. The assignment of the absolute configuration *S* to the levorotatory and *R* to the dextrorotatory form of 13 was straightforward, because the absolute configuration of ifosfamide was established^{17,23} and the conversions 1 → 19 and 19 → 13 both occur without affecting the sense of chirality at the phosphorus atom (that means without cleavage of any bond attached to the phosphorus atom). Albeit enantiomeric forms of compounds 14-18 are available, their preparation was abandoned in the light of biological test results obtained with racemic compounds 14-18 (vide infra).

Compounds 13-18 were isolated as described in the Experimental Section and are characterized by elemental analyses, mass spectrometry, ¹H NMR, ³¹P NMR, ¹³C NMR, and TLC. Characteristic ¹³C NMR chemical shifts

- (18) Stokes, J. B.; Woods, C. W.; Borckovec, A. B. *Phosphorus Sulfur* 1981, 10, 139.
 (19) Misiura, K.; Silverton, J. V.; Stec, W. J. *J. Org. Chem.* 1985, 50, 1815.
 (20) Takamizawa, A.; Matsumoto, S.; Iwata, T.; Makino, I.; Yamaguchi, K.; Uchida, N.; Kasai, H.; Shiratori, O.; Takase, S. *J. Med. Chem.* 1978, 21, 208.

- (21) Stec, W. J. *Organophosphorus Chem.* 1982, 13, 145.
 (22) Kusnierczyk, H.; Radzikowski, C.; Paprocka, M.; Budzynski, W.; Kinas, R.; Misiura, K.; Stec, W.; *J. Immunopharmacol.* 1986, 8, 455.
 (23) Adamiak, D. A.; Gdaniec, M.; Pankiewicz, K.; Stec, W. J. *Angew. Chem., Int. Ed. Engl.* 1980, 19, 549.

Table I. Selected Carbon-13 Chemical Shifts^a and Coupling Constants^b of Analogues 13–18


compd	C-4	C-5	C-6	C-1'	C-2'	C-1''	C-2''
13	42.0 (4.9)	26.3 (4.9)	67.1 (7.3)	43.0 (4.8)	34.1	47.6	50.1
14	42.0 (3.7)	26.3 (4.2)	67.1 (7.0)	41.4 (6.6)	67.7 (t, 5.7)	47.6	50.1 (2.0)
15	42.1 (3.7)	26.3 (4.2)	67.1 (7.3)	41.5 (7.0)	68.2 (t, 5.7)	47.7	50.2 (2.0)
16	42.1 (3.7)	26.3 (4.9)	67.2 (7.3)	40.4	70.5	47.7	50.0
17	42.0 (4.9)	26.4 (4.9)	67.1 (7.3)	40.3	64.6 (4.9)	47.6	50.1
18	42.1 (4.9)	26.4 (4.9)	67.1 (7.3)	41.6 (20) ^c	83.9 (167) ^c	47.7	50.1

^aChemical shifts are in parts per million with respect to TMS. ^b*J*(PC) values are in parentheses. ^c*J*(FC) values.**Table II.** Antitumor Activity against L 1210 Leukemia in Mice^a

compd	dose, mg/kg	% ILS	survivors on day 60/no. of mice tested (%)
IF (1)	120	74	0/24 (0)
	150	123	4/44 (9)
	180	>250	6/21 (29)
	200	>350	16/38 (42)
	300	>400	17/30 (56)
	400	>500	18/30 (60)
	500	>500	18/30 (60)
CP (2)	600	>480	9/18 (50)
	120	86	1/12 (8)
	150	89	1/19 (5)
	200	>550	8/12 (67)
	300	>550	16/18 (89)
13	400	>425	7/18 (39)
	50	74 ^b	0/7 ^b
	60	88	4/60 (7)
	80	110	7/25 (28)
	100	>430, >200 ^b	24/46, 1/7 ^b (52)
	150	>550	16/23 (70)
	200	>550, >600 ^b	23/27, 7/7 ^b (85)
	300	>550	14/19 (74)
	400	>500	8/12 (67)
	120	114	0/6 (0)
(R)-(+)-13	150	>500	4/6 (67)
	200	>550	10/12 (83)
	300	>550	11/12 (92)
	400	>550	11/12 (92)
	500	206	0/6 (0)
(S)-(-)-13	60	140	1/12 (8)
	100	>550	8/12 (67)
	150	>550	6/6 (100)
	200	>550	12/12 (100)
	300	>350	7/12 (58)
14	400	62	0/6 (0)
	30–180	0–12	0/6
15	30–180	0–12	0/6
16	30–180	13	0/6
17	30–180	13	0/6
18	15–60	10	0/6
	100	–40	0/6
	200	–40	0/6

^aFemale CD2F₁ mice, 10–12 weeks old, were inoculated ip with 10⁵ cells of L 1210 leukemia on day 0. Test preparations were dissolved in sterile saline prior to use and were given ip on day 1 in volume of 0.01 mL/g of body weight. Test and control groups consisted of 6–8 and of 18–24 mice, respectively. ^bData provided by Otsuka Pharmaceutical Co. Preparation was given ip on day 2 to 6-week-old male BDF₁ mice bearing L 1210 leukemia.

δ and coupling constants with ³¹P and ¹⁹F (shown in Table I) were in agreement with the assigned structures. The chemical shifts of the ring carbons C-4, C-5, and C-6 as well as the coupling constants ²*J*(PNC), ²*J*(POC) and ³*J*(PC) are almost the same (within experimental error ± 1.2 Hz), indicating that the tetrahydro-2H-1,3,2-oxazaphosphorin ring adopts the chair conformation.²⁴ Only δ values for

Table III. Toxicity, Effective Doses, and Therapeutic Indices of Bromofosfamidates as Compared to Cyclophosphamide and Ifosfamide

compd	ED ₅₀ ^a mean \pm SD, mg/kg	n ^b	LD ₅₀ ^c mg/kg	TI ^d
IF (1)	88.6 \pm 13.2	7	679 (646–718)	7.7
CP (2)	90.6 \pm 14.4	4	445 (427–465)	4.9
rac-13	36.2 \pm 5.5	9	470 (453–489)	13.0
(R)-(+)-13	66.2 \pm 11.6	5	541 (522–562)	8.2
(S)-(-)-13	27.7 \pm 1.8	5	368 (356–381)	13.3

^aEffective doses providing 50% increase in life span were determined for each therapeutic experiment separately. Appropriate dose ranges, providing % ILS between 15% and 100%, were given to L 1210 leukemia bearing mice (details described in Table II). Since a linear relationship within these dose ranges were observed, each ED₅₀ point was calculated by the method of linear regression analysis of drug dose (four to six levels, six mice/dose) plotted against percentage of ILS. ^bNumber of experiments. ^cLethal doses required to kill 50% of the population were determined after single ip injection of test preparations or reference drugs into healthy female CD2F₁ mice. Dose-level groups providing mortality between 0% and 100% (five to six for one evaluation) consisted of 10 or 15 animals. Observation was carried out for 21 days. The LD₅₀ values and their 95% confidence limits were calculated by using the probit analysis method. ^dTherapeutic index = LD₅₀/ED₅₀.

carbons at positions C-1' and C-2' were affected by the introduction of the X substituent.

Anticancer Screening Data. Cyclophosphamide (2), ifosfamide (1), and the newly synthesized ifosfamide congeners 13–18 were tested for their in vivo anticancer activity against L 1210 lymphoid leukemia in mice, according to NCI standard screening protocols²⁵ at the Institute of Immunology and Experimental Therapy. Some tests were performed at the Otsuka Pharmaceutical Co. From the comparative data included in Table II it can be seen that racemic 13 and enantiomers (R)-(+)-13 and (S)-(-)-13 caused higher than 500% increase in lifespan and cured up to 100% of the mice with L 1210 leukemia. Racemic bromofosfamide and its enantiomers were clearly more effective than IF, and to achieve the same maximal curative effects as provided by IF, an approximately two times lower dose of 13 was required. Moreover, the bromofosfamidates caused a significant number of cures at a wider dose range than CP. In contrast to the bromofosfamidates, compounds 14–18 were inactive in this tumor system.

Effective doses (ED₅₀) of racemic 13, (R)-(+)-13, and (S)-(-)-13 were determined as 36.2, 66.2, and 27.2 mg/kg, respectively (Table III), and were lower than those ob-

(24) Al-Rawi, J. M. A.; Ayed, N.; Kraemer, R.; Navech, J. *Magn. Reson. Chem.* **1985**, *23*, 925.

(25) *In Vivo Cancer Models, 1976–1982*; NIH Publication No. 84–2635, February 1984; p 15.

tained for CP (90.6 mg/kg) or IF (88.6 mg/kg). Although all three bromofosfamides were more toxic than IF, their cancerotoxic selectivity²⁶ expressed as a therapeutic index (TI) was higher (about 1.7 times for 13 and (S)-(-)-13). In comparison to CP, only (S)-(-)-13 was more toxic, but the therapeutic indices of this enantiomer and the racemate were about 2.7 times higher.

Conclusions. The introduction of the bromine substituent instead of chlorine in the *N*-(2-chloroethyl) group of 1 most probably accelerates the rate of aziridinium ion formation²⁷ from the phosphorodiamidic acid, metabolite of 13, and leads to an improved therapeutic index. This finding could suggest that cancerotoxic selectivity is connected not only with the primary activation process (1 → 3, 4 and reaction of 4 with sulfhydryl compounds with formation of 4-thioxazaphosphorins) but also with the toxification process (4 → 5) and subsequent aziridinium ion formation (5 → 10). The increase of the rate of aziridinium ion formation, which is the rate-limiting step of alkylation of biomolecules by 5, could lead to change of the sites and extent of alkylation of nucleophilic centers of DNA, protein, or DNA-polymerase⁶ and, hence, the change of cancerotoxic selectivity. The lack of antitumor activity of compounds 14–18 may be due to their poor tendency, after metabolic transformation, to form aziridinium ion.²⁷ However, other factors, such as rate of oxidation, formation of the aldehydic metabolites, rate of fragmentation, and lipophilicity, might affect the metabolism of compounds 14–18, and should have some consequence on anticancer activity.

Detailed studies on the kinetics of decomposition of 5 and its analogues and their interactions with DNA are in progress.

Experimental Section

General Procedures. Melting points were determined in open glass capillary tubes and are uncorrected. Elemental analyses were performed by CMMS Microanalytical Laboratory. Mass spectra were recorded on a LKB 2091 mass spectrometer at 70 eV ionizing energy. Optical rotations were measured with a Perkin-Elmer 241 MC polarimeter. ¹³C(¹H) NMR and ¹H NMR spectra were recorded at 75.5 and 300 MHz, respectively, with a Bruker MSL-300 instrument. δ values refer to TMS. ³¹P(¹H) NMR spectra were recorded at 24.24 MHz with a JEOL C-60H instrument. Optical purity of enantiomeric bromofosfamides was determined by recording ³¹P NMR spectra at 80.8 MHz with a Varian XL-200 spectrometer. ³¹P NMR δ values refer to external 85% H₃PO₄. Silica gel 60 F₂₅₄ plates were used for thin-layer chromatography; spots were visualized with iodine vapor or 5% ethanolic H₃PO₄·12MoO₃. Silica gel for column chromatography was 230–400 mesh, Merck Art. 9385. Solvents and commercial reagents were distilled and dried by conventional methods before use. Organic extracts were dried over MgSO₄. Enantiomeric forms of ifosfamide,¹⁷ diethyl hydrogen phosphate²⁸ and 3-aziridino-propanol²⁹ were prepared according to the procedures described previously.

2-Aziridino-3-(2-chloroethyl)tetrahydro-2H-1,3,2-oxazaphosphorin 2-Oxide (19) and Its Enantiomers (R)-(+)-19 and (S)-(-)-19. Ethyleneimides 19, (R)-(+)-19, and (S)-(-)-19 were

prepared in the reaction of racemic, dextrorotatory ($[\alpha]^{25}_D = +46.2^\circ$), and levorotatory ($[\alpha]^{25}_D = -45.8^\circ$) ifosfamides, respectively, with sodium hydride in anhydrous THF according to a described procedure.¹⁷ Racemic 19: mp 25–26 °C; ³¹P NMR (CHCl₃) δ 17.8. (R)-(+)-19: mp 64–66 °C; $[\alpha]^{25}_D = +25.5^\circ$ (c 3.0, MeOH); ³¹P NMR (CHCl₃) δ 17.8. (S)-(-)-19: mp 64–66 °C; $[\alpha]^{25}_D = -25.8^\circ$ (c 3.0, MeOH); ³¹P NMR (CHCl₃) δ 17.8.

N-(2-Bromoethyl)-3-(2-chloroethyl)tetrahydro-2H-1,3,2-oxazaphosphorin-2-amine 2-Oxide (13, Bromofosfamide) and Its Enantiomers (R)-(+)-13 and (S)-(-)-13. A water solution of hydrobromic acid (6.6 mL of 6%, 5.0 mmol) was added dropwise to a stirred solution of 19 (1.12 g, 5.0 mmol) in chloroform (10 mL) at room temperature. Stirring was continued for 15 min. The organic layer was separated, and the water layer was extracted twice with chloroform (2 × 5.0 mL). The combined chloroform solutions were dried and concentrated on a rotary evaporator. The solidified material was recrystallized from ether-*n*-pentane to give racemic 13 (1.17 g, 77%): mp 59–60 °C; ³¹P NMR δ 11.4; ¹H NMR (CDCl₃) δ 1.90–2.04 (m, 2), 3.16–3.54 (m, 9), 3.63–3.76 (m, 2), 4.20–4.44 (m, 2); TLC *R*_f 0.44 (CHCl₃-EtOH, 9:1); MS, *m/z* 304 (0.3), 306 (0.3), 225 (30), 83 (84), 31 (100). Anal. (C₇H₁₃BrClN₂O₃P) C, H, Br, Cl, N, P. (R)-(+)-13 (1.1 g, 81%) was obtained by starting from (R)-(+)-19 (1.0 g, 4.45 mmol), $[\alpha]^{25}_D = +25.5^\circ$ (c 3.0, MeOH), and using an analogous procedure: $[\alpha]^{25}_D = +41.8^\circ$ (c 3.0, MeOH); mp 89–90 °C; other parameters (³¹P NMR, ¹H NMR, ¹³C NMR, *R*_f, and MS) as for racemic 13. Anal. C, H, N, P. In the same way, (S)-(-)-19 [1.0 g, 4.45 mmol; $[\alpha]^{25}_D = -25.8^\circ$ (c 3.3, MeOH)] was converted to (S)-(-)-13 (1.1 g, 81%): $[\alpha]^{25}_D = -41.7^\circ$ (c 3.0, MeOH); mp 89–90 °C; other parameters (³¹P NMR, ¹H NMR, ¹³C NMR, *R*_f, and MS) as for racemic 13. Anal. C, H, N, P.

3-(2-Chloroethyl)-N-[2-[(diethoxyphosphinyl)oxy]ethyl]tetrahydro-2H-1,3,2-oxazaphosphorin-2-amine 2-Oxide (14). A solution of diethyl hydrogen phosphate (0.71 g, 5 mmol) in chloroform (2 mL) was added dropwise to a stirred solution of 19 (0.55 g, 2.5 mmol) in CHCl₃ (10 mL) at room temperature. Stirring was continued for 1 h, and then the mixture was diluted with chloroform (8 mL), washed with 10% KHCO₃ (10 mL), dried, and concentrated on a rotary evaporator. The residue was chromatographed on silica gel (25 g) with CHCl₃-EtOH (18:1) as eluent. Product was isolated as a colorless oil: ³¹P NMR (CHCl₃) δ 11.9 (P-2), 1.3 (P-2'); ¹H NMR (CDCl₃) δ 1.34 (t, *J* = 7.1 Hz, 6), 1.89–1.96 (m, 2), 2.15 (s, 1), 3.15–3.29 (m, 5), 3.39–3.46 (m, 1), 3.62–3.68 (m, 2), 4.07–4.37 (m, 8); TLC *R*_f 0.38 (CHCl₃-EtOH, 9:1); MS, *m/z* 381 ([M + 3]⁺, 1.3), 380 ([M + 2]⁺, 1.3), 379 ([M + 1]⁺, 3.4), 378 (M⁺, 2.0), 343 ([M - Cl]⁺, 17), 342 ([M - HCl]⁺, 14), 329 ([M - CH₂Cl]⁺, 7.5), 70 (100). Anal. (C₁₁H₂₅ClN₂O₆P₂) C, H, N, P.

3-(2-Chloroethyl)-N-[2-[[bis(benzyloxy)phosphinyl]oxy]ethyl]tetrahydro-2H-1,3,2-oxazaphosphorin-2-amine 2-Oxide (15). By a procedure essentially identical with that applied above to the diethyl phosphate analogue 14, 19 (0.55 g, 2.5 mmol) and dibenzyl hydrogen phosphate (1.05 g, 3.73 mmol) were converted into 15 (0.9 g, 74%). Product was isolated after column chromatography as colorless oil: ³¹P NMR (CHCl₃) δ 12.0 (P-2), 0.8 (P-2'); ¹H NMR (CDCl₃) δ 1.88–1.94 (m, 2), 3.09–3.28 (m, 6), 3.36–3.43 (m, 1), 3.60–3.64 (m, 2), 4.00–4.33 (m, 4), 5.04–5.12 (m, 4), 7.37 (s, 10); TLC *R*_f 0.33 (CHCl₃-EtOH, 9:1); MS, *m/z* 504 ([M + 2]⁺, 0.3), 502 (M⁺, 0.7), 213 ([M + 2 - CH₂OP(O)(OBz)₂]⁺, 4.6), 211 ([M - CH₂OP(O)(OBz)₂]⁺, 12.4). Anal. (C₂₁H₂₉ClN₂O₆P₂) C, H, N, P.

3-(2-Chloroethyl)-N-[2-(tosyloxy)ethyl]tetrahydro-2H-1,3,2-oxazaphosphorin-2-amine 2-Oxide (16). A solution of *p*-toluenesulfonic acid hydrate (2.0 g, 10.5 mmol) in a chloroform-acetone mixture (3:1, 100 mL) was added dropwise to a stirred solution of 19 (2.24 g, 10.0 mmol) in ether (40 mL) at room temperature. Stirring was continued for 30 min. The mixture was concentrated at reduced pressure, and the residue was dissolved in chloroform (100 mL). The chloroform solution was washed with 5% NaHCO₃ (50 mL) and water (50 mL). The aqueous washings were extracted with chloroform (50 mL), and the combined chloroform solutions were dried and concentrated on a rotary evaporator. The residue was chromatographed on silica gel (90 g) with CHCl₃-EtOH (18:1) as eluent. Product 16 (2.7 g 68%) was isolated as a colorless oil: ³¹P NMR (CHCl₃) δ 11.5; ¹H NMR (CDCl₃) δ 1.88–1.93 (m, 2), 2.34 (s, 1), 2.44 (s, 3),

(26) Brock, N.; *J. Cancer Res. Clin. Oncol.* 1986, 111, 1. Brock, N.; *Cancer Treat. Rev.* 1983, 10 (Suppl. A), 3.

(27) Our unpublished ³¹P NMR kinetic studies (Kinas, R. W.; Pilichowska, S.; Stec, W. J.) of the decomposition of the cyclohexylammonium salt of *N,N'*-bis(2-chloroethyl)phosphorodiamidic acid and *N,N'*-bis(2-bromoethyl)phosphorodiamidic acid have indicated a half-life of ca. 85 and 16 min, respectively (pH 7.4, 37 °C). Under the same experimental conditions,¹⁵ the cyclohexylammonium salt of *N,N'*-bis[2-[(diethoxyphosphinyl)oxy]ethyl]phosphorodiamidic acid does not undergo decomposition.

(28) Zwierzak, A. *Roc.* 1965, 39, 1411.

(29) Okruszek, A. Verkade, J. G. *Phosphorus Sulfur* 1979, 7, 235.

3.10-3.42 (m, 6), 3.58-3.64 (m, 2), 4.04-4.31 (m, 4), 7.35 (d, $J = 8.2$ Hz, 2), 7.78 (d, $J = 8.2$ Hz, 2); TLC R_f 0.48 (CHCl_3 -EtOH, 9:1); MS, m/z 347 ($[\text{M} - \text{CH}_2\text{Cl}]^+$, 1.0), 243 ($[\text{M} + 2 - \text{Ts}]^+$, 1.6), 241 ($[\text{M} - \text{Ts}]^+$, 4.3), 213 ($[\text{M} + 2 - \text{TsOCH}_2]^+$, 34), 211 ($[\text{M} - \text{TsOCH}_2]^+$, 100). Anal. ($\text{C}_{14}\text{H}_{22}\text{ClN}_2\text{O}_5\text{PS}$) C, H, N, P.

N-(2-Acetoxyethyl)-3-(2-chloroethyl)tetrahydro-2H-1,3,2-oxazaphosphorin-2-amine 2-Oxide (17). A solution of 19 (1.1 g, 5.0 mmol) in acetic acid (10 mL) was stirred at room temperature for 1 h and then concentrated under reduced pressure. The solid material was dissolved in chloroform (30 mL), and this solution was washed with 10% KHCO_3 (25 mL). The aqueous layer was extracted twice with chloroform (2×30 mL). The combined chloroform solutions were dried and concentrated on a rotary evaporator. TLC (CHCl_3 -acetone, 1:1) revealed two compounds, unreacted 19 (R_f 0.46) and 17 (R_f 0.25). Elution of the mixture from silica gel (30 g) with CHCl_3 -acetone (1:1) gave unreacted 19 (0.2 g) and 17 (0.6 g, 43%), which was subsequently purified by crystallization from acetone-*n*-pentane to give material (0.5 g) of mp 83-84 °C: ^{31}P NMR (CHCl_3) δ 12.4; ^1H NMR (CDCl_3) δ 1.86-2.18 (m, 2), 2.08 (s, 3), 3.13-3.15 (m, 1), 3.16-3.32 (m, 5), 3.39-3.50 (m, 1), 3.64-3.70 (m, 2), 4.11-4.41 (m, 4); TLC R_f 0.47 (CHCl_3 -EtOH, 9:1); MS, m/z 287 ($[\text{M} + 3]^+$, 0.6), 285 ($[\text{M} + 1]^+$, 1.3), 235 ($[\text{M} - \text{CH}_2\text{Cl}]^+$, 43), 226 ($[\text{M} + 3 - \text{AcO}]^+$, 66), 224 ($[\text{M} + 1 - \text{AcO}]^+$, 20), 215 ($[\text{M} + 2 - \text{AcOCH}_2]^+$, 60), 213 ($[\text{M} - \text{AcOCH}_2]^+$, 20). Anal. ($\text{C}_9\text{H}_{18}\text{ClN}_2\text{O}_4\text{P}$) C, H, N, P.

N-(2-Fluoroethyl)phosphoramidic Dichloride (23). A stirred suspension of 2-fluoroethylamine hydrochloride (5.0 g, 50 mmol) in POCl_3 (25 mL) was heated under reflux for 3 h. The excess of POCl_3 was removed by rotary evaporation, and the residue was distilled under reduced pressure to give 22 (4.6 g, 50%)

as a colorless oil: bp 117-118 °C (7 Pa); ^1H NMR (CDCl_3) δ 3.0-3.9 (m, 2, NCH_2), 4.5 (m, $^2J_{\text{HF}} = 48$ Hz, $^3J_{\text{HH}} = 6$ Hz, $^4J_{\text{HP}} = 2$ Hz, 2, CH_2F), 5.8 (s, 1, NH); ^{31}P NMR (CHCl_3) δ 16.6; MS, m/z 179 (M^+ , 1), 146 ($[\text{M} - \text{CH}_2\text{F}]^+$, 100), 117 ($[\text{M} - \text{FCH}_2\text{CH}_2\text{NH}]^+$, 28).

3-(2-Chloroethyl)-N-(2-fluoroethyl)tetrahydro-2H-1,3,2-oxazaphosphorin-2-amine 2-Oxide (18). To a stirred solution of 23 (3.6 g, 20 mmol) in dioxane (50 mL) was added at room temperature a mixture of 3-aziridinopropanol (2.0 g, 20 mmol), triethylamine (2.2 g, 22 mmol), and dioxane (50 mL). The mixture was stirred for 18 h, then filtered, and concentrated to dryness. A solution of the residue in CHCl_3 (100 mL) was washed with water (2×20 mL), dried, and concentrated under reduced pressure. This material was purified by silica gel (40 g) column chromatography with CHCl_3 -EtOH (18:1) as eluent and recrystallized from Et₂O. Compound 23 (1.2 g) was obtained in 30% yield: mp 56-57 °C; ^{31}P NMR (CHCl_3) δ 12.2; ^1H NMR δ 1.89-2.00 (m, 2), 2.96 (s, 1), 3.16-3.36 (m, 5), 3.44-3.56 (m, 1), 3.63-3.70 (m, 2), 4.19-4.29 (m, 1), 4.34-4.44 (m, 1), 4.48 (m, $J = 48.4$ Hz, $J = 4.8$ Hz, 2); MS, m/z 195 ($[\text{M} - \text{CH}_2\text{Cl}]^+$, 41), 134 (16), 83 (43), 31 (100). Anal. ($\text{C}_7\text{H}_{15}\text{ClFN}_2\text{O}_2\text{P}$) C, H, N, P.

Acknowledgment. This project was financially supported by the Polish National Cancer Program CPBR-11.5.110 and 11.5.112.

Registry No. 13, 104149-14-6; (R)-(+)-13, 104149-16-8; (S)-(-)-13, 104149-15-7; 14, 110971-88-5; 15, 110971-89-6; 16, 110971-90-9; 17, 110971-91-0; 18, 110971-92-1; 19, 29102-47-4; (R)-(+)-19, 72578-73-5; (S)-(-)-19, 72578-74-6; 23, 110971-93-2; 2-fluoroethylamine hydrochloride, 460-08-2; 3-aziridinopropanol, 31190-87-1.

Spiro Hydantoin Aldose Reductase Inhibitors

Reinhard Sarges,* Rodney C. Schnur, John L. Belletire,[†] and Michael J. Peterson

Pfizer Central Research, Groton, Connecticut 06340. Received June 8, 1987

Sorbitol formation from glucose, catalyzed by the enzyme aldose reductase, is believed to play a role in the development of certain chronic complications of diabetes mellitus. Spiro hydantoins derived from five- and six-membered ketones fused to an aromatic ring or ring system inhibit aldose reductase isolated from calf lens. In vivo these compounds are potent inhibitors of sorbitol formation in sciatic nerves of streptozotocinized rats. Optimum in vivo activity is reached in spiro hydantoins derived from 6-halogenated 2,3-dihydro-4H-1-benzopyran-4-ones (4-chromanones). In 2,4-dihydro-6-fluorospiro[4H-1-benzopyran-4,4'-imidazolidine]-2',5'-dione, the activity resides exclusively in the 4S isomer, compound 115 (CP-45,634, USAN: sorbinil). This compound is currently being used to test, in humans, the value of aldose reductase inhibitors in the therapy of diabetic complications.

Aldose reductase, the NADPH-linked enzyme that reduces aldehydic substrates, converts glucose to sorbitol; sorbitol can be converted in turn by polyol dehydrogenase to fructose. Because the K_M of aldose reductase for glucose is high, the hypothesis has been put forward^{1,2} that significant flux through this "polyol pathway" occurs only at high glucose levels, for instance, in diabetics whose plasma and tissue glucose levels may often exceed the normal range.^{2a} Since sorbitol and fructose do not readily diffuse out of cells, it has been postulated that accumulation of sorbitol and fructose in tissues such as lens, nerve, or retina contributes to the development of certain chronic complications of diabetes mellitus, such as cataracts, neuropathies, or retinopathy.^{1,2} If this hypothesis were correct, inhibitors of aldose reductase should be of value in preventing or treating these diabetic complications.

The search for aldose reductase inhibitors with which to test this hypothesis has been under way for many years. In 1965 several simple C_6 - C_8 fatty acids and their derivatives were disclosed as inhibitors of calf lens aldose re-

ductase.³ Tetramethyleneglutaric acid (TMG) was reported in 1968 to be an aldose reductase inhibitor which, in contrast to the C_6 - C_8 fatty acids, was active and also less toxic in lens cultures,^{4a} although TMG failed to dem-

- (1) For recent reviews of the polyol pathway hypothesis, see: (a) Lipinski, C. A.; Hutson, N. J. *Annu. Rep. Med. Chem.* 1984, 19, 169. (b) Kador, P. F.; Robison, W. G.; Kinoshita, J. H. *Annu. Rev. Pharmacol. Toxicol.* 1985, 25, 691. (c) Kador, P. F.; Kinoshita, J. H.; Sharpless, N. E. *J. Med. Chem.* 1985, 28, 841. (d) Benfield, P. *Drugs* 1986, 32 (Suppl. 2), 43. (e) Sarges, R. *Trends in Medicinal Chemistry, Proceedings of the 9th International Symposium on Medicinal Chemistry*, Berlin, 1986; Mutschler, E., Winterfeldt, E., Eds.; VCH: Weinheim, 1987; pp 551-564.
- (2) (a) Van Heyningen, R. *Nature (London)* 1959, 184, 194. (b) Kinoshita, J. H. *Invest. Ophthalmol.* 1965, 4, 786. (c) Gabbay, K. H.; O'Sullivan, J. B. *Diabetes* 1968, 17, 239. (d) Beaumont, P.; Hollows, F. C.; Schofield, P. J.; Williams, J. F.; Steinbeck, W. *Lancet* 1971, 1, 579. (e) Sarges, R. *Prog. Med. Chem.* 1981, 18, 191.
- (3) Hayman, S.; Kinoshita, J. H. *J. Biol. Chem.* 1965, 240, 877.
- (4) (a) Kinoshita, J. H.; Dvornik, D.; Kraml, M.; Gabbay, K. H. *Biochim. Biophys. Acta* 1968, 158, 472. (b) Hutton, J. C.; Schofield, P. J.; Williams, J. F.; Hollows, F. C. *Biochem. Pharmacol.* 1974, 23, 2991.

[†] Current address: Department of Chemistry, University of Cincinnati, Cincinnati, OH 45221.