

165. Synthesis and Ruthenium-Catalyzed Enantioselective Hydrogenation of 3-*O*-Substituted 1,3-Dihydroxypropan-2-ones

Part 2¹⁾

by Edoardo Cesarotti and Patrizia Antognazza

Dipartimento di Chimica Inorganica, Metallorganica e Analitica, Centro CNR, Università di Milano,
via Venezian 21, I-20133 Milano

and Marco Pallavicini* and Luigi Villa

Istituto di Chimica Farmaceutica e Tossicologica, Università di Milano,
viale Abruzzi 42, I-20131 Milano

(30.III.93)


A number of 3-*O*-substituted 1,3-dihydroxypropan-2-ones have been synthesized in view of their potential use as prochiral precursors of optically active glycerols. Indeed, the oxo-ethers have been reduced to the corresponding 3-*O*-substituted glycerols *via* chiral Ru complexes derived from (*S*)-binap, (= (–)-(*S*)-2,2'-bis(diphenylphosphino)-1,1'-binaphthalene) with enantiomeric excesses up to 93%. The steric course of the catalytic reduction appears to be essentially dependent on the steric encumbrance of the substituents; indeed, a dramatic increase of the enantiomeric excess is observed when the bulky trityl group is substituted by the less encumbering benzyl or octadecyl groups.

Introduction. – 3-*O*-Substituted glycerols are versatile intermediates in organic synthesis due to the presence, on a C₃ unit, of three vicinal chemically different O-atoms. Their importance is even more evident, considering that the glycerol skeleton constitutes the basic subunit of many biologically active compounds synthetically obtainable from 3-*O*-substituted glycerols [2], in which the ether acts as a protecting group or a definitive substituent.


On the basis of the reports on enantioselective chemical [3] and biological [4] reductions of α-hydroxy ketones and of our previous investigations concerning the enantioselective hydrogenation of 1,3-*O*-disubstituted 1,3-dihydroxypropan-2-ones [1] [5], the mono-ethers of the same dihydroxy-ketone were considered as potential precursors for optically active 3-*O*-substituted glycerols.

This paper focuses on the preparation of a number of such prochiral ketones, in particular the compounds **1a–e**, which have been selected as target molecules, considering their importance as synthons of the respective glycerol derivatives and on the catalytic reduction with chiral Ru complexes. The catalysts chosen are the dimeric complex [Ru₂((*S*)-binap)₂Cl₄]NEt₃ [6] (**3**) and the 'preformed' [Ru((*S*)-binap)Cl₂] (**4**), recently developed by the *Noyori*'s group [7].

¹⁾ Part 1: [1].



- a** R = PhCH₂
b R = Ph₃C
c R = [(4-MeO)C₆H₄]Ph₂C
d R = C₁₈H₃₇
e R = CH₂=CHCH₂

((*S*)-binap)


- 3** [Ru₂((*S*)-binap)₂Cl₄]NEt₃
4 {Ru((*S*)-binap)Cl₂}_n
 (from [Ru(Benzene)Cl₂]_n and ((*S*)-binap)

Results and Discussion. – The literature methods for the synthesis of 3-*O*-substituted 1,3-dihydroxypropan-2-ones are based on: *a*) the rearrangement of 3-*O*-substituted glyceraldehydes in hot pyridine [8]; *b*) the chemical or biological oxidation of 3-*O*-substituted glycerols [9] [10]; *c*) the replacement of the halogen with an OH moiety in 1-alkoxy-3-chloropropan-2-ones [11] [12]; *d*) the oxidation of glycerol derivatives bearing an ester and an ether group at C(1) and C(3), followed by a sequence of appropriate transformations of the keto, ester, and ether moieties [13]; *e*) the etherification of 1-*O*-trityl-2,2-dimethoxy-3-hydroxypropane, prepared from acetyl dihydroxyacetone dimethyl ketal, and the successive removal of trityl and ketal groups [14].

More simply, our approach involves the 3-*O*-substitution of 1,3-dihydroxypropan-2-one or of its easily obtainable dimethyl ketal. To date, the only example of such a strategy in the literature is the low-yield (17%) monotritylation of 1,3-dihydroxypropan-2-one [15]. Nevertheless, it seemed worthwhile to explore the practicability of this synthetic route further, on account of its attractive shortness.

The sequence of reactions used to synthesize the title compounds is outlined in the *Scheme*.

1,3-Dihydroxypropan-2-one was directly converted to **1b** and **1c** by treatment with trityl chloride and (*p*-anisyl)(chloro)diphenylmethane, respectively, in the presence of pyridine and a catalytic amount of 4-(dimethylamino)pyridine. To prepare **1a**, **1d**, and **1e**, 1,3-dihydroxypropan-2-one was previously transformed into the dimethyl ketal **5** by reaction in MeOH, with TsOH as a catalyst. The successive treatment of **5** with PhCH₂Cl, allyl bromide, and octadecyl methanesulfonate in *t*-BuOH in the presence of *t*-BuOK



gave the mono-ethers **6–8**, which were converted finally into **1a**, **1d**, and **1e** by hydrolysis of the protecting ketal function.

The yields of the 3-*O*-substitution, which represent the critical step of the whole synthetic sequence, range between 45 and 55%.

The results of the enantioselective hydrogenation of 3-*O*-substituted 1,3-dihydroxypropan-2-ones **1a–e** are compiled in the *Table*.

Table. *Enantioselective Hydrogenation of 3-O-Substituted 1,3-Dihydroxypropan-2-ones*^{a)}

Entry	Catalyst	Substrate	Solvent	<i>t</i> [h]	Conversion [%] ^{b)}	e.e. [%] ^{c)}	Configuration ^{d)}
1	3	1a	CH ₂ Cl ₂	24	> 98	87	(<i>R</i>)
2	4	1a	CH ₂ Cl ₂	48	> 98	93	(<i>R</i>)
3	4	1a	EtOH	40	> 98	70	(<i>R</i>)
4	3	1b	CH ₂ Cl ₂	72	> 98	0	–
5 ^{e)}	3	1b	CH ₂ Cl ₂	72	> 98	0	–
6 ^{f)}	4	1b	CH ₂ Cl ₂	70	95	6.5	(<i>R</i>)
7	4	1b	CH ₂ Cl ₂	45	54	0	–
8	3	1c	CH ₂ Cl ₂	90	> 98	0	–
9 ^{f)}	4	1c	CH ₂ Cl ₂	45	50	0	–
10	4	1d	CH ₂ Cl ₂	45	83	76 ^{g)}	(<i>R</i>)
11	4	1e	CH ₂ Cl ₂	90	–	–	–

^{a)} Substrate concentrations 0.1M; catalyst concentration 0.001M; molar substrate/catalyst ratio 100:1; pressure 100 Kgw·cm^{–2} (1 Kgw·cm^{–2} = 9.81 × 10⁴ Pa).

^{b)} Isolated yield after flash chromatography.

^{c)} Determined by HPLC on chiral stationary phase.

^{d)} Determined by the sign of rotation.

^{e)} 2 equiv. of NEt₃ with respect to the catalyst.

^{f)} 1 equiv. of NEt₃ with respect to the catalyst.

^{g)} Specific optical rotation (+)-(*R*)-**2d**: [α]_D²⁰ = +2.36 (*c* = 7, THF) [16].

1-(Benzyloxy)-3-hydroxypropan-2-one (**1a**) is reduced to the corresponding alcohol **2a** by the dimer **3** and by the complex **4** in excellent yields with a remarkably high e.e. % (*Entries 1 and 2*). In EtOH, however, the enantioselectivity is somewhat lower (*Entry 3*), confirming that protic solvents are not the medium of choice when hydrogenating substrates which do not coordinate strongly [5]. A good enantioselectivity is obtained also in the reduction of 3-hydroxy-1-(octadecyloxy)propan-2-one (**1d**; *Entry 10*). Due to the absence of chromophores in **2d**, making UV detection impossible in HPLC, instead of the e.e. %, the optical purity is given, calculated on the specific optical rotation of (+)-(*R*)-**2d**: $[\alpha]_D = +2.36$ ($c = 7$, THF) [16].

Rather surprisingly, the substrates 3-hydroxy-1-trityloxypropan-2-one (**1b**) and 3-hydroxy-1-[(4-methoxyphenyl)diphenylmethoxy]propan-2-one (**1c**) are reduced in high yields but without enantioselectivity (*Entries 4–9*). Only the catalyst **4** reduces **1b** with a moderate 6.5 % e.e. in the presence of 1 equiv. of NEt_3 as co-catalyst (*Entry 6*). It is worth mentioning that, albeit with disappointingly low enantioselectivities, the cleavage of the O-trityl bond in both the substrates **1b** and **1c** occurs in traces in CH_2Cl_2 (less than 2%) both in the presence and in the absence of NEt_3 as cocatalyst.

The substrate 1-allyloxy-3-hydroxypropan-2-one (**1e**) undergoes a different fate: only the C=C bond is selectively reduced after 90 h giving the corresponding 3-hydroxy-1-(propyloxy)propan-2-one (*Entry 11*) in 50 % yield. The reduction of the C=C bond is not accompanied by the hydrogenation of the C=O group as one could expect; indeed it would seem as if the reduction of the C=C bond is followed by the inactivation of the catalyst towards the reduction of the C=O group. Recently, it has been shown that similar catalytic systems can affect ketone reduction without alkene hydrogenation in selected β -keto-esters [17]. Such chemoselectivity is coupled with a high enantioselectivity in spite of the slightly elevated temperatures at which the reactions are performed. All these results seem to confirm our previous observations on the non-innocent role of the substrate in directing the equilibrium mixture of different catalytically active Ru species towards the prevalence of one species over the others [5]. It is also very likely that the different substrates can drive the reaction between the Ru-phosphine complexes and the hydrogen towards the irreversible formation of different catalytic species.

In spite of the fact that α -hydroxy-ketones have been generally considered good substrates in the enantioselective hydrogenation [2] [7], our results seem to indicate that this statement can only be true when the substituents are not particularly sterically hindering; the presence of bulky trityloxy groups in fact leads to a dramatic and apparently inexplicable decrease in the enantioselectivity.

These findings remind one of the earlier observations in the field of asymmetric syntheses catalyzed by transition-metal complexes: the interaction between a chiral complex and a prochiral or achiral substrate consists of a total interaction, reflected in the $\Delta G^\ddagger$, and a part, necessarily lower, which is the diastereotopic interaction $\Delta\Delta G^\ddagger$ [18]. Maximization of the total interaction increasing the hindrance of the substituents, does not affect necessarily the diastereotopic interaction $\Delta\Delta G^\ddagger$ to whom the stereochemical control is related; indeed this term may become so small with respect to the total interaction that the intermediates fail to 'see' a difference in reaction pathway and produce a racemic mixture. Thus, increasing the steric hindrance of the substituents does not necessarily lead to an increased efficiency in the asymmetric synthesis.

We wish to express our sincere thanks to Dr. Suzanne Mulley (University of Liverpool and Milan) for helpful discussions and revision of the manuscript and to Johnson Matthey for the generous loan of RuCl_3 (to E.C.).

Experimental Part

General. (–)-(S)-2,2'-Bis(diphenylphosphino)-1,1'-binaphthalene, ((–)-(S)-binap) is commercially available and was purchased from *Fluka AG*. The Ru complexes used in enantioselective reduction were prepared according to literature procedures: they were the dimer $[\text{Ru}_2((\text{S})\text{-binap})_2\text{Cl}_4]\text{NEt}_3$ (**3**) [6] and the 'preformed' $[\text{Ru}((\text{S})\text{-binap})\text{Cl}_2]_n$ (**4**), prepared from the combination of (S)-binap and $[\text{Ru}(\text{benzene})\text{Cl}_2]_2$ in DMF [7]. All reactions with air- and moisture-sensitive solvents and reagents were carried out under a positive pressure of dry Ar. Solvents were distilled under dry N_2 and degassed with Ar prior to use. Solvent mixtures for TLC and column flash chromatography are specified by volume-to-volume ratio. HPLC analyses were performed on a *Chiralcel-OD* column (250 × 4.6 mm i.d.) from *Daicel*. $^1\text{H-NMR}$ were recorded on a *Bruker WP80* and a *Varian XL200* in CDCl_3 with TMS as an internal standard.

3-O-Substituted 1,3-Dihydroxypropan-2-ones. *1-Hydroxy-3-(triphenylmethoxy)propan-2-one* (**1b**) [15]. Freshly distilled 1,3-dihydroxypropan-2-one (0.52 g, 5.75 mmol), trityl chloride (1.72 g, 6.16 mmol), and 4-(dimethylamino)pyridine (54 mg) were dissolved in pyridine and stirred at r.t. for 15 h. After removing pyridine *in vacuo*, H_2O and AcOEt were added. The org. layer was separated and concentrated. The resulting residue was chromatographed on silica gel (hexane/AcOEt 75:25) yielding 0.86 g (2.59 mmol; 45%) of **1b** as a colorless oil, which solidified on standing. $^1\text{H-NMR}$ (200 MHz, CDCl_3): 3.0 (t, OH); 3.9 (s, CH_2OTr); 4.6 (d, CH_2OH); 7.2–7.5 (m, 3 Ph).

1-Hydroxy-3-[(4-Methoxyphenyl)diphenylmethoxy]propan-2-one (**1c**). Freshly distilled 1,3-dihydroxypropan-2-one (1 g, 11.1 mmol), (*p*-anisyl)(chloro)diphenylmethane (3.43 g, 11.1 mmol) and 4-(dimethylamino)pyridine (100 mg) were dissolved in pyridine and stirred at r.t. for 2 h. Workup was carried out as described for the preparation of **1b** and was followed by chromatography on silica gel (hexane/AcOEt 80:20), giving 1.95 g (5.38 mmol; 48.5%) of **1c** as a colorless oil, which solidified on standing. $^1\text{H-NMR}$ (200 MHz, CDCl_3): 3.0 (t, OH); 3.8 (s, CH_3O); 3.9 (s, $\text{CH}_2\text{OC}(4\text{-MeO-C}_6\text{H}_4)\text{Ph}_2$); 4.6 (d, CH_2OH); 6.85, 7.25–7.45 (d and m, $(4\text{-MeO-C}_6\text{H}_4)\text{Ph}_2$).

2,2-Dimethoxypropane-1,3-diol (**5**). A mixture of 1,3-dihydroxypropan-2-one dimer (25 g, 138.8 mmol), trimethyl orthoformate (30 ml, 0.274 mmol), and TsOH (100 mg) in MeOH (300 ml) was shaken at r.t. overnight. After adding Na_2CO_3 (300 mg), the soln. was concentrated and the resulting residue chromatographed on silica gel ($\text{CH}_2\text{Cl}_2/\text{MeOH}$ 93:7) yielding 28.6 g (210.1 mmol; 75.7%) of **5** as a white waxy solid. $^1\text{H-NMR}$ (200 MHz, CDCl_3): 2.55 (t, 2 OH); 3.3 (s, 2 CH_3); 3.7 (d, 2 CH_2).

3-(Benzyloxy)-2,2-dimethoxypropan-1-ol (**6**). PhCH_2Cl (2.79 ml, 24.3 ml) was added dropwise to a stirred mixture of **5** (3 g, 22.03 mmol) and *t*-BuOK (2.97 g, 26.5 mmol) in *t*-BuOH at 40° over 20 min. The mixture was shaken for 30 min at that temp. and then refluxed for 30 min more. After cooling to r.t., Et_2O , H_2O were added and the layers separated. The aq. phase was extracted twice with Et_2O . The combined Et_2O layers were dried (Na_2SO_4) and evaporated under reduced pressure. The resulting residue was chromatographed on silica gel (cyclohexane/AcOEt 65:35) yielding 2.73 g (12.1 mmol; 54.8%) of **6** as a colorless oil. $^1\text{H-NMR}$ (200 MHz, CDCl_3): 2.0 (t, OH); 3.3 (s, 2 CH_3); 3.55 (s, CH_2OBn); 3.7 (d, CH_2OH); 4.6 (s, CH_2Ph); 7.35 (m, Ph).

2,2-Dimethoxy-3-(octadecyloxy)propan-1-ol (**7**) [19]. Octadecyl methanesulfonate (5.63 g, 16.2 mmol) dissolved in THF was added dropwise to a stirred mixture of **5** (2 g, 14.7 mmol) and *t*-BuOK (1.98 g, 17.6 mmol) in *t*-BuOH at 40°. The mixture was refluxed for 30 min and then cooled to r.t. (*i*-Pr) $_2\text{O}$ and H_2O were added. The upper phase was evaporated and the residue chromatographed on silica gel (hexane/AcOEt 70:30) yielding 2.58 g (6.6 mmol; 45.2%) of **7** as a white solid. $^1\text{H-NMR}$ (80 MHz, CDCl_3): 0.9 (t, CH_2CH_3); 1.3 (br. s, 16 CH_2); 2.3 (OH); 3.3 (s, 2 CH_3O); 3.35–3.8 (CH_2OCH_2 , CH_2OH).

3-(Allyloxy)-2,2-dimethoxypropan-1-ol (**8**). Allyl bromide (1.4 ml, 16.2 mmol) was added dropwise to a stirred mixture of **5** (2 g, 14.7 mmol) and *t*-BuOK (1.98 g, 17.6 mmol) in *t*-BuOH at 40°. The mixture was shaken for 15 min at that temp. and then cooled to r.t. AcOEt and H_2O were added and the layers separated. The aq. phase was extracted twice with AcOEt. The combined org. extracts were dried (Na_2SO_4) and concentrated. The residue was chromatographed on silica gel (hexane/AcOEt 60:40), yielding 1.25 g (7.1 mmol; 48.3%) of **8** as a colorless oil. $^1\text{H-NMR}$ (80 MHz, CDCl_3): 2.65 (OH); 3.3 (s, 2 CH_3); 3.55 (s, $\text{CH}_2\text{O(allyl)}$); 3.7 (s, CH_2OH); 4.05 (d, $\text{CH}_2\text{CH}=\text{CH}_2$); 5.1–5.5 (m, $\text{CH}=\text{CH}_2$); 5.7–6.1 (m, CH).

1-(Benzyloxy)-3-hydroxypropan-2-one (**1a**). The compound **6** (0.55 g, 2.43 mmol) was dissolved in 3N HCl (15 ml) and shaken at r.t. for 30 min. The mixture was neutralized with NaHCO_3 , extracted with AcOEt, and the extract dried (Na_2SO_4). Filtration and removal *in vacuo* of the solvent afforded 0.43 g (2.39 mmol; 98%) of **1a** as a colorless oil, which solidified on standing. $^1\text{H-NMR}$ (200 MHz, CDCl_3): 3.0 (t, OH); 4.2 (s, CH_2OBn); 4.5 (d, CH_2OH); 4.6 (s, CH_2Ph); 7.35 (m, Ph).

1-Hydroxy-3-(octadecyloxy)propan-2-one (**1d**) [14]. The compound **7** (0.98 g, 2.52 mmol) was dissolved in dioxane (20 ml). 3N HCl (3 ml) was added and the mixture stirred at r.t. for 2 h. After adding H_2O and NaHCO_3 (2 g), the mixture was extracted with CHCl_3 . The org. extracts were concentrated *in vacuo* yielding 0.83 g (2.42

mmol; 96%) of **1d** as a white waxy solid. ¹H-NMR (200 MHz, CDCl₃): 0.9 (*t*, CH₃); 1.3 (*br. s*, 15 CH₂); 1.6 (*m*, OCH₂CH₂); 3.05 (*t*, OH); 3.5 (*t*, OCH₂CH₂); 4.15 (*s*, CH₂OC₁₈H₃₇); 4.45 (*d*, CH₂OH).

1-(Allyloxy)-3-hydroxypropan-2-one (1e). The compound **8** (1.2 g, 6.81 mmol) was dissolved in 3*N* HCl (15 ml) and shaken at r.t. for 30 min. Workup as described for the preparation of **1a** gave 0.82 g (6.30 mmol; 93%) of **1e** as a colorless oil. ¹H-NMR (200 MHz, CDCl₃): 3.0 (*br. s*, OH); 4.05 (*dt*, CH₂CH=CH₂); 4.2 (*s*, CH₂O(allyl)); 4.45 (*s*, CH₂OH); 5.15–5.4 (*m*, CH=CH₂); 5.8–6.0 (*m*, CH).

Catalytic Reduction. A weighed amount of catalyst was dissolved in a known amount of the solvent in a *Schlenk* tube under Ar. In another *Schlenk* tube, more than 1 mmol of the substrate was dissolved in the same solvent under Ar. With a syringe, 1.1 mmol of the substrate and 0.011 mmol of the catalyst were transferred into a third *Schlenk* tube, the solvent was added to obtain 11 ml of soln. which was stirred for 30 min under Ar. A stainless steel autoclave was pressurized to 100 Kg/cm² and vented five times; 10 ml of the substrate/catalyst soln. were introduced into the autoclave with a syringe through a serum cap; the autoclave was pressurized to 100 Kg/cm² and stirred in a thermoregulated oil bath at 20°. After the reaction, the autoclave was carefully vented, the resulting orange-yellow soln. was concentrated on a rotatory evaporator, and the oily residue was separated by column flash chromatography. The workup conditions were as follows:

3-(Benzyloxy)propane-1,2-diol (2a) [16]: AcOEt/toluene 85:15. ¹H-NMR (200 MHz, CDCl₃, D₂O): 3.60 (*m*, 4H); 3.80 (*m*, 1H); 4.46 (*s*, 2H); 7.33 (*s*, Ph).

3-(Triphenylmethoxy)propane-1,2-diol (2b) [20]: AcOEt/toluene 1:1. ¹H-NMR (200 MHz, CDCl₃, D₂O): 3.25 (*m*, 2H); 3.6 (*m*, 2H); 3.8 (*m*, 1H); 7.2–7.4 (*m*, 15H).

3-[(4-Methoxyphenyl)diphenylmethoxy]propane-1,2-diol (2c) [21]: AcOEt/toluene 1:1. ¹H-NMR (200 MHz, CDCl₃, D₂O): 3.3 (*m*, 2H); 3.67 (*m*, 1H); 3.80 (*s*, 3H); 3.92 (*m*, 2H); 6.8–6.9 (*d*, 2H); 7.3 (*m*, 12H).

3-(Octadecyloxy)propane-1,2-diol (2d) [16]: AcOEt/toluene 1:1. ¹H-NMR (200 MHz, CDCl₃, D₂O): 0.9 (*t*, 3H); 1.26 (*br. m*, 32H); 3.40–3.70 (*m*, 7H).

1-Hydroxy-3-propyloxypropan-2-one from the reduction of *1-(allyloxy)-3-hydroxypropan-2-one (1e)*. AcOEt/hexane 7:3. ¹H-NMR (200 MHz, CDCl₃, D₂O): 0.9 (*t*, 3H); 1.39 (*q*, 2H); 3.0 (*br. s*, OH); 4.05 (*t*, 2H); 4.2 (*s*, 2H); 4.45 (*s*, 2H).

The enantiomeric composition was determined by HPLC on a chiral stationary phase under the following conditions: **2a–c**: hexane/*i*-PrOH 9:1, flow rate: 1 ml/min.

The enantiomeric composition of **2d** was determined by comparing the optical rotation value with that reported in [16].

REFERENCES

- [1] E. Cesarotti, P. Antognazza, A. Mauri, M. Pallavicini, L. Villa, *Helv. Chim. Acta* **1992**, *75*, 2563.
- [2] J. Jurczak, S. Pikul, T. Bauer, *Tetrahedron* **1986**, *42*, 447.
- [3] R. Nojori, T. Ohkuma, M. Kitamura, H. Takaya, N. Sayo, H. Kumobayashi, S. Akutagawa, T. Ohta, S. Inoue, *J. Am. Chem. Soc.* **1988**, *110*, 629.
- [4] F. Aragozzini, E. Maconi, D. Potenza, C. Scolastico, *Synthesis* **1989**, 225.
- [5] E. Cesarotti, A. Mauri, M. Pallavicini, L. Villa, *Tetrahedron Lett.* **1991**, *32*, 4384.
- [6] T. Ikariya, Y. Ishii, H. Kawano, T. Arai, M. Saburi, S. Yoshikawa, S. Akutagawa, *J. Chem. Soc., Chem. Commun.* **1985**, 922.
- [7] M. Kitamura, M. Tokunaga, T. Ohkuma, R. Nojori, *Tetrahedron Lett.* **1991**, *32*, 4163.
- [8] H. O. L. Fischer, E. Baer, *Chem. Ber.* **1932**, *65B*, 345.
- [9] I. S. Neuberger, *Biochem. Z.* **1931**, *238*, 459.
- [10] I. S. Neuberger, *Biochem. Z.* **1932**, *255*, 6.
- [11] A. Grün, W. Stoll, U.S.P. 2, 374, 283, 1945.
- [12] *J. R. Geigy AG*, Swiss 230, 364, 1943.
- [13] C. Piantadosi, K. Chae, K. S. Ishaq, F. Snyder, *J. Pharm. Sci.* **1973**, *62*, 320.
- [14] C. Piantadosi, K. S. Ishaq, F. Snyder, *J. Pharm. Sci.* **1970**, *59*, 1201.
- [15] L. M. Strawn, R. E. Martell, R. U. Simpson, K. L. Leach, R. E. Counsell, *J. Med. Chem.* **1989**, *32*, 2104.
- [16] G. Hirth, R. Barner, *Helv. Chim. Acta* **1982**, *65*, 1059.
- [17] D. F. Taber, L. J. Silverberg, *Tetrahedron Lett.* **1991**, *32*, 4227.
- [18] B. Bosnich, M. D. Fryzuk, 'Topics in Stereochemistry', Eds. N. L. Allinger and E. L. Eliel, J. Wiley, New York, 1981, Vol. 12.
- [19] C. Piantadosi, K. S. Ishaq, R. L. Wykle, F. Snyder, *Biochemistry* **1971**, *10*, 1417.
- [20] J. Gigg, R. Gigg, *J. Chem. Soc. (C)* **1967**, 1865.
- [21] G. Hirth, H. Saroka, W. Baumwarth, R. Barner, *Helv. Chim. Acta* **1983**, *66*, 1210.