Gold Catalysis

Gold(I)-Catalyzed Asymmetric Cycloisomerization of Eneallenes into Vinylcyclohexenes**

Michael A. Tarselli, Anthony R. Chianese, Stephen J. Lee, and Michel R. Gagné*

The use of gold(I) catalysts for organic synthesis continues to be the subject of considerable attention, as unique reactivity patterns are often observed.^[1] Of the remarkably diverse array of transformations catalyzed by Au^I, the number that have asymmetric variants remains low,^[2-4] most likely as a result of the enforced linear coordination geometry of gold(I), which spatially separates ligand chirality from the reacting substrate.^[2]

The divergent reactivity of Au^{I} catalysts came into view during efforts to logically extend a project on diene cycloisomerization^[5] to 1,6-eneallenes. Although the initially tested Pt^{II} catalysts were not effective, compound **1** was found to react with 10 mol% of the catalyst generated from [(Ph₃P)AuCl] and AgOTf to give vinylcyclohexene **2**,^[6,7] a rare six-membered-ring product of eneallene cycloisomerization [Eq. (1)]. Although numerous studies of eneallene



cycloisomerization reactions have been reported (e.g. with Rh,^[8] Pd,^[9] Ru,^[10] Ni/Cr^[11]), five-membered (typically) and seven-membered (less common) rings are usually obtained, and apparently, none of these reactions tolerates internal substitution at the alkene.^[12] This unknown selectivity for cyclohexene derivatives was therefore pursued.^[13]

Although the reaction rate was good and the PPh₃-based catalyst well-behaved, we decided to use this success as a starting point for the discovery of heretofore unknown enantioselective variants. Since the phosphine ligand provided the obvious point for creating a chiral catalyst environ-



- [**] The authors thank Prof. Ross Widenhoefer (Duke) for a sample of [(R)-tBu₂MeO-BIPHEP(AuCl)₂] and Prof. Wenbin Lin (UNC) for a sample of 4,4'-(TMS)₂binap. Postdoctoral funding from the National Research Council (A.R.C.) and from the National Institutes of General Medicine (GM-60578) is also greatly appreciated.
- Supporting information for this article is available on the WWW under http://www.angewandte.org or from the author.

6670

© 2007 Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim

ment, a preliminary screen of chiral phosphines was initiated. From this examination, $[3,5-xylyl-binap(AuCl)_2]$ (3,5-xylyl-binap = 2,2'-bis(di(3,5-xylyl)phosphino)-1,1'-binaphthyl) was identified as a promising catalyst precursor, and additional optimization studies were carried out with this complex.^[14]

The regioselectivity was moderately dependent on solvent polarity, though the enantioselectivities were strongly dependent on this parameter (Table 1). Attempts to further optimize these two parameters by modifying the counterion were partially successful, but the counterion that delivered the best regioselectivity for the formation of 2 (OTs⁻) was not the same as that which provided the best enantioselectivity (OTf⁻; Table 1).

Table 1: Optimization of reaction conditions for the conversion of 1 to 2a,b using the [(*R*)-3,5-xylyl-binap(AuCl)₂] precursor.^[a]

 MeO_2C CO_2Me

MeO₂C_CO₂Me

MeO₂C₂CO₂Me

	→ 	2a + 2b		
Solvent	Activator	Regioselectivity 2a:2b		
CH ₂ Cl ₂	AgNTf ₂ ^[b]	2.5:1	30	
toluene	AgNTf ₂	1:2	29	
chlorobenzene	AgNTf ₂	2:1	51	
EtNO ₂	AgNTf ₂	3:1	42	
MeNO ₂	AgNTf ₂	4:1	65	
MeNO ₂	AgSbF ₆	4:3	57	
MeNO ₂	AgBF₄	incomplete conversion	30	
MeNO ₂	AgOTs ^[c]	10:1	50	
MeNO ₂	AgPF ₆	9:1	65	
MeNO ₂	AgOTf ^[d]	4:1	72	

[a] Reaction conditions: 5 mol% [(R)-3,5-xylyl-binap(AuCl)₂], 15 mol% AgX, 0.1 \bowtie 1,.RT, 16 h. The best regioselectivites and *ee* values are shown in bold. [b] NTf₂ = N(SO₂CF₃)₂. [c] OTs = OSO₂(4-MeC₆H₄). [d] OTf = OSO₂CF₃. [e] The *ee* values are an average of **2a** and **2b** on chiral-phase GC; see the Supporting Information.

Using the conditions providing optimum enantioselectivities with 3,5-xylyl-binap, a variety of chiral diphosphine ligands were re-examined; however, none provided a better enantioselectivity (Table 2). Interestingly, DTBM-SEG-PHOS [(4,4'-bi-1,3-benzodioxole)-5,5-diylbis(di(3,5-di-*tert*butyl-4-methoxyphenyl)phosphine)], previously shown to be optimum in related gold(I) allene chemistry,^[15] was unreactive in this system. Also intriguing was the reversed regioselectivity in the [QUINAP(AuCl)] case, though the enantioselectivity was poor.

Table 2: Chiral ligand screening for the conversion of 1 to 2a/2b.^[a]

Catalyst	ee [%] ^[b]	Regioselectivity 2a:2b ^[c]
[(S,S)-DIOP(AuCl) ₂] ^[d]	4	1:1
[(R)-QUINAP(AuCl)] ^[e]	10	1:3
[(R)-xylyl-PHANEPHOS(AuCl) ₂] ^[f]	16	2.5:1
[(R)-xylyl-SOLPHOS(AuCl) ₂] ^[g]	28	3.5:1
[(R)-SYNPHOS(AuCl) ₂] ^[h]	33	3:1
$[(R)-4,4'-TMS-binap(AuCl)_2]$	66	1:1.5
[(R)-SEGPHOS(AuCl) ₂] ^[i]	66	3.3:1
$[(R)-3,5-xylyl-binap(AuCl)_2]$	72	3.5:1

[a] Only two products seen by GC after isolation; conversion monitored by ¹H NMR. The best regioselectivity and *ee* are shown in bold. [b] Average of both regioisomers. [c] GC peak integration. [d] DIOP= 1,4-bis (diphenylphosphino)-1,4-dideoxy-2,3-O-isopropylidene-D-threitol. [e] QUINAP=1-(2-diphenylphosphino-1-naphthyl)isoquinoline. [f] xylyl-PHANEPHOS = 4,12-bis[di (3,5-xylyl)phosphino]-[2.2]-paracyclophane. [g] SOLPHOS = 7,7'-bis[di (3,5-xylyl)phosphino]-3,3',4,4'-tetrahydro-4,4'dimethyl-8,8'-bi (2H-1,4-benzoxazine). [h] SYNPHOS = [(5,6),(5',6')bis (ethylenedioxy)biphenyl-2,2'-diyl]bis (diphenylphosphine). [i] SEG-PHOS = 4,4'-bi-1,3-benzodioxole-5,5'-diylbis (diphenylphosphine).

Eneallenes for examining the scope of this chemistry were readily available^[8,16] and enabled several structural variations to be examined. Alkenes that were only monosubstituted at the internal position were unreactive, though substitution at the alkene terminus was tolerated. In the latter cases, this tolerance provided sufficient bias to favor one product regioisomer (10, 12, Table 3). In the case of methallyl- (1) or phenallyl-substituted substrates (7) lacking this element, a mixture of regioisomers was obtained. This result, however, was not universal, as the sulfone and urea variants of the methallyl substrate were quite regioselective for the 1,2product. Substitution at the internal allene hydrogen position, on the other hand, led to lower yields and enantioselectivities, along with competitive isomerization of starting material (not shown).

Enantioselectivities were found to be moderate for most substrates, and a slight increase in *ee* was possible when the reaction was run at -12 °C (77% *ee* for 2). Sulfone 3 was the unusual case, as 4 was obtained in virtually racemic form. Nevertheless, these results represent a significant subset of the known gold-catalyzed asymmetric transformations of allenes.

The skeletal divergence of Au^I in this cycloisomerization suggested a significantly different mechanism from that typically found for Rh,^[8] Pd,^[9] and Ru^[10] catalysts, which are prone to oxidative C–C coupling processes and give *exo,exo*cyclopentene products.^[17] Our current mechanistic hypothesis explaining the observed reactivity patterns involves the electrophilic activation of the internal allene double bond by Au^{I,[6,18]} In this scenario, the alkene acts as a nucleophile to generate putative carbenium ion **A** (Scheme 1); the catalytic cycle is then completed by 2,3- or 2,1-elimination and protodeauration. In cases where C1 and C2 are each substituted, elimination to the more highly substituted alkene product provides the observed major isomer (**10** and **12**). The role of the alkene as a nucleophile also explains why internal disubstitution of the alkene is essential, since this

Substrate	Product	Yield [%] ^[a] (isomer ratio) ^[b]	ee [%] ^[c]
MeO ₂ C_CO ₂ Me	MeO ₂ C CO ₂ Me		
	2a,b	83 (3.5:1)	72 (77) ^{[d}
PhO ₂ S_SO ₂ Ph	PhO ₂ SSO ₂ Ph		
3		82	6
0 II	0		
HNÂNH	HNÂNH		
5	6	82	57 ^[e]
MeO ₂ C_CO ₂ Me	MeO ₂ CCO ₂ Me		
Ph 7	Ph	70 (1.5:1)	45
MeO ₂ C、CO ₂ Me	MeO ₂ C、CO ₂ Me		
e	10	80 (5:1)	59
MeO ₂ C_CO ₂ Me	MeO ₂ C_CO ₂ Me		
		70	65
MeOOMe	MeO OMe		
		77 (1:1)	64
1.4	14 "		

[a] Catalyst generated in situ using standard reaction conditions, see the Experimental Section. [b] 2,1:2,3 olefin isomer. [c] Major regioisomer. [d] When cooled to -12 °C, 0.08 м, 24 h. [e] Poor resolution (chiral-phase supercritical fluid chromatography) limits the accuracy of this measurement ($\pm 10\%$).



Scheme 1. Proposed mechanism for eneallene cycloisomerization. [Au] represents the active catalytic species.

ensures the generation of a tertiary carbenium ion in the putative intermediate. This mechanism is also consistent with the observed regiochemical sensitivity to counterions, since they can presumably act as a weak general bases in the β -elimination step.

Since asymmetric induction in Au^I-catalyzed processes is relatively rare, we obtained a crystal structure of the dinuclear catalyst precursor (Figure 1).^[19,20] Interestingly, a key π - π stacking interaction between two P-bound aryl groups (plane-

Communications



Figure 1. X-ray structure of [3,5-xy|y|-binap $(AuCl)_2]$; one of two molecules in the asymmetric unit is shown; hydrogen atoms are omitted for clarity.

to-plane distance of 3.7 Å) lends the structure of this precursor a degree of rigidity and thus establishes a welldefined chiral environment in the solid state. The same conformational preference was observed in a tol-binap digold structure (tol = tolyl),^[3] which suggests that it may be a common structural feature capable of sculpting the reactive environment of a complex that is otherwise constrained to a linear geometry.^[21] Several observations that complicate the identification of the active catalyst occurred while investigating the cyclization of 1 to 2 with an isolated catalyst. When 5 mol % isolated [(R)-3,5-xylyl-binap(AuOTf)₂] was utilized under the standard conditions, slow reactions and low enantioselectivities were obtained (21% ee, more than 24 h for completion), contrasting with studies of in situ generated catalysts. Suspecting a role for Ag⁺ in these processes, we repeated the reactions using $5 \mod \%$ [(R)-3,5-xylyl-binap-(AuOTf)₂] and a 15 mol% excess of AgOTf. Although a fast rate returned, the enantioselectivity remained low (21%). Unexpectedly, a control reaction using $5 \mod \%$ [(R)-3,5xylyl-binap(AuOTf)₂] and added AgCl (15 mol%) increased the ee to 34%. While we do not yet have an explanation for these observations, the in situ protocol reproducibly provides products with good selectivities.

In conclusion, an enantioselective gold(I) catalyst has been developed for the cycloisomerization of eneallenes into vinylcyclohexene products. This carbon skeleton is unusual, as five- and seven-membered-ring products are the norm with other metal catalysts. The Au¹ catalyst is tolerant of functional groups, it appears to selectively activate the allene over the alkene, and it can be used to generate bicyclic products.

Experimental Section

In a glovebox charged with nitrogen, an oven-dried 1-dram vial was charged with [(R)-3,5-xylyl-binap(AuCl)₂] (5.0 mg, 4.2 mmol), silver triflate (3.0 mg, 12.6 mmol), and nitromethane (0.8 mL). After stirring the suspension for 5 min, **1** (20 mg, 84 mmol) was added. A yellow color was usually noted within 10 min, and the reaction was usually blue-gray upon completion. After consumption of the starting material (determined by ¹H NMR spectroscopy), the reaction mix-

ture was loaded directly onto a silica gel column for purification and eluted with hexanes/ethyl acetate (8:1).

Received: May 3, 2007 Revised: May 31, 2007 Published online: July 27, 2007

Keywords: asymmetric synthesis \cdot C–C coupling \cdot gold \cdot homogeneous catalysis

- [1] For recent reviews see: a) A. Fürstner, P. W. Davies, Angew. Chem. 2007, 119, 3478-3519; Angew. Chem. Int. Ed. 2007, 46, 3410-3449; b) N. Marion, S. P. Nolan, Angew. Chem. 2007, 119, 2806-2809; Angew. Chem. Int. Ed. 2007, 46, 2750-2752; c) A. S. K. Hashmi, G. J. Hutchings, Angew. Chem. 2006, 118, 8064-8105; Angew. Chem. Int. Ed. 2006, 45, 7896-7936; d) S. Ma, S. Yu, Z. Gu, Angew. Chem. 2006, 118, 206-209; Angew. Chem. Int. Ed. 2006, 45, 200-203; e) R. A. Widenhoefer, X. Han, Eur. J. Org. Chem. 2006, 4555-4563; f) C. Bruneau, Angew. Chem. 2005, 117, 2380-2386; Angew. Chem. Int. Ed. 2005, 44, 2328-2334; g) A. Hoffmann-Röder, N. Krause, Org. Biomol. Chem. 2005, 3, 387-391; h) A. S. K. Hashmi, Angew. Chem. 2005, 117, 7150-7154; Angew. Chem. Int. Ed. 2005, 44, 6990-6993; i) A. M. Echavarren, C. Nevado, Chem. Soc. Rev. 2004, 33, 431-436; j) C. Aubert, O. Buisine, M. Malacria, Chem. Rev. 2002, 102, 813-834.
- [2] Enantioselective ene-yne cycloisomerization processes have been reviewed, see: I. J. S. Fairlamb, *Angew. Chem.* 2004, *116*, 1066–1070; *Angew. Chem. Int. Ed.* 2004, *43*, 1048–1052.
- [3] For a recent enantioselective alkyne activation, see: M. Paz Muñoz, J. Adrio, J. C. Carretero, A. M. Echavarren, *Organometallics* 2005, 24, 1293–1300.
- [4] For recent enantioselective allene activations, see: a) C. Liu, R. A. Widenhoefer, Org. Lett. 2007, 9, 1935–1938; b) R. L. LaLonde, B. D. Sherry, E. J. Kang, F. D. Toste, J. Am. Chem. Soc. 2007, 129, 2452–2453; c) Z. Zhang, R. A. Widenhoefer, Angew. Chem. 2007, 119, 287–289; Angew. Chem. Int. Ed. 2007, 46, 283–285.
- [5] J. A. Feducia, A. N. Campbell, M. Q. Doherty, M. R. Gagné, J. Am. Chem. Soc. 2006, 128, 13290–13297.
- [6] For selected examples of gold-catalyzed heteronucleophile addition to allenes, see: a) A. K. Buzas, F. M. Istrate, F. Gagosz, Org. Lett. 2007, 9, 985–988; b) N. Morita, N. Krause, Angew. Chem. 2006, 118, 1930–1933; Angew. Chem. Int. Ed. 2006, 45, 1897–1899; c) B. Gockel, N. Krause, Org. Lett. 2006, 8, 4485–4488; d) A. S. K. Hashmi, M. C. Blanco, D. Fischer, J. W. Bats, Eur. J. Org. Chem. 2006, 1387–1389; e) Z. Zhibin, C. Liu, R. E. Kinder, X. Han, H. Qian, R. A. Widenhoefer, J. Am. Chem. Soc. 2006, 128, 9066–9073; f) N. Nishina, Y. Yamamoto, Angew. Chem. 2006, 118, 3392–3395; Angew. Chem. Int. Ed. 2006, 45, 3314–3317; g) A. Buzas, F. Istrate, F. Gagosz, Org. Lett. 2006, 8, 1957–1959; h) N. Morita, N. Krause, Org. Lett. 2004, 6, 4121–4123; i) A. S. K. Hashmi, Angew. Chem. 2000, 112, 3737–3740; Angew. Chem. Int. Ed. 2000, 39, 3590–3593.
- [7] For examples of gold-catalyzed additions of carbon nucleophiles to allenes, see: a) J. H. Lee, F. D. Toste, Angew. Chem. 2007, 119, 930–932; Angew. Chem. Int. Ed. 2007, 46, 912–914;b) G. Lemiere, V. Gandon, K. Cariou, T. Fukuyama, A.-L. Dhimane, L. Fensterbank, M. Malacria, Org. Lett. 2007, 9, 2207–2209; c) G. Lemiere, V. Gandon, N. Agenet, J.-P. Goddard, A. D. Kozak, C. Aubert, L. Fensterbank, M. Malacria, Angew. Chem. 2006, 118, 7758–7761; Angew. Chem. Int. Ed. 2006, 45, 7596–7599; d) A. Buzas, F. Gagosz, J. Am. Chem. Soc. 2006, 128, 12614–12615; e) Z. Liu, A. Wasmuth, S. G. Nelson, J. Am. Chem. Soc. 2006, 128, 10352–10353. See also footnote [6i].
- [8] T. Makino, K. Itoh, J. Org. Chem. 2004, 69, 395-405.



- [9] K. Närhi, J. Franzén, J.-E. Bäckvall, Chem. Eur. J. 2005, 11, 6937-6943.
- [10] a) C. Mukai, R. Itoh, *Tetrahedron Lett.* 2006, 47, 3971–3974;
 b) S.-K. Kang, B.-S. Ko, D.-M. Lee, *Tetrahedron Lett.* 2002, 43, 6693–6696.
- [11] B. M. Trost, J. M. Tour, J. Am. Chem. Soc. 1988, 110, 5231-5233.
- [12] Itoh reported that internal alkene substituents completely inactivate Rh-catalyzed eneallene cycloisomerizations, see footnote [8].
- [13] Six-membered-ring products have been observed from Pauson-Khand-type reactions and thermal [2+2] cyclizations of eneallenes. See, for example: a) F. Inagaki, C. Mukai, Org. Lett. 2006, 8, 1217–1220; b) A. Padwa, M. Meske, S. S. Murphee, S. H. Watterson, Z. Ni, J. Am. Chem. Soc. 1995, 117, 7071–7080; c) H. Ohno, T. Mizutani, Y. Kadoh, K. Miyamura, T. Tanaka, Angew. Chem. 2005, 117, 5243–5245; Angew. Chem. Int. Ed. 2005, 44, 5113–5115.
- [14] AgNTf₂ has previously been utilized to generate reactive Au¹ cations, see: N. Mézailles, L. Ricard, F. Gagosz, *Org. Lett.* 2005, 7, 4133–4136.
- [15] The [xylyl-binap(Au(3,5-NO₂-benzoate)₂)] catalyst recently reported for the cyclization of allenylsulfonamides was inactive for the reactions described herein (20 h at 50 °C; see footnote [6f]). The *t*Bu₂MeO-BIPHEP digold catalyst reported by Widenhoefer and co-workers for the cyclization of allenyl

alcohols gave full conversion but low *ee* in the present system (see footnote [6e]).

- [16] P. A. Wender, M. P. Croatt, N. M. Deschamp, Angew. Chem. 2006, 118, 2519–2522; Angew. Chem. Int. Ed. 2006, 45, 2459– 2462.
- [17] For a discussion of cycloisomerization mechanisms, see: G. C. Lloyd-Jones, Org. Biomol. Chem. 2003, 1, 215–236.
- [18] A related ionic pathway is proposed for the Au^I-catalyzed asymmetric alkoxy cyclization of ene-ynes, see reference [3].
- [19] For structural information, contact Dr. Peter White in the UNC– Chapel Hill X-ray Facility (pswhite@email.unc.edu).
- [20] Crystal data for $[3,5-xylyl-binap(AuCl)_2]$ at 100 K: $C_{52}H_{48}Au_2Cl_2P_2$, 1/2 H₂O, $M_r = 1200.76 \text{ gmol}^{-1}$, orthorhombic, space group $P2_12_12_1$, a = 12.6526(5), b = 19.0582(7), c = 38.1147(13) Å, V = 9190.8(6) Å³, Z = 8, $\rho_{calcd} = 1.746 \text{ mgm}^{-3}$, $R_1 = 0.0423$ (0.0527), $wR_2 = 0.0933$ (0.0996), for 4688 reflections with $I > 2 \sigma(I)$ (for 26863 reflections ($R_{int} = 0.0604$) with a total of 219861 measured reflections), goodness-of-fit on $F^2 = 1.029$, largest diff peak (hole) = 4.909 (-4.305 e Å^{-3}).
- [21] The L-Au^I-L bond angle is typically in the 170–180° range. See, for example: a) A. A. Mohamed, J. Chen, A. E. Bruce, M. R. M. Bruce, J. A. K. Bauer, D. T. Hill, *Inorg. Chem.* 2003, 42, 2203–2205; b) J. R. Stork, D. Rios, D. Pham, V. Bicocca, M. M. Olmstead, A. L. Balch, *Inorg. Chem.* 2005, 44, 3466–3472.