

Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/gpss20>

A CONVENIENT SYNTHESIS 2-AMINO-5-METHYL-3- NITROBENZENETHIOL AND ITS CONVERSION INTO PHENOTHIAZINES VIA A SMILES REARRANGEMENT

K. G. Ojha^a, Priyanka Mathur^a, Neera Jaisinghani^a
& Purnima Adwani^a

^a Department of Applied Chemistry, M.D.S.
University, Ajmer, 305 001, INDIA

Published online: 24 Sep 2006.

To cite this article: K. G. Ojha, Priyanka Mathur, Neera Jaisinghani & Purnima Adwani (1999) A CONVENIENT SYNTHESIS 2-AMINO-5-METHYL-3-NITROBENZENETHIOL AND ITS CONVERSION INTO PHENOTHIAZINES VIA A SMILES REARRANGEMENT, Phosphorus, Sulfur, and Silicon and the Related Elements, 152:1, 77-81, DOI: [10.1080/10426509908031619](https://doi.org/10.1080/10426509908031619)

To link to this article: <http://dx.doi.org/10.1080/10426509908031619>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and

are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

A CONVENIENT SYNTHESIS OF 2-AMINO-5-METHYL-3- NITROBENZENETHIOL AND ITS CONVERSION INTO PHENOTHIAZINES VIA A SMILES REARRANGEMENT

**K.G. OJHA^{*}, PRIYANKA MATHUR, NEERA JAISINGHANI and
PURNIMA ADWANI**

Department of Applied Chemistry, M.D.S.University AJMER 305 001, INDIA[†]

(Received October 11, 1998; In final form January 23, 1999)

The synthesis of some substituted 3-methyl-1-nitro phenothiazines via a Smiles rearrangement is reported. 2-Amino-5-methyl 3-nitrobenzenethiol was prepared by alkaline hydrolysis of 2-amino-6-methyl-4-nitrobenzothiazole and condensed with o-halonitrobenzenes to obtain diaryl sulphides. They were converted by formic acid into formyl derivatives which by a Smiles rearrangement give the corresponding phenothiazines.

Keywords: phenothiazines; Smiles rearrangement; Antitumor; Alkaline hydrolysis

Phenothiazines^[1] occupy an important place in medicinal chemistry as they find a number of uses as anticancer drugs, tranquilizers, antiinflammatory agents. In addition they exhibit antitumor activity, tuberculostatic activity, and in industry phenothiazene are used as heat stabilizers, antioxidants and analytical reagents.


In connection with our work on phenothiazines we required a reliable and rapid method for the synthesis of substituted 2-aminobenzenethiols. A literature survey showed that in earlier studies reduction of o-nitrophenyl disulphides obtained from o-chloronitrobenzenes and sodium polysulphide was used as a common route to 2-aminobenzenethiols. However this method can not be used to prepare aminonitrothiophenols because of concomitant reduction of the nitro group.

^{*} Correspondence Author.

[†] Address for correspondence B-6 Mayur Colony Alwar gate, Ajmer 305 008 [INDIA]

The Herz reaction has been studied for the synthesis of 2-aminobenz-
enethiols from aromatic amines but it gives only very low yields^[2].

2-Amino-5-methyl-3-nitrobenzenethiol (IV) required in the synthesis of
phenothiazines described in this paper was prepared by hydrolytic cleav-
age of 2-amino-6-methyl-4-nitrobenzothiazole (III) with 50% aqueous
KOH solution. Benzothiazole was obtained by thiocyanogenation of
4-methyl-2-nitroaniline hydrochloride (I) following a process reported
earlier^[3]. (SCHEME 1).



SCHEME 1


3-Methyl-1-nitrophenothiazines (VIII) were prepared by a Smiles rear-
rangement of 5-methyl-2-formamido-2,3-dinitrodiphenyl sulphides (VII)
with alcoholic KOH. Diphenyl sulphides (VII) were obtained by the
formylation of 2-amino-5-methyl-2',3-dinitrodiphenyl sulphides (VI)
which are the condensation products of 2-amino-5-methyl-3-nitroben-
zenethiol (IV) and substituted o-halonitrobenzenes (V) (SCHEME 2)

EXPERIMENTAL

Melting points were taken in open capillaries and are uncorrected. IR spec-
tra have been scanned in KBr. The purity of compounds was checked by
TLC on plates coated with silica gel (0.25 mm) using various nonaqueous
solvent system. All the compounds gave satisfactory C, H & N analyses.

Preparation of 2-amino-5-methyl-3 nitrobenzenethiol (IV)

2-Amino-6-methyl-4-nitrobenzothiazole (III) was prepared by a method
reported earlier (m.p 147°C ; N% Found 20.07, Calcd 20.09; yield 70%). It
was refluxed with aqueous KOH (five times by the weight of benzothia-



SCHEME 2

zole) and water (ten times by the weight of benzothiazole) till evolution of ammonia gas ceased (24 hrs). Contents were diluted with water and filtered, the clear solution was neutralised by acetic acid to get (IV). (M.P. 127°C, yield 32%, N% Found 15.19 Calcd. 15.21).

Preparation of Diaryl Sulphide (VIa-c)

A solution of anhydrous sodium acetate (0.01 mol) in ethanol (5 ml) was added to a hot solution of (IV) (0.012 mol). An alcoholic solution of halo-nitrobenzene (V, 0.01 mol) was then slowly added with stirring. The solution was refluxed for 4 hrs, concentrated and cooled in ice over night.

After filtration the solid was washed with water until the filtrate was colourless. Recrystallisation from methanol afforded the desired product (VI). The physical data are summarized in Table I.

TABLE I

Compound	Yield %	Molecular Formula	M.P.°C	N%	
				Found	Calcd.
VI _a	70	C ₁₃ H ₁₀ N ₃ ClS	193	12.35	12.37
VI _b	65	C ₁₃ H ₁₀ N ₃ BrS	201	10.92	10.93
VI _c	70	C ₁₄ H ₁₃ N ₃ O ₄ S	233	13.15	13.16
VII _a	65	C ₁₄ H ₁₀ N ₃ O ₅ ClS	203	11.43	11.42
VII _b	72	C ₁₄ H ₁₀ N ₃ O ₅ BrS	212	10.18	10.19
VII _c	68	C ₁₅ H ₁₃ N ₃ O ₅ S	257	12.11	12.10
VIII _a	60	C ₁₃ H ₉ N ₂ O ₂ ClS	207	9.56	9.57
VIII _b	65	C ₁₃ H ₉ N ₂ O ₂ BrS	219	8.31	8.30
VIII _c	58	C ₁₄ H ₁₂ N ₂ O ₂ S	280	10.28	10.29

Preparation of 2-formamidodiarly sulphide (VIIa-c)


The diarylsulphides (VI a-c, 0.01 mol) were dissolved in 90% formic acid (20–25 ml) and heated under reflux for 6 hrs and poured on to crushed ice. The solid so obtained was collected, washed and mixed with crushed ice. After collecting the solid was washed with cold water until the filtrate was neutral and recrystallized from benzene (Table I).

Preparation of 1-Nitrophenothiazines (VIIIa-c)

An alcoholic solution of KOH (0.2 gm in 5 ml ethanol) was added to the refluxing solution of the formly derivatives (VII a-c, 0.01 mol) dissolved in acetone (15 ml). The colour of the reaction mixture darkened immediately. After refluxing for half an hour. The second lot of ethanolic solution of KOH (ethanol 5 ml, KOH 0.2 gm) was added and refluxing was continued for 2 hrs. The contents were poured into a beaker containing crushed ice. The precipitate was filtered off, washed with cold water and finally with 30% ethanol then it was recrystallised from benzene to get the desired compounds (VIII a-c).

IR Spectra

In all of the compounds (VIa-c, VIIa-c, VIIIa-c) two peaks were observed in the region 1560–1530 $\{\text{cm}^{-1}\}$ for asymmetric and symmetric valance vibrations of the nitro group. Phenothiazines have been reported to exhibit a single sharp peak at 3450 $\{\text{cm}^{-1}\}$ due to the NH stretching vibration. But 1-nitrophenothiazines exhibit this characteristic peak in the region 3325–3320 $\{\text{cm}^{-1}\}$. This shift to a lower frequency suggests the possibility of a six membered chelate ring of high stability through a strong -NH ---- O = N – intramolecular hydrogen bond which could arise^[4-7] due to the proximity of the NH proton to one of the – NO₂ group at position – 1.



A sharp peak at 745–730 $\{\text{cm}^{-1}\}$ exhibited by (VIa, VIIa, and VIIIa) could be attributed to C-Cl stretching vibrations. All the sulphides (VIa-c) exhibit two peaks in the region of 3380–3360 $\{\text{cm}^{-1}\}$ and 3480–3445 $\{\text{cm}^{-1}\}$ which probably is due to symmetric and asymmetric vibrations of the primary amino group.

Acknowledgements

We are thankful to head of the department of Applied Chemistry, M.D.S. University, AJMER, for providing necessary research facility and encouragement.

References

- [1] R.R. Gupta and M.Kumar; 'Phenothiazines' in 'Phenothiazines and 1,4- Benzothiazines Chemical and Biomedical Aspects' (Ed. R.R. Gupta), Elsevier (1988) pp 1.
- [2] P. Kirby, S.B. Soloway, J.H. Davies and S.B. Webb., J. Chem. Soc. (c) 2250 (1970).
- [3] R.R. Gupta, K.G. Ojha, and M. Kumar, J. Heterocyclic. Chem, 17, 1325 (1980).
- [4] O.L. Brady and C. Waller, Can. J. Chem. Soc., 1218 (1930).
- [5] K.C. Roberts and H.B. Clark, J. Chem. Soc., 1312 (1935).
- [6] O.R. Roding, R.E. Collier, and R.K. Schatzer, J. Med. Chem., 9, 116 (1966).
- [7] L.J. Bellamy (Ed.) The infrared spectra of complex molecules Methuem and Company Ltd., London 1964, pp. 254.