

Accepted Article

Title: Enantioselective Nickel-Catalyzed Intramolecular Allylic Alkenylations Enabled by Reversible Alkenylnickel E/Z Isomerization

Authors: Connor Yap, Gabriel MJ Lenagh-Snow, Somnath N Karad, William Lewis, Louis J Diorazio, and Hon Wai Lam

This manuscript has been accepted after peer review and appears as an Accepted Article online prior to editing, proofing, and formal publication of the final Version of Record (VoR). This work is currently citable by using the Digital Object Identifier (DOI) given below. The VoR will be published online in Early View as soon as possible and may be different to this Accepted Article as a result of editing. Readers should obtain the VoR from the journal website shown below when it is published to ensure accuracy of information. The authors are responsible for the content of this Accepted Article.

To be cited as: Angew. Chem. Int. Ed. 10.1002/anie.201703380 Angew. Chem. 10.1002/ange.201703380

Link to VoR: http://dx.doi.org/10.1002/anie.201703380 http://dx.doi.org/10.1002/ange.201703380

WILEY-VCH

COMMUNICATION

WILEY-VCH

10.1002/anie.201703380

Enantioselective Nickel-Catalyzed Intramolecular Allylic Alkenylations Enabled by Reversible Alkenylnickel *E/Z* Isomerization

Connor Yap, Gabriel M. J. Lenagh-Snow, Somnath Narayan Karad, William Lewis, Louis J. Diorazio, and Hon Wai Lam*

Abstract: Enantioselective nickel-catalyzed arylative cyclizations of substrates containing a *Z*-allylic phosphate tethered to an alkyne are described. These reactions give multisubstituted chiral aza- and carbocycles, and are initiated by the addition of an arylboronic acid to the alkyne, followed by cyclization of the resulting alkenylnickel species onto the allylic phosphate. The reversible *E*/*Z* isomerization of the alkenylnickel species is essential for the success of the reactions.

*E*nantioselective metal-catalyzed allylic substitutions of achiral or racemic substrates using carbon-centered nucleophiles are a major class of reactions for preparing enantioenriched chiral compounds.^[1] Although numerous developments have been described,^[1] there are only a few reports of the enantioselective allylation of alkenyl nucleophiles.^[2] Chiral copper–*N*-heterocyclic carbene catalysts are highly effective in the enantioselective additions of alkenylaluminum,^[2a-c] alkenylboron,^[2d,f,h] and allenylboron^[2e] reagents to achiral allylic phosphates. Chiral iridium^[2g] and rhodium^[2i] catalysts are also effective in enantioselective additions of alkenylboron reagents to racemic allylic alcohols^[2g] and allylic halides, ^[2i] respectively.

While the aforementioned examples provide valuable enantioenriched chiral 1,4-diene building blocks,^[2] several aspects remain underdeveloped. For example, reactions involving fully substituted alkenyl nucleophiles are rare.^[3] The integration of these reactions into domino processes that form more than one new carbon–carbon bond is also not wellestablished.^[3] Murakami and co-workers have partially addressed these issues by developing rhodium-catalyzed cyclizations of 1,6-enynes, in which the reaction is triggered by addition of an arylboronic acid to the alkyne (Scheme 1A).^[3,4] These reactions give cyclopentanes containing a

[*] C. Yap, Dr. S. N. Karad, Prof. H. W. Lam The GSK Carbon Neutral Laboratories for Sustainable Chemistry, University of Nottingham, Jubilee Campus, Triumph Road, NG7 2TU, United Kingdom E-mail: hon.lam@nottingham.ac.uk Homepage: http://www.nottingham.ac.uk/~pczhl/

C. Yap, Dr. G. M. J. Lenagh-Snow, Dr. S. N. Karad, Dr. W. Lewis, Prof. H. W. Lam School of Chemistry, University of Nottingham, University Park,

Nottingham, NG7 2RD, United Kingdom

Dr. L. J. Diorazio AstraZeneca, Pharmaceutical Technology and Development, Etherow F53/20, Charter Way, Macclesfield, Cheshire, SK10 2NA, United Kingdom

۲

Supporting information for this article is available on the WWW under http://dx.doi.org/



B. Cyclization where metal is distal to electrophile (this work)





Scheme 1. Arylative cyclizations of enynes.

tetrasubstituted exocyclic alkene.^[3a,b] However, only two enantioselective reactions were reported, and low selectivity was observed in the initial addition to the alkyne, which led to other products being formed.^[3b] Therefore, the availability of other methods to meet these challenges would significantly enhance the utility of domino alkyne carbometalation–allylic alkenylations.

Here, we describe highly enantioselective intramolecular alkenylations of allylic phosphates with fully substituted alkenylnickel species, which are themselves generated by the nickel-catalyzed addition of arylboronic acids to internal alkynes (Scheme 1B). Notably, this process gives chiral 1,4-diene-containing hetero- and carbocycles that are inaccessible from these substrates with established conditions using rhodium catalysis.^[3] These include 4,5-diaryl 1,2,3,6-tetrahydropyridines, which are seen in naturally occurring alkaloids such as (–)-septicine and (–)-tylophorine (Scheme 1C).^[5]

Our studies began with the arylative cyclization of substrate **1a**, which contains a *Z*-allylic phosphate tethered to an alkyne (Table 1). Guided by our recent work,^[6] we anticipated that arylnickelation of the alkyne to place nickel distal to the electrophile, followed by reversible E/Z isomerization of the alkenylnickel species, would provide a species capable of cyclizing onto the allylic phosphate to give products of formal *anti*-carbometalation (Scheme 1B).^[7] However, of the existing reports of enantioselective nickel-catalyzed allylic substitutions

COMMUNICATION

WILEY-VCH



[a] Reactions were conducted using 0.05 mmol of **1a**. [b] Determined by ¹H NMR analysis of the crude reactions. [c] Determined by ¹H NMR analysis using 1,4-dimethoxybenzene as an internal standard. [d] Determined by HPLC analysis on a chiral stationary phase. [e] Major product was the enantiomer of **2a**. [f] Conducted at 100 °C.

with carbon-centered nucleophiles,^[8,9] none describe the use of alkenyl nucleophiles, and success in this ring-closing step was therefore uncertain.

We were pleased to observe that reaction of 1a with PhB(OH)₂ (2.0 equiv) in the presence of Ni(OAc)₂·4H₂O (10 mol%) and various chiral P,N-ligands (10 mol%) in mixtures of MeCN with THF or 2-MeTHF did indeed provide the anticarbometallative cyclization product 2a. However, these reactions proceeded in low conversions (<10%) and 2a was accompanied by comparable amounts of pyrrolidine 3a, resulting from arylnickelation of the alkyne with the opposite regioselectivity. No reaction was observed in other solvents such dioxane, and toluene. Fortunately, in as DMF, 2,2,2-(TFE), with trifluoroethanol reactions conducted phosphinooxazoline (PHOX) ligands L1-L3^[10] gave 2a in moderate NMR yields and in 36-98% ee, accompanied by only a small quantity of pyrrolidine 3a (entries 1-3).[11] (S,S)-t-Bu-FOXAP (L4) gave no reaction (entry 4). Improved selectivities (>19:1) in favor of 2a were obtained using NeoPHOX ligands^[12] L5 and L6 (entries 5–7).^[13] Although the enantioselectivity was higher using (S)-t-Bu-NeoPHOX (L6) (compare entries 5 and 6), the conversion was modest (entry 6). However, increasing the temperature to 100 °C gave a significantly higher yield of 2a with only a small decrease in enantioselectivity (entry 7). The conditions of entry 7 were subsequently employed to test the generality of this process.[14]

The scope of this reaction with respect to the alkynetethered allylic phosphate was then explored in reactions with PhB(OH)₂, which gave products **2a–2i** in 45–92% yield and 49– 99% ee (Scheme 2). High selectivities (\geq 14:1) in favor of the sixmembered ring products were observed. Regarding the substituent on the alkyne, the reaction is compatible with a phenyl group (**2a**), various *para*- (**2b** and **2c**), *meta*- (**2d**), and *ortho*-substituted benzenes (**2e**), and a 2-thienyl group (**2f**). An



Scheme 2. Scope of 1,6-enynes. Reactions were conducted using 0.30 mmol of **1**. Yields are of isolated products. Values in parentheses refer to the ratio of **2:3** as determined by ¹H NMR analysis of the crude reactions. Unless stated otherwise, the minor isomers **3** were not evident in the isolated products. Enantiomeric excesses were determined by HPLC analysis on a chiral stationary phase. [a] Product **2i** contained trace quantities of inseparable, unidentified impurities, and the ratio of **2i:3i** could not be determined.

alkenyl group on the alkyne is also tolerated, though the product **2g** was formed in 49% *ee*. The reaction of a substrate containing a methyl group on the alkyne gave only a complex mixture of products. Replacement of the *p*-toluenesulfonyl group with a 4-nitrobenzenesulfonyl group is possible (**2h**). Finally, changing the linking group between the alkyne and the allylic phosphate to an all-carbon tether enabled the formation of carbocycle **2i** in 51 yield and 96% *ee*.

Pleasingly, this process is compatible with a range of other (hetero)arylboronic acids, which gave products **2j–2r** in 47–66% yield and 96–99% *ee* from three different 1,6-enynes (Scheme 3). The scope includes *para-* (**2p** and **2q**), *meta-* (**2j** and **2r**), *ortho-* (**2m**), and disubstituted phenylboronic acids (**2k** and **2n**), containing methyl (**2k** and **2p**), halide (**2j**, **2m**, and **2n**), alkoxy (**2n**), nitrile (**2q**), or ester groups (**2r**). In addition, 3-thienylboronic acid (**2l**) and 2-naphthylboronic acid (**2o**) are also tolerated.

Further studies of the scope of the alkyne-tethered allylic phosphate are shown in Eq. (1–3). A trimethylsilyl-substituted alkyne **1j** gave low conversions under the standard conditions, but replacing ligand **L6** with **L5** gave **2s** in 70% yield and 69% ee [Eq. (1)]. The *Z*-stereochemistry of the allylic phosphate appears to be essential, as *E*-allylic phosphate **4** only underwent hydroarylation to give **5** in 63% yield, as a 2:1 ratio of geometric isomers [Eq. (2), stereochemistry of **5** not assigned]. Although the reasons for this observation are not understood, one possibility is that the steric demands of this reaction are better accommodated by *Z*-allylic phosphates. The standard conditions were ineffective in the formation of a cyclopentene from 1,5-enyne **6**, but use of (*R*)-Ph-PHOX (**L1**) in place of **L6** gave **7** in 64% yield and 42% ee [Eq. (3)].

COMMUNICATION



Scheme 3. Scope of boronic acids. Reactions were conducted using 0.30 mmol of **1.** Yields are of isolated products. Values in parentheses refer to the ratio of **2:3** as determined by ¹H NMR analysis of the crude reactions. Unless stated otherwise, the minor isomers **3** were not evident in the isolated products. Enantiomeric excesses were determined by HPLC analysis on a chiral stationary phase. [a] Products **2p–2r** contained trace quantities of inseparable, unidentified impurities, and the ratio of **2:3** could not be determined.



Finally, replacing the arylboronic acid with other pronucleophilic species was examined. Although no reaction occurred with methylboronic acid, (E)-2-phenylvinylboronic acid reacted with **1a** to give **8** in 98% *ee*, albeit in 13% yield [Eq. (4)].

WILEY-VCH



Scheme 4 illustrates a possible catalytic cycle for these reactions, using 1a and PhB(OH)₂ as representative reactants. Transmetalation of PhB(OH)₂ with the chiral nickel species 9, which could have hydroxide, acetate, 2,2,2-trifluoroethoxide, or diethylphosphate ligands resulting from the possible species in the reaction, gives arylnickel species 10. Syn-phenylnickelation of 1a gives alkenylnickel species (E)-11, which undergoes reversible E/Z isomerization to give (Z)-11.[6,7a, 15] The mechanism of *E/Z* isomerization is not currently known, but may involve the intermediacy of zwitterionic carbene-type species.^[15a, 16] Migratory insertion of the alkene of the allylic phosphate into the carbon-nickel bond of (Z)-11 gives alkylnickel species 12, from which β-phosphate elimination would liberate the product 2a, regenerating the nickel species 9. This mechanism for allylic substitution^[17] is similar to that proposed by Murakami and co- workers for arylative cyclizations of 1.6-envnes. [3a,b] Furthermore, it stands in contrast to other related Ni-catalyzed allylic substitutions, which are thought to proceed through allylnickel intermediates. [1j,8j-l, 18] However, although we have proposed that nickel remains in the +2 oxidation state throughout, we do not rule out alternative mechanisms involving Ni(I) species. [7a]



Scheme 4. Postulated catalytic cycle.

In conclusion, we have reported enantioselective nickelcatalyzed allylic alkenylations of allylic phosphates which provide a range of chiral 1,4-diene-containing hetero- and carbocycles in high enantiomeric excesses. This reaction further demonstrates the power of the reversible E/Z isomerization of alkenylnickel species in providing products that would otherwise be inaccessible using *syn*-selective alkyne carbometalation processes.

Received: ((will be filled in by the editorial staff)) Published online on ((will be filled in by the editorial staff))

WILEY-VCH

COMMUNICATION

Acknowledgements

We thank the EPSRC and AstraZeneca (Industrial CASE Studentship to C.Y., grant number EP/N509309/1), the EPSRC Centre for Doctoral Training in Sustainable Chemistry (grant number EP/L015633/1), the European Commission (Marie Skłowdowska-Curie Fellowship to S.N.K., project number 702386), the University of Nottingham, and GlaxoSmithKline for support of this work.

Keywords: allylic substitution • asymmetric catalysis • cyclization • isomerization • nickel

- Selected reviews: a) B. M. Trost, D. L. Van Vranken, *Chem. Rev.* 1996, 96, 395-422. b) B. M. Trost, M. L. Crawley, *Chem. Rev.* 2003, 103, 2921-2944. c) A. Alexakis, J. E. Bäckvall, N. Krause, O. Pàmies, M. Diéguez, *Chem. Rev.* 2008, 108, 2796-2823. d) S. R. Harutyunyan, T. den Hartog, K. Geurts, A. J. Minnaard, B. L. Feringa, *Chem. Rev.* 2008, 108, 2824-2852. e) J. F. Hartwig, L. M. Stanley, *Acc. Chem. Res.* 2010, 43, 1461-1475. f) W.-B. Liu, J.-B. Xia, S.-L. You, *Top. Organomet. Chem.* 2012, 38, 155-207. g) C. Moberg, *Top. Organomet. Chem.* 2012, 38, 209-234. h) J.-B. Langlois, A. Alexakis, *Top. Organomet. Chem.* 2012, 38, 235-268. i) P. Tosatti, A. Nelson, S. P. Marsden, *Org. Biomol. Chem.* 2012, 10, 3147-3163. j) A. H. Cherney, N. T. Kadunce, S. E. Reisman, *Chem. Rev.* 2015, 115, 9587-9652.
- [2] a) Y. Lee, K. Akiyama, D. G. Gillingham, M. K. Brown, A. H. Hoveyda, J. Am. Chem. Soc. 2008, 130, 446-447. b) K. Akiyama, F. Gao, A. H. Hoveyda, Angew. Chem., Int. Ed. 2010, 49, 419-423. c) F. Gao, K. P. McGrath, Y. Lee, A. H. Hoveyda, J. Am. Chem. Soc. 2010, 132, 14315-14320. d) R. Shintani, K. Takatsu, M. Takeda, T. Hayashi, Angew. Chem., Int. Ed. 2011, 50, 8656-8659. e) B. Jung, A. H. Hoveyda, J. Am. Chem. Soc. 2012, 134, 1490-1493. f) F. Gao, J. L. Carr, A. H. Hoveyda, Angew. Chem., Int. Ed. 2012, 51, 6613-6617. g) J. Y. Hamilton, D. Sarlah, E. M. Carreira, J. Am. Chem. Soc. 2013, 135, 994-997. h) F. Gao, J. L. Carr, A. H. Hoveyda, J. Am. Chem. Soc. 2014, 136, 2149-2161. i) M. Sidera, S. P. Fletcher, Nature Chem. 2015, 7, 935-939.
- a) T. Miura, M. Shimada, M. Murakami, J. Am. Chem. Soc. 2005, 127, 1094-1095. b) T. Miura, M. Shimada, M. Murakami, Chem. Asian. J 2006, 1, 868-877. For a related process, see: c) M. Shimada, T. Harumashi, T. Miura, M. Murakami, Chem. Asian. J 2008, 3, 1035-1040.
- For reviews of domino carbometalation sequences triggered by the rhodium-catalyzed addition of organoboron reagents, see: (a) T. Miura, M. Murakami, *Chem. Commun.* 2007, 217-224. (b) S. W. Youn, *Eur. J. Org. Chem.* 2009, 2597-2605.
- [5] D. L. Comins, X. Chen, L. A. Morgan, J. Org. Chem. 1997, 62, 7435-7438.
- [6] C. Clarke, C. A. Incerti-Pradillos, H. W. Lam, J. Am. Chem. Soc. 2016, 138, 8068-8071.
- [7] For related Ni-catalyzed *anti*-carbometallative cyclizations, see: a) X.
 Zhang, X. Xie, Y. Liu, *Chem. Sci.* 2016, 7, 5815-5820. b) X. Wang, Y.
 Liu, R. Martin, *J. Am. Chem. Soc.* 2015, 137, 6476-6479. c) M.
 Börjesson, T. Moragas, R. Martin, *J. Am. Chem. Soc.* 2016, 138, 7504-7507.
- [8] Selected examples of enantioselective Ni-catalyzed allylic alkylations or arylations: a) G. Consiglio, F. Morandini, O. Piccolo, *Helv. Chim. Acta* 1980, 63, 987-989. b) G. Consiglio, O. Piccolo, L. Roncetti, F. Morandini, *Tetrahedron* 1986, 42, 2043-2053. c) T. Hiyama, N. Wakasa, *Tetrahedron Lett.* 1985, 26, 3259-3262. d) G. Consiglio, A. Indolese, *Organometallics* 1991, 10, 3425-3427. e) A. F. Indolese, G. Consiglio, *Organometallics* 1994, 13, 2230-2234. f) U. Nagel, H. G. Nedden, *Inorg. Chim. Acta* 1998, 269, 34-42. g) E. Gomez-Bengoa, N. M. Heron, M. T. Didiuk, C. A. Luchaco, A. H. Hoveyda, *J. Am. Chem. Soc.* 1998, 120, 7649-7650. h) K.-G. Chung, Y. Miyake, S. Uemura, *J. Chem. Soc. Perkin Trans* 1 2000, 15-18. i) S. Son, G. C. Fu, *J. Am. Chem. Soc.* 2008, 130, 2756-2757. j) J. D. Shields, D. T. Ahneman, T. J. A. Graham, A. G. Doyle, *Org. Lett.* 2014, 16, 142-145. k) Z. Zeng, D. Yang, Y. Long, X. Pan, G. Huang, X. Zuo, W. Zhou, *J. Org. Chem.* 2014, 79, 5249-

5257. I) Y. Kita, R. D. Kavthe, H. Oda, K. Mashima, *Angew. Chem., Int. Ed.* **2016**, *55*, 1098-1101.

- [9] For examples of enantiospecific rather than enantioselective Nicatalyzed allylic or benzylic substitutions, see: a) B. L. H. Taylor, E. C. Swift, J. D. Waetzig, E. R. Jarvo, J. Am. Chem. Soc. 2011, 133, 389-391. b) B. L. H. Taylor, M. R. Harris, E. R. Jarvo, Angew. Chem., Int. Ed. 2012, 51, 7790-7793. c) M. R. Harris, L. E. Hanna, M. A. Greene, C. E. Moore, E. R. Jarvo, J. Am. Chem. Soc. 2013, 135, 3303-3306. d) Q. Zhou, H. D. Srinivas, S. Dasgupta, M. P. Watson, J. Am. Chem. Soc. 2013, 135, 3307-3310. e) H. M. Wisniewska, E. C. Swift, E. R. Jarvo, J. Am. Chem. Soc. 2013, 135, 9083-9090. f) I. M. Yonova, A. G. Johnson, C. A. Osborne, C. E. Moore, N. S. Morrissette, E. R. Jarvo, Angew. Chem., Int. Ed. 2014, 53, 2422-2427. g) H. D. Srinivas, Q. Zhou, M. P. Watson, Org. Lett. 2014, 16, 3596-3599. h) E. J. Tollefson, D. D. Dawson, C. A. Osborne, E. R. Jarvo, J. Am. Chem. Soc. 2014, 136, 14951-14958. i) E. J. Tollefson, L. E. Hanna, E. R. Jarvo, Acc. Chem. Res. 2015, 48, 2344-2353.
- [10] a) P. von Matt, A. Pfaltz, *Angew. Chem., Int. Ed. Engl.* **1993**, *32*, 566-568. b) J. Sprinz, G. Helmchen, *Tetrahedron Lett.* **1993**, *34*, 1769-1772. c) J. V. Allen, S. J. Coote, G. J. Dawson, C. G. Frost, C. J. Martin, J. M. J. Williams, *J. Chem. Soc., Perkin Trans.* **1 1994**, 2065-2072.
- [11] The exact reasons for the beneficial effect of TFE are not known at the present time, but one possibility is that TFE engages in hydrogenbonding with the phosphate, activating it as a leaving group in allylic substitution. Use of TFE in combination with other solvents such as THF, dioxane, or MeCN gave much lower yields.
- [12] M. G. Schrems, A. Pfaltz, Chem. Commun. 2009, 6210-6212.
- [13] The absolute configuration of 2a obtained using ligand L5 was determined by X-ray crystallography, and the absolute configurations of the remaining products were assigned by analogy. CCDC 1539326 contains the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Centre.
- [14] Repeating the experiment of Table 1, entry 7, but in the absence of L6, gave none of 2a. Instead, 3a was formed in 12% NMR yield, along with a 10% NMR yield of what appeared to be the product of hydroarylation of 1a without cyclization.
- [15] a) J. M. Huggins, R. G. Bergman, J. Am. Chem. Soc. 1981, 103, 3002-3011. b) A. Yamamoto, M. Suginome, J. Am. Chem. Soc. 2005, 127, 15706-15707. c) M. Daini, A. Yamamoto, M. Suginome, Asian. J. Org. Chem. 2013, 2, 968-976.
- a) M. Murakami, T. Yoshida, S. Kawanami, Y. Ito, *J. Am. Chem. Soc.* **1995**, *117*, 6408-6409. b) P.-S. Lin, M. Jeganmohan, C.-H. Cheng, *Chem. Eur. J.* **2008**, *14*, 11296-11299. c) T. Sperger, C. M. Le, M. Lautens, F. Schoenebeck, *Chem. Sci.* **2017**, *8*, 2914-2922.
- [17] Sawamura and co-workers have described stereospecific Pd-catalyzed reactions between allylic esters and arylboronic acids, in which allylic substitution is proposed to occur by migratory insertion of the alkene followed by β-oxygen elimination: a) H. Ohmiya, Y. Makida, T. Tanaka, M. Sawamura, *J. Am. Chem. Soc.* 2008, *130*, 17276-17277. b) H. Ohmiya, Y. Makida, D. Li, M. Tanabe, M. Sawamura, *J. Am. Chem. Soc* 2010, *132*, 879-889. c) D. Li, T. Tanaka, H. Ohmiya, M. Sawamura, *Org Lett.* 2010, *12*, 3344-3347.
- [18] a) B. M. Trost, M. D. Spagnol, J. Chem. Soc., Perkin Trans 1 1995, 2083-2097. b) Y. Kobayashi, R. Mizojiri, E. Ikeda, J. Org. Chem. 1996, 61, 5391-5399. c) Y. Kobayashi, K. Watatani, Y. Kikori, R. Mizojiri, Tetrahedron Lett. 1996, 37, 6125-6128.

WILEY-VCH

COMMUNICATION

COMMUNICATION



Enantioselective nickel-catalyzed arylative cyclizations of substrates containing a *Z*-allylic phosphate tethered to an alkyne are described. These reactions give multisubstituted chiral aza- and carbocycles, and are initiated by the addition of an arylboronic acid to the alkyne, followed by cyclization of the resulting alkenylnickel species onto the allylic phosphate. The reversible E/Z isomerization of the alkenylnickel species is essential for the success of the reactions.

C. Yap, G. M. J. Lenagh-Snow, S. N. Karad, W. Lewis, L. J. Diorazio, H. W. Lam*

Page No. – Page No.

Enantioselective Nickel-Catalyzed Intramolecular Allylic Alkenylations Enabled by Reversible Alkenylnickel *E/Z* Isomerization