

Tetrahedron Letters 40 (1999) 4059-4062

Copper Salt - Crown Ether Systems as Catalysts for the Oxidation of Cumene with 1-Methyl-1-phenylethylhydroperoxide to bis(1-Methyl-1-phenylethyl)peroxide

Jan Zawadiak*, Danuta Gilner and Roman Mazurkiewicz

Institute of Organic Chemistry and Technology, Silesian Technical University, 44-100 Gliwice, ul. Krzywoustego 4, Poland

Received 8 February 1999; accepted 29 March 1999

Abstract: Evidence for phase transfer catalysis in the oxidation of cumene with 1-methyl-1-phenylethylhydroperoxide to bis-(1-methyl-1-phenylethyl)peroxide in the presence of copper salt – crown ether catalysts is given. © 1999 Elsevier Science Ltd. All rights reserved.

Organic peroxides are important components in the polymer industry. They can be obtained from catalytic reactions of hydrocarbons with an active hydrogen and hydroperoxides. Salts of Cu(1), Cu(11), Co(11), $Mn(11)^{1,2,3,4,5}$ are usually used as catalysts. The reaction proceeds according to the following equation (Scheme 1):

 $2ROOH + R'H \xrightarrow{cataliss} ROOR' + ROH + H_2O$ Scheme 1

It is known from our previous studies, that the synthesis of ditertiary peroxides is much more efficient in the presence of a catalytic system composed of a variable valence metal salt and an alkylammonium or alkylphosphonium salt, and can be carried out in moderate conditions, without an additional solvent.^{6,7}

Feldberg and Sasson⁸ have also applied a catalytic system consisting of copper(II) chloride and tetrabutylammonium bromide in the oxidation of tetralin to 1-(*tert*-butylperoxy)tetralin by tert-butylhydroperoxide (as a 70% solution in water). The process was carried out in methylene chloride, using a two- phase aqueous/organic system.

Barak and Sasson⁹ stated that quaternary ammonium salts may play a dual role in the metal-assisted oxidation of styrene to benzaldehyde or acetophenone by hydrogen peroxide. It extracts both metal and hydrogen peroxide into organic solvents of low polarity. It also modifies the nature of the metal by complexation, resulting in an improved catalytic activity and higher resistance of the metal towards reduction. An extraction mechanism involving the formation of a lipophilic ion pair, soluble in an organic solvent, is postulated for many types of reactions occurring in the presence of catalysts consisting of metal salts and onium salts.¹⁰

Our studies on the reaction of 1-methyl-1-phenylethyl hydroperoxide (1) with cumene (2) in the presence of catalysts composed of Cu(II), Cu(I), Co(II), Mn(II) salts and alkylammonium or alkylphosphonium salts¹¹ also indicate such a kind of catalysis (transfer of the catalyst in the form of a complex or ion pair with onium salt to the organic phase).

In this paper we present some experimental evidence for phase transfer catalysis in the oxidation of cumene (2) with 1-methyl-1-phenylethyl hydroperoxide (1) to bis(1-methyl-1-phenylethyl)peroxide (3) in the presence of copper salts and crown ethers (Scheme 2).



12-Crown-4, 15-crown-5, 18-crown-6 as well as Kryptofix 21 and Kryptofix 22 were applied with $CuCl_2 \cdot 2H_2O$ or CuCl. The inner diameters of 12-crown-4 (1,1÷1,4Å) and 15-crown-5 or Kryptofix 21 (1,7÷2,2Å) are adequate for the $Cu^{2\oplus}$ (1,38Å) and Cu^{\oplus} (1,92Å) ionic diameters. The process was carried out without any additional solvent and the aqueous phase was formed during the reaction (water is one of the products of this reaction).

In a typical procedure 0.23mol of (1) (as a 70% solution in cumene) was added with a dosage rate of $0.27 \text{ cm}^3/\text{min}$ to 0.14 mol of (2) with $4.1 \cdot 10^{-3}$ mol of CuCl₂·2H₂O (or CuCl), $8.4 \cdot 10^{-5}$ mol of crown ether (or tetrabutylammonium bromide) and in some experiments $4.1 \cdot 10^{-3}$ mol of LiCl (or NaCl, or KCl). The reaction was carried out at 70°C for 3h. The reaction rate as well as the composition of a post-reaction mixture were analysed using the iodometric method¹² and a HPLC method.¹³ The results are presented in Table 1.

We have found that the best results in the reaction of (1) with (2) were obtained in the presence of catalysts composed of $CuCl_2 \cdot 2H_2O$ and crown ethers with inner diameters larger than the ionic diameters of copper ions (15-crown-5 as well as Kryptofix 22). Moreover, we have also found that the presence of both components of the catalyst, i.e. copper salt and crown ether, is essential. No reaction was detected (under the applied conditions) in the absence of copper salt; on the other hand, only a 4% conversion of CHP was observed without crown ether.

The efficiency of the $CuCl_2 \cdot 2H_2O$ - crown ether catalytic system was enhanced markedly by adding some complexing salts like KCl, NaCl, LiCl, with the cation diameter corresponding to the inner diameter of the crown ether. The most striking results were obtained adding LiCl to a hardly active catalyst composed of $CuCl_2 \cdot 2H_2O$ and 12-crown-4. In the presence of LiCl both the reaction rate and the yield of peroxide increased considerably. The possibility of a catalytic activity of the 12-crown-4/LiCl and 18-crown-6/KCl systems was excluded by demonstrating that no oxidation takes place without $CuCl_2 \cdot 2H_2O$.

Copper salt	Crown ether	Complexing salt Conversion of 1 [%]		Yield of 3 [%]
CuCl ₂ ·2H ₂ O	_	-	4	1
-	18-crown-6	-	0	0
CuCl ₂ ·2H ₂ O	12-crown-4	-	9	8
CuCl ₂ ·2H ₂ O	12-crown-4	LiCl	84	46
-	12-crown-4	LiCl	0	0
CuCl ₂ ·2H ₂ O	15-crown-5	-	82	42
CuCl ₂ ·2H ₂ O	15-crown-5	NaCl	82	48
CuCl ₂ ·2H ₂ O	18-crown-6	-	72	33
CuCl ₂ ·2H ₂ O	18-crown-6	KCl	88	46
-	18-crown-6	KCl	0	0
CuCl ₂ ·2H ₂ O	Kryptofix22	-	80	45
$CuCl_2 \cdot 2H_2O$	Kryptofix21	-	82	36
CuCl	15-crown-5	-	6	4
CuCl ₂ 2H ₂ O	TBAB	-	83	63

Table 1. The influence of crown ethers on the oxidation of cumene with 1-methyl-1-phenylethylhydroperoxide to bis(1-methyl-1-phenylethyl)peroxide in the presence of CuCl₂·2H₂O

lonic diameters: Cu[#] 1,92Å; Cu^{2#} 1,38Å; Li[#] 1,20Å; Na[#] 1,90Å; K[#] 2,66Å; Inner diameter of crown ethers: 12-crown-4 1,1+1,4Å; 15-crown-5 1,7+2,2Å; 18-crown-6 2,6+3,2Å; Kryptofix 21 1,7+2,2Å; Kryptofix 22 2,6+3,2Å

The results obtained confirm strongly that phase transfer catalysis operates in the reaction of (1) with (2). In general, phase transfer catalysis consists in the transfer of copper ions to the organic layer. The copper ions can be transferred in the form of complexes of a copper cation with a crown ether. Our results prove that copper cations may be transferred to the organic layer also as complexes of Li^{\oplus} , Na^{\oplus} or K^{\oplus} cations with the crown ether and copper bonded in $CuCl_4{}^{2\Theta}$, $CuCl_3{}^{\Theta}$ or $CuCl_2{}^{\Theta}$ counterions. Copper ions can also be transferred in the form of complexes $Cu[CuCl_3]$, $Cu[CuCl_4]$. The complexes discussed are known from literature.^{14, 15}

The compositions of the post-reaction mixtures in the cases of applying 15-crown-5, 15-crown-5+NaCl and TBAB in the reactions of (1) with (2) in the presence of $CuCl_2 \cdot 2H_2O$ were compared (Table 2). If the 15-crown-5 + $CuCl_2 \cdot 2H_2O$ system is applied, nearly twice the amount of alcohol is formed (compared with that formed in the presence of TBAB + $CuCl_2 \cdot 2H_2O$). It results from intensive activity of the catalyst in the decomposition of (1) to by-products (particularly to (4)). The addition of NaCl to 15-crown-5 + $CuCl_2 \cdot 2H_2O$

does not affect any change in the composition of the product. Investigations have proved that in the presence of crown ether 15-crown-5, at 70°C, after 5 hours peroxide is not decomposed.

Table 2. Composition of the reaction mixture after the oxidation of cumene with 1-methyl-1phenylethylhydroperoxide in the presence of catalytic systems CuCl₂·2H₂O+15-crown-5 or CuCl₂·2H₂O+15-crown-5+NaCl or CuCl₂·2H₂O+TBAB

Catalyst	1[mol]	3 [mol]	4 [mol] 0,20	PhAc* [mol] 0,04
15-crown-5+CuCl ₂ ·2H ₂ O	0,04	0,05		
15-crown-5 +CuCl ₂ ·2H ₂ O +NaCl	0,04	0,06	0,21	0,04
TBAB+CuCl ₂ ·2H ₂ O	0,02	0,07	0,13	0,05
*Dh A a= A cetonhonone				

*PhAc=Acetophenone

References:

- 1. Kharasch, M.S.; Fono, A.; J. Org. Chem. 1959, 24, 72.
- 2. Antonovskii, V.L.; Emelin, Y.D.; Zh. V. Khim. Obshch. 1964, 9(4), 471.
- 3. Kulicki, Z.; Zawadiak, J.; PL Patent 1980, 105 264.
- 4. Kulicki, Z.; Baj, S.; Zawadiak, J.; Pol. J. Chem., 1980, 54, 2337.
- 5. Komai, T.; Sawada, H.; Kato, M.; Jap. Patent, 1989, 1 019 062.
- Zawadiak, J.; Stec, Z.; Kulicki, Z.; Burghardt, A.; Staniowski, B.; Jóźwicki, R.; Stolarczyk, G.; PL. Patent 1988, 156 813.
- 7. Zawadiak, J.; Gilner, D.; Pol.J.Appl.Chem. 1995, 39(2), 225.
- 8. Feldberg, L.; Sasson, Y.; Tetrahedron Lett. 1996, 37(12), 2063.
- 9. Barak, G.; Sasson, Y.; J. Chem. Soc., Chem. Commun. 1987, 1266.
- Sasson, Y.; Neumann, R.; Handbook of Phase Transfer Catalysis; Blackie Academic and Professional; London, 1997; pp. 317; 524-525.
- 11. Zawadiak, J.; Zeszyty Naukowe Politechniki Śląskiej, s. Chemia, 1991, 126.
- 12. Zawadiak, J.; Gilner, D.; Kulicki, Z.; Baj, S.; Analyst 1993, 118 (8), 1081.
- 13. Baj, S.; Kulicki, Z.; J. Chrom., 1991, 588, 33.
- 14. Gmelins Handbuch der Anorganischen Chemie; 8 Auflage, Teil **B 60**, Liferung 1, Verlag Chemie GmbH, Weinheim, Germany, 1958, pp. 251-321.
- Ullmann's Encyclopedia of Industrial Chemistry, V Ed., Vol. A 7, VCH Verlagsgesellschaft mBH, Weinheim, Germany, 1991; H.Wayne Richardson, Copper Compounds, pp. 567-587.