



Pergamon

Tetrahedron Letters 40 (1999) 1137-1140

---

---

TETRAHEDRON  
LETTERS

---

---

# Enantioselective $\alpha$ -Hydroperoxylation of Long-Chain Fatty Acids with Crude Enzyme of Marine Green Alga *Ulva pertusa*

Yoshihiko Akakabe\*, Kenji Matsui and Tadahiko Kajiwara

Department of Biological Chemistry, Faculty of Agriculture, Yamaguchi University, 1677-1 Yoshida, Yamaguchi 753-8515, Japan

Received 28 September 1998; accepted 27 November 1998

**Abstract:** When palmitic acid was incubated with crude enzyme of marine green alga *Ulva pertusa*, (*R*)-2-hydroperoxyhexadecanoic acid was formed in high enantiomeric purity (>99%ee).

© 1999 Elsevier Science Ltd. All rights reserved.

Long-chain fatty acids can be metabolised to the corresponding aldehydes by  $\alpha$ -oxidation in a wide range of organisms<sup>1-6</sup>. Although  $\alpha$ -oxidation of fatty acids appears to be important for the physiological functions, the reaction pathway has not been fully characterised<sup>7,8</sup>.

In an essential oil, which prepared from marine green alga *Ulva pertusa* by simultaneous distillation extraction (SDE) manipulation, long-chain fatty aldehydes such as pentadecanal, (8*Z*)-heptadecenal, (8*Z*, 11*Z*)-heptadecadienal and (8*Z*, 11*Z*, 14*Z*)-heptadecatrienal, were identified by comparison of GC-MS data with those of authentic samples<sup>9</sup>. As expected, long-chain fatty acids; palmitic acid, oleic acid, linoleic acid, and linolenic acid, were converted to the corresponding aldehydes with a loss of one carbon atom<sup>10,11</sup>. Although a partly purification of long-chain aldehyde forming enzyme was examined, the enzyme has not been sufficiently characterised<sup>12,13</sup>. Thus, to elucidate the mechanism and the physiological functions of  $\alpha$ -oxidation in *U. pertusa*, the biogenetic intermediate has to be determined. In this paper, we describe highly enantioselective  $\alpha$ -hydroperoxylation of long-chain fatty acids with crude enzyme of *U. pertusa*.

The alga *U. pertusa* was collected in the intertidal zone of Aio, Yamaguchi, Japan on 31 July 1997 and immediately frozen at -20°C. The frozen tissue (100g fresh weight) was cut into small pieces and homogenized with 0.1M phosphate buffer (250ml, pH 6.0), which contained 0.1% Triton X-100, in a blender for 10 min. After filtering, the resulting homogenate was centrifuged at 150000 x *g* for 60 min. The pellet was resuspended in the same buffer (50ml) containing 0.1% Triton X-100. The crude enzyme solution was stirred at 5°C for 10 min, and then

---

E-mail : akakabe@agr.yamaguchi-u.ac.jp




Figure 1 Comparison of 2-oxygenated compounds from extracts of *U. pertusa* with synthetic samples. Condition : column Mightysil RP-18 GP ; eluent CH<sub>3</sub>CN / 0.1M CH<sub>3</sub>CO<sub>2</sub>NH<sub>4</sub> (9 / 1) ; flow rate 1.0ml / min ; detector fluorescence Ex 365nm, Em 412nm. A : synthetic samples, I 2-hydroperoxyhexadecanoic acid ; II 2-hydroxyhexadecanoic acid ; III 2-oxohexadecanoic acid ; IV pentadecanoic acid ; V palmitic acid. B : The crude enzyme of *U. pertusa* was stirred with palmitic acid. C : The crude enzyme was incubated in the absence of substrate. D : Heat treatment of the crude enzyme was carried out at 90°C prior to addition of substrate.

palmitic acid (5mg, 0.02mmol) in DMSO (0.5ml) was administered. After stirring at 5°C for 30 min, (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>, NaCl, and THF were added to the reaction mixture. The whole mixture was centrifuged at 2000 x g for 10 min, the organic layer was separated. The layer was washed with sat. NaCl solution and dried over MgSO<sub>4</sub>. A portion of the extract was treated with 9-anthryldiazomethane (ADAM) at 0°C for 10 min<sup>14</sup>. Then the reaction mixture was concentrated *in vacuo* and the residue was subjected to HPLC analysis of oxygenated products<sup>15</sup>. With palmitic acid, two major peaks of ADAM esters appeared as shown in Figure 1-B, i.e. the ADAM esters of 2-hydroperoxyhexadecanoic acid (I ; Rt 21.0 min) and palmitic acid (V; Rt 64.0 min)<sup>16</sup>. These compounds were identified by comparison with the ADAM ester of synthetic samples<sup>17</sup>. Indeed, when a portion of the extract was treated with PPh<sub>3</sub> prior to HPLC analysis, peak I of 2-hydroperoxy form disappeared whereas peak II of 2-hydroxy form increased (Figure 2). In the separate experiment, the extract was esterified with diazomethane at 0°C for 10 min, and the resulting ester was purified by preparative TLC to give methyl ester of 2-hydroperoxyhexadecanoic acid, which exhibited identical properties (TLC<sup>18</sup> and LC-MS<sup>19</sup>) with those of a synthetic sample. Thus, the HPLC and LC-MS data revealed that the addition of palmitic acid to the crude enzyme solution led to an




Figure 2 Confirmation of compounds involving 2-hydroperoxy group with PPh<sub>3</sub>. For conditions, see Figure 1 legend.




Figure 3 HPLC enantioseparation of 2-hydroperoxyhexadecanoic acid. condition : column Develosil ODS-5 ; eluent CH<sub>3</sub>CN ; flow rate 1.0ml/min ; detector fluorescence Ex 365nm, Em412nm. A : separation of (±)-2-hydroperoxyhexadecanoic acid. B : enantiopurity analysis of 2-hydroperoxyhexadecanoic acid obtained from homogenate of *U. pertusa*.


increased yield of the 2-hydroperoxy acid. In turn, incubation of heat-treated crude enzyme prior to addition of palmitic acid did not produce the 2-hydroperoxy acid (Figure 1-D), thus indicating that the 2-oxygenated compound I is formed enzymatically .

On the other hand, a solution of the ADAM ester I in THF was treated with Noe's reagent<sup>20</sup> and TsOH (catalytic amount) at room temperature for 5 min. The reaction mixture was concentrated *in vacuo*, and the concentrate, without further purification, was circulated to determine the absolute configuration and enantiomeric purity by HPLC. On the basis of the HPLC elution pattern

Table 1 Determination of optical purity of 2-hydroperoxy acids using Noe's reagent

fatty acids	2-hydroperoxy acids	
	%ee	confign
myristic acid (C14)	>99	<i>R</i>
pentadecanoic acid (C15)	>99	<i>R</i>
palmitic acid (C16)	>99	<i>R</i>
heptadecanoic acid (C17)		n.d.*
stearic acid (C18)		n.d.

\* not detected



Scheme 1 Highly enantioselective  $\alpha$ -hydroperoxylation of long-chain fatty acids with crude enzyme of *U. pertusa*.

in Figure 3, the peracetal of I from the crude enzyme was attributed to (*R*)-enantiomer of synthetic 2-hydroperoxyhexadecanoic acid and the enantiomeric excess of the product was shown to be >99%<sup>21</sup>.

The enantiospecificity of the hydroperoxylation using several other long-chain fatty acids was investigated for the crude enzyme (Table 1). With pentadecanoic and myristic acid, (*R*)-2-hydroperoxypentadecanoic and (*R*)-2-hydroperoxytetradecanoic acid were obtained with an enantiomeric excess of >99%, respectively. However, heptadecanoic and stearic acid were poor substrates.

In conclusion, our results show that long-chain fatty acids (C14-C16) are  $\alpha$ -hydroperoxylated with the crude enzyme of *U. pertusa* to afford (*R*)-2-hydroperoxy acids with excellent enantiomeric excess (Scheme 1).

**Acknowledgment.** We thank Dr Hironori Masui of Ymaguchi Women's University for LC-MS measurements.

### References and Note

- Hitchcock, C. and James, A. T., *Biochim. Biophys. Acta* **1966**, 116, 413-424.
- Laties, G. G. and Hoelle, C., *Phytochemistry* **1967**, 6, 49-57.
- Shine, W. E. and Stumpf, P. K., *Arch. Biochem. Biophys.* **1974**, 162, 147-157.
- Galliard, T. and Matthew, J. A., *Biochim. Biophys. Acta* **1976**, 424, 26-35.
- Hanna, M. -S., Campa, A. and Cilento, G., *Photochem. Photobiol.* **1987**, 45, 849-854.
- Adam, W., Lazarus, M., Möler, C. R. -S. and Schreier, P., *Tetrahedron : Asymmetry* **1996**, 7, 2287-2292.
- Borge, G. I. A., Slinde, E. and Nilsson, A., *Biochim. Biophys. Acta* **1997**, 1344, 47-58.
- Matsunaga, I., Yamada, M., Kusunose, E., Miki, T. and Ishihara, K., *J. Biochemistry* **1998**, 124, 105-110.
- Fujimura, T., Kawai, T., Shiga, M., Kajiwar, T. and Hatanaka, A., *Phytochemistry* **1990**, 29, 745-747.
- Kajiwar, T., Yoshikawa, H., Saruwatari, T., Hatanaka, A., Kawai, T., Ishihara, M. and Tsuneya, T. *Phytochemistry* **1988**, 27, 1643-1645.
- Kajiwar, T., Kashibe, M., Matsui, K. and Hatanaka, A., *Phytochemistry* **1991**, 30, 193-195.
- Kajiwar, T., Hatanaka, A., Matsui, K., Tomoi, T. and Idohara, T., *Phytochemistry* **1994**, 35, 55-57.
- Kajiwar, T., Matsui, K. and Akakabe, Y., In "Biotechnology for Improved Foods and Flavors (ACS Symposium Series No. 637, edited by G. R. Takeoka, R. Teranishi, P. J. Williams, and A. Kobayashi, American Chemical Society)" **1996**, 146-166.
- Nimura, N. and Kinoshita, T. *Anal. Lett.* **1980**, 13, 191.
- A Mightysil RP-18 GP column (3.0 x 250mm) was used for analysis of oxygenated compounds formed by the  $\alpha$ -oxidation, while a Develosil ODS-5 column (4.6 x 250mm) was used to evaluate the absolute configuration of 2-hydroperoxy and 2-hydroxyhexadecanoic acid produced from palmitic acid.
- In addition, minor peaks were observed and retention times were follows : 24.5 min, 2-hydroxyhexadecanoic acid II ; 34.0 min, 2-oxohexadecanoic acid III ; 46.5 min, pentadecanoic acid IV.
- Incidentally, racemic 2-hydroperoxyhexadecanoic acid can be conveniently prepared from palmitic acid through sequential formation of dianion with lithium diisopropylamide (LDA) and O<sub>2</sub> bubbling process at -78°C : Konen, D. A., Silbert, L. S. and Pfeffer, P.E. *J. Org. Chem.* **1975**, 40, 3253-3258. Of course, the 2-hydroxy acid can be also obtained by reduction of the 2-hydroperoxy acid with PPh<sub>3</sub>.
- Confirmation of compounds involving hydroperoxy acids was monitored by TLC, using an *N, N'*-dimethyl-*p*-phenylenediamine indicator.
- LC-MS of the purified compound gave a pseudomolecular ion at *m/z* 320 [(M+NH<sub>4</sub>)<sup>+</sup>] in accordance with the synthetic sample.
- Noe, C. R., *Chem. Ber.* **1982**, 115, 1591-1606 ; Noe, C. R., Knollmüller, M., Oberhauser, B., Steinbauer, G. and Wagner, E., *Chem. Ber.* **1986**, 119, 729-743 ; Knollmüller, M., Noe, C. R. and Oberhauser, B., *Monatsh. Chem.* **1986**, 117, 407-419.
- Furthermore, reduction of the 2-hydroperoxy acid of I from the crude enzyme with PPh<sub>3</sub>, followed by treatment with Noe's reagent, gave the corresponding acetal, which coincided with the (*R*)-form of synthetic 2-hydroxy acid : Mori, K. and Funaki, Y., *Tetrahedron* **1985**, 41, 2379-2386 ; Sugai, T. and Ohta, H., *Agric. Biol. Chem.* **1990**, 54, 3337-3338.