

# A Journal of the Gesellschaft Deutscher Chemiker A Deutscher Chemiker GDCh International Edition www.angewandte.org

## **Accepted Article**

Title: Photoredox/Nickel-Catalyzed Single-Electron Tsuji-Trost Reaction: Development and Mechanistic Insight

Authors: Gary Molander, Jennifer Matsui, Alvaro- Gutierrez-Bonet, Madeline Rotella, Rauful Alam, and Osvaldo Gutierrez

This manuscript has been accepted after peer review and appears as an Accepted Article online prior to editing, proofing, and formal publication of the final Version of Record (VoR). This work is currently citable by using the Digital Object Identifier (DOI) given below. The VoR will be published online in Early View as soon as possible and may be different to this Accepted Article as a result of editing. Readers should obtain the VoR from the journal website shown below when it is published to ensure accuracy of information. The authors are responsible for the content of this Accepted Article.

To be cited as: Angew. Chem. Int. Ed. 10.1002/anie.201809919 Angew. Chem. 10.1002/ange.201809919

Link to VoR: http://dx.doi.org/10.1002/anie.201809919 http://dx.doi.org/10.1002/ange.201809919

## WILEY-VCH

COMMUNICATION

#### WILEY-VCH

## Photoredox/Nickel-Catalyzed Single-Electron Tsuji-Trost Reaction: Development and Mechanistic Insight

Jennifer K. Matsui,<sup>†</sup> Álvaro Gutiérrez-Bonet,<sup>†</sup> Madeline Rotella,<sup>‡</sup> Rauful Alam,<sup>†</sup> Osvaldo Gutierrez<sup>‡\*</sup> and Gary A. Molander<sup>†\*</sup>

Abstract: A regioselective, nickel-catalyzed photoredox allylation of secondary, benzyl, and  $\alpha$ -alkoxy radical precursors is disclosed. Through this manifold, a variety of linear allylic alcohols and allylated monosaccharides are accessible in high yields under mild reaction conditions. Quantum mechanical calculations [DFT and DLPNO-CCSD(T)] support the mechanistic hypothesis of a Ni(0) to Ni(II) oxidative addition pathway followed by radical addition and inner-sphere allylation.

Over the course of the past 30 years, the Tsuji-Trost allylation has been a workhorse reaction within the field of synthetic organic chemistry.<sup>1</sup> A particularly striking advantage of the Tsuji-Trost reaction is the ability to forge C(sp<sup>3</sup>)–C(sp<sup>3</sup>) bonds under mild reaction conditions, typically using palladium as the transition metal in conjunction with an appropriate phosphine ligand.<sup>2</sup> Although the breadth of electrophilic allyl partners has been quite thoroughly explored, nucleophilic partners have primarily been limited to two-electron carbon- or nitrogencentered moieties (Scheme 1).<sup>3</sup> Specifically, for C–C bond formation, "soft" nucleophiles (e.g., malonates) are frequently employed, wherein addition occurs by an outer-sphere mechanism.<sup>1</sup>

The choice of nucleophile in these transformations is primarily limited by pKa constraints, because anions derived from pronucleophiles with pKa >25 are not appropriately tuned to react with the incipient  $\pi$ -allyl transition metal intermediate.<sup>4</sup> Recent reports by Walsh and coworkers have pointed to the utility of diarylmethane nucleophiles (pKa up to 32) in Pdcatalyzed allylic substitution reactions, broadening the nucleophile window of Tsuji-Trost reactions.<sup>5</sup> Alternatively, Zn,<sup>6</sup> Sn, B, Mg, Si, and Li-based enolates have been used successfully as nucleophilic partners related in transformations.<sup>2b,12</sup> However, a major limitation of enolatebased methods is the prerequisite for highly basic conditions. To overcome these limitations, pioneering work by Tunge shifted efforts toward radical-based intramolecular allylations via a

[a] Dr. J. K. Matsui, Dr. A. Gutiérrez-Bonet, Dr. R. Alam, Prof. G. A. Molander Department of Chemistry University of Pennsylvania Roy and Diana Vagelos Laboratories, Philadelphia, Pennsylvania 19104-6323. United States of America E-mail: <u>gmolandr@sas.upenn.edu</u>
[b] M. Rotella, Prof. O. Gutierrez Department of Chemistry and Biochemistry University of Maryland College Park, Maryland

20742. United States of America

Email: <u>ogs@umd.edu</u> Supporting information for this article is given via a link at the end of the document. palladium/photoredox cross-coupling, facilitating the decarboxylative allylation of  $\alpha$ -amino carboxylic acids. Unfortunately, when Tunge and coworkers shifted to *intermolecular* allylation chemistry, only unsubstituted allyl groups were successfully coupled with alkyl carboxylic acid partners.<sup>10</sup>

Given the excellent precedent for merging photoredox and nickel catalysis in alkylation reactions established by our laboratory and others," we sought to expand this paradigm to radical-based Tsuji-Trost reactions (Scheme 1). In particular, numerous radical (e.g., precursors alkyltrifluoroborates, carboxylates, alkylsilicates, and 1,4-dihydropyridines) derived from diverse commodity chemicals have been developed.<sup>15</sup> Although such reagents have been extensively utilized in conjunction with photoredox/Ni dual cross-coupling with aryl-16 and alkenyl17 electrophiles, we have also begun to expand the scope of electrophilic partners for such transformations (e.g., using isocyanate, 18 acyl chloride, 19 carboxylic acid, 20 and acyl imide electrophiles<sup>21</sup>). In a continuation of that effort, the cost-effective nature of Ni-based catalysts and the slow  $\beta$ -hydride elimination of alkylnickel species relative to that of palladium was considered to investigate how this approach could be applied to the Tsuji-Trost reaction.22



Scheme 1. Exploring radical-based Tsuji-Trost reactivity under photoredox conditions.

Vinyl epoxides have been employed previously in palladiumcatalyzed allylations,<sup>33</sup> but to the best of our knowledge, reports using a nickel-based catalyst — a non-precious, inexpensive catalyst — are scarce,<sup>34</sup> with only a few examples being reported involving cycloaddition reactions. At the outset of substrate exploration, a variety of benzyltrifluoroborates were selected; early results demonstrated a diversity of electron-withdrawing and electron-donating groups were accommodated (Table 1). Additionally, modestly sterically hindered substrates could also be used. For example, ortho-substitution afforded 62% of desired product **1f**. As the radical precursor search was expanded,  $\alpha$ alkoxytrifluoroborates were next targeted. Generally, *E/Z* selectivity was high, although the ratio was severely diminished to 53:47 when *tert*-butyl ether moieties were explored (**1j**).

Moving forward, the breadth of carbon-centered radicals was expanded by targeting unstabilized secondary alkyl radical

#### COMMUNICATION

precursors. Under the standard and modified reaction conditions, secondary alkyltrifluoroborates formed complex mixtures (e.g., homocoupling, aldehydic side products).<sup>25</sup> Therefore, DHPs (1,4-dihydropyridines) were investigated as alternative radical precursors <sup>26</sup> <sup>- 27</sup> with 4CzIPN [2,4,5,6-tetra(9H-carbazol-9-yl)isophthalonitrile] organophotocatalyst.





**Table 1.** Reactions were conducted on 0.50 mmol scale. Standard reaction conditions employed butadiene monoxide (1.0 equiv), alkyltrifluoroborate (1.5 equiv), Ir[dFCF<sub>3</sub>ppy]<sub>2</sub>(bpy)PF<sub>6</sub> (2 mol %), [Ni(dtbbpy)(H<sub>2</sub>O)<sub>4</sub>]Cl<sub>2</sub> (3 mol %), and 2,6-lutidine (3.5 equiv) in acetone/MeOH (9:1). <sup>a</sup>For α-alkoxyalkyltrifluoroborates, a solvent mixture of acetonitrile/*tert*-butanol (10:1) was used.

During the exploration of various DHPs, heterocyclic (**2a** and **2e**) and alkenyl moieties (**2c**) were successfully incorporated within the coupling partners under the optimized reaction conditions. A desire to integrate greater functional density within the radical precursor led to the exploration of monosaccharide moieties that were recently highlighted in publications from our group. <sup>as</sup> Pleasingly, pyranose DHP afforded the allylated products **2i-j** in high E/Z ratios and diastereoselectivity.

Subsequently, substitution on the vinyl epoxide moiety was probed, starting with methyl substitution on the requisite epoxide (**2f-2h**). Notably, the stereochemical outcome of the photoredox/Ni dual catalyzed alkylation is complementary to that of the Pd-catalyzed transformations. The latter provide net retention via double inversion,<sup>26</sup> whereas in the present case 1,2-epoxy-3,4-cyclohexene afforded primarily the trans isomer (**2m**), suggesting the nickel coordinates the alkene and adds to the opposite face of the epoxide, but then the resulting nickel complex engages the radical and undergoes reductive elimination by a stereoretentive, inner-sphere mechanism. This hypothesis was further supported by both DFT and DLPNO-CCSD(T) quantum mechanical calculations (vide infra).

**Table 2.** Reactions were conducted on 0.50 mmol scale. Standard reaction conditions employed butadiene monoxide (1.0 equiv), 1,4-dihydropyridine (1.5 equiv), 4CzIPN (2 mol %), and [Ni(dtbbpy)(H<sub>2</sub>O)<sub>4</sub>]Cl<sub>2</sub> (3 mol %) in acetone. For trifluoroborates, standard reaction conditions employed butadiene monoxide (1.0 equiv), alkyltrifluoroborate (1.5 equiv), Ir[dFCF<sub>3</sub>ppy]<sub>2</sub>(bpy)PF<sub>6</sub> (2 mol %), [Ni(dtbbpy)(H<sub>2</sub>O)<sub>4</sub>]Cl<sub>2</sub> (3 mol %), and 2,6-lutidine (3.5 equiv) in acetone/MeOH (9:1). For  $\alpha$ -alkoxyalkyltrifluoroborates, a solvent mixture of acetonitrile/tert-butanol (10:1) was used.

Finally, although the allylic alcohol products themselves are versatile moieties,<sup>39</sup> we aimed to make the protocol more general by exploring electrophiles as  $\pi$ -allylnickel precursors. First, various leaving groups were surveyed (i.e., bromide, chloride,

### COMMUNICATION

and carbonate), with allyl bromide affording the highest yield of product under the initial reaction conditions.



Table 3. Expanding the allylic handle.

Motivated by the advantageous biological properties demonstrated by molecules containing alkyl-substituted carbohydrate substructures, " DHP monosaccharides were successfully allylated under the dual catalytic conditions with modest to high diastereoselectivities (Table 4). Pyranose and furanose backbones with benzyl-, methoxy-, and dioxolane protecting groups were conserved. Further studies on substituent effects were conducted using difluoroallyl bromide. Under the established reaction conditions, **4c** was isolated as a single regioisomer. Presumably, oxidative addition to allyl bromide followed by bromide expulsion generates the key  $\pi$ -allylnickel complex which, via an inner-sphere mechanism (vide infra), intercepts the  $\alpha$ -alkoxy radical to undergo diastereoselective reductive elimination, forming the allylated products.

Intrigued by the stereoselectivity of these processes and questions regarding the order of reaction events, quantum mechanical calculations were undertaken to gain insight into



**Table 4.** All reactions were performed using 0.75 mmol of allyl bromide, 0.25mmol of 1,4-dihydropyridine, 4CzIPN (2 mol %), [Ni(dtbbpy)(H<sub>2</sub>O)<sub>4</sub>]Cl<sub>2</sub> (3 mol %) in acetone with blue LED irradiation for 24 h.

the underlying mechanism of this transformation.<sup>31</sup> As shown in Figure 1 (squares), complexation of Ni(0) to vinyl epoxide (A") is energetically favored over that of complexation to benzyl radical (A'; presumably formed from the photocatalytic cycle, see Supporting Information for energetics) by 8.9 kcal/mol. Subsequently, this vinyl epoxide-Ni(0) (A'') complex can quickly undergo an S<sub>N</sub>2-like ring-opening (via A"-TS; barrier is only 3.1 kcal/mol from  $A^{**}$  to form  $\pi$ -allylnickel(II) intermediate **B**<sup>\*\*</sup>. As anticipated by previous calculations,<sup>32</sup> the nickel(II) intermediate B" can quickly engage the benzyl radical, generated by SET oxidation of BnBF3K by the photocatalyst (see Supporting Information for energetics), to form Ni(III) intermediate C (via B"-C-TS). Overall, radical addition is facile (the barrier is ca. 1-3 kcal/mol, dependent on the method). Finally, inner-sphere  $C(sp^2)$ - $C(sp^3)$  bond formation via C-TS<sub>L</sub> (the barrier is 8.9 kcal/mol from C) leads to formation of the linear product (P<sub>L</sub>), 50.3 kcal/mol downhill in energy.



Figure 1. Competing Ni(0)/Ni(1)/Ni(11) (circles), Ni(0)/Ni(11)/Ni(111) (squares), Ni(0)/Ni(11) (squares-triangles), pathways leading to C(sp<sup>3</sup>)-C(sp<sup>3</sup>) bond formation

### COMMUNICATION

A series of reductive elimination transition states leading to formation of branched product (e.g., C-TS<sub>B</sub>) were also located, but these were much higher in energy (>5 kcal/mol) than  $C-TS_L$ , presumably because of unfavorable electronic repulsion and steric interactions between the charged oxygen and the benzyl group. Moreover, C(sp<sup>2</sup>)-C(sp<sup>3</sup>) bond formation via an outer sphere pathway (squares-triangles) was also found higher in energy. Formation of Ni(III) complex C from a Ni(0)-Ni(I)-Ni(III) pathway (Figure 1, circle) is also overall higher in energy than the Ni(0)-Ni(II)-Ni(III) pathway (squares). Albeit low, the barrier for radical addition (A-TS) from Ni(0) is still 7.3 kcal/mol higher in energy than oxidative addition to vinyl epoxide (via A"-TS). However, it is likely that formation of C is highly dependent on the local concentration of benzyl radical and vinyl epoxide. Overall, DFT and DLPNO-CCSD(T) calculations support a Ni(0)-Ni(II)-Ni(III) reaction pathway and an inner sphere reductive elimination as the product-determining step, leading to formation of the linear product as the major product.33 This model is in accord with the stereochemical results observed with cyclohexadiene monoxide (Table 2, 2m). Further, although computational studies favored the linear product, the branched pathway is only 5 kcal/mol higher in energy, suggesting branched products may be observed in some cases (likely dependent on the nature of the substrate, ligand, etc.). To test this hypothesis, we subjected sterically hindered arylsubstituted vinyl epoxides to the standard reaction conditions and indeed found a mixture of linear and branched products, albeit in low yields (Scheme 2).



Scheme 2. Observation of linear and branched mixtures

In summary, a dual catalytic allylation reaction is reported that demonstrates complementarity to traditional Tsuji-Trost reactions. In juxtaposition with classical reactivity in the Pd-catalyzed transformations (stabilized anionic carbon nucleophiles), carbon-centered radicals were employed, enabling the allylation of secondary, benzyl, and  $\alpha$ -alkoxy radical precursors. Additionally, net inversion of stereochemistry is observed in the photoredox/Ni dual catalyzed

systems, as opposed to the net retention via double inversion observed in the traditional Pd-catalyzed processes.

To gain insight into the mechanistic pathway, studies incorporating quantum mechanical calculations [using dispersion corrected (U)DFT and DLPNO-CCSD(T) methods] were conducted to determine the energetic favorability of the oxidative addition step. Here, a Ni(0) to Ni(II) oxidative addition step was calculated to occur before radical addition to the nickel center; this is a deviation from previous mechanistic studies of photoredox/Ni dual cross-coupling pathways where Ni(0) to Ni(I) has been involved in the putative pathway.<sup>32</sup> Given the mild reaction conditions and availability of numerous radical precursors, the disclosed method provides a useful complement to the traditional, palladium-catalyzed Tsuji-Trost reaction.

#### Acknowledgements

The authors are grateful for the financial support provided by NIGMS (R01 GM 113878 to G.M.) and NSF (CAREER, 1751568 to O.G.). J.K.M. is supported by the Bristol-Myers Squibb Graduate Fellowship for Synthetic Organic Chemistry. O.G. is grateful to the University of Maryland College Park for start-up funds and computational resources from UMD Deepthought2 and MARCC/BlueCrab HPC clusters and XSEDE (CHE160082 and CHE160053). We thank Dr. Charles W. Ross, III (University of Pennsylvania) for assistance in obtaining HRMS data. Additionally, Dr. Jun Gu (University of Pennsylvania) and Matt Bunner (University of Pennsylvania) assisted in the acquisition and analysis of NMR data. The alkyltrifluoroborates and the iridium (for photocatalyst synthesis) were generously donated by Frontier and Johnson-Matthey, respectively.

#### Abbreviations

DHP, 1,4-dihydropyridine; SET, Single Electron Transfer; 4CzIPN, 2,4,5,6-tetra(9H-carbazol-9-yl)isophthalonitrile.

#### **Conflict of interest**

The author declare no conflict of interest.

**Keywords:** Photoredox, Nickel Catalysis, 1,4-Dihydropyridines, Allylation, Tsuji-Trost

### COMMUNICATION

#### References

- <sup>1</sup> B. M. Trost, Acc. Chem. Res. **1980**, *13*, 385–393.
- <sup>2</sup> (a) J. Tsuji, Organic Synthesis with Palladium Compounds, Springer: New York, 1980. (b) B. M. Trost, D. L. Van Vranken, 1996, 96, 395–422.
- <sup>3</sup> (a) G. Consiglio, R. M. Waymouth, Chem. Rev. 1989, 89, 257–276. (b) A. Heumann, M. Reglier, Tetradedron 1995, 51, 975–1015.
- <sup>4</sup> B. M. Trost, M. R. Machacek, A. Aponick, Acc. Chem. Res. 2006, 39, 747.
- <sup>5</sup> S. -C. Sha, J. Zhang, P. J. Carroll, P J. Walsh, J. Am. Chem. Soc. 2013, 135, 17602–17609.
- <sup>6</sup> S. Son, G. C. Fu, J. Am. Chem. Soc. 2008, 130, 2756–2757.
- <sup>7</sup> (a) B. M. Trost, C. R. Self, J. Org. Chem. 1984, 49, 468–473. (b) B. M. Trost, E. Keinan, Tetrahedron Lett. 1980, 21, 2591–2594.
- <sup>8</sup> E. Negishi, H. Matsushita, S. Chatterjee, R. A. John, J. Org. Chem. 1982, 47, 3188–3190.
- <sup>9</sup> M. Braun, T. Meier, Angew. Chem. Int. Ed. 2006, 45, 6952–6955.
- <sup>10</sup> T. Graening, J. F. Hartwig, J. Am. Chem. Soc. 2005, 127, 17192–17193.
- <sup>11</sup> B. M. Trost, G. M. Schroeder, J. Am. Chem. Soc. 1999, 121, 6759-6760.
- <sup>12</sup> M. Braun, T. Meier, Angew. Chem. Int. Ed. 2006, 45, 6952–6955.
- <sup>13</sup> S. B. Lang, K. M. O'Nele, J. A. Tunge, J. Am. Chem. Soc. 2014, 136, 13606–13609.

<sup>14</sup> (a) J. C. Tellis, D. N. Primer, G. A. Molander, *Science* **2014**, *345*, 433–436. (b) Z. Zuo, D. T. Ahneman, L. Chu, J. A. Terett, A. G. Doyle, D. W. C. MacMillan, *Science* **2014**, *345*, 437–440.

<sup>15</sup> (a) V. Corce, L. –M. Chamoreau, E. Derat, J. –P. Goddard, C. Ollivier, L. Fensterbank, *Angew. Chem. Int. Ed.* 2015, 54, 11414–11418. (b)
 M. Jouffroy, D. N. Primer, G. A. Molander, *J. Am. Chem. Soc.* 2016, *138*, 475–478. (c) K. Nakajima, S. Nojima, Y. Nishibayashi, *Angew. Chem. Int. Ed.* 2016, *55*, 14106–14316. (d) A. Gutierrez-Bonet, J. C. Tellis, J. K. Matsui, B. A. Vara, G. A. Molander, *ACS Catal.* 2016, *6*, 8004–8008.
 <sup>16</sup> Photoredox catalysis reviews: (a) J. C. Tellis, C. B. Kelly, D. N. Primer, M. Jouffroy, N. R. Patel, G. A. Molander, *Acc. Chem. Res.* 2016, *49*, 1429–1439. (b) J. K. Matsui, S. B. Lang, D. R. Heitz, G. A. Molander, *ACS Catal.* 2017, *7*, 2563–2575. (c) C. K. Prier, D. A. Rankic, D. W. C.

MacMillan, Chem. Rev. 2013, 113, 5322–5363. (d) M. N. Hopkinson, B. Sahoo, J. –L. Li, F. Glorius, Chem. Eur. J. 2014, 20, 3874–3886.

- <sup>17</sup> N. R. Patel, C. B. Kelly, M. Jouffroy, G. A. Molander, *Org. Lett.* **2016**, *18*, 764–767.
- <sup>18</sup> S. Zheng, D. N. Primer, G. A. Molander, *ACS Catal.* **2017**, *7*, 7957–7961.
- <sup>19</sup> (a) J. Amani, E. Sodagar, G. A. Molander, Org. Lett. **2016**, *18*, 732–735. (b) J. Amani, G. A. Molander, Org. Lett. **2017**, *82*, 1856–1863.
- <sup>20</sup> J. Amani, G. A. Molander, *Org. Lett.* **2017**, *19*, 3612–3615.
- <sup>21</sup> J. Amani, R. Alam, S. Badir, G. A. Molander, *Org. Lett.* **2017**, *1*9, 2426–2429.
- <sup>22</sup> S. Z. Tasker, E. A. Standley, T. F. Jamison, *Nature* **2014**, *509*, 299–309.

<sup>23</sup> B. M. Trost, G. A. Molander, J. Am. Chem. Soc. **1981**, 103, 5969–5972. For an extensive review article, see: J. He, J. Ling, P. Chiu, Chem. Rev. **2014**, 114, 8037–8128.

<sup>24</sup> S. Crotti, F. Bertolini, F. Macchia, M. Pineschi, *Org. Lett.* **2009**, *11*, 3762. During the course of our studies, Gong and coworkers published a method describing a reductive allylation of tertiary alkyl halides using allylic carbonates: H. Chen, X. Jia, Y. Yu, Q. Qian, H. Gong, *Angew. Chem. Int. Ed.* **2017**, *56*, 13103–13106. Their report includes deuterium studies supporting the formation of a π-allylnickel intermediate.

<sup>25</sup> Secondary alkyltrifluoroborates have significantly high reduction potentials at +1.50 V vs SCE (see D. N. Primer, K. Karakaya, J. C. Tellis, G. A. Molander, *J. Am. Chem. Soc.* **2015**, *137*, 2195–2198).

<sup>26</sup> (a) N. Tewari, N. Dwivedi, R. Tripathi, *Tetrahedron Lett.* 2004, 45, 9011–9014. (b) H. Adibi, H. A. Samimi, M. Beygzadeh, *Catal. Commun.* 2007, 8, 2119–2124. (c) A. Heydari, S. Khaksar, M. Tajbakhsh, H. R. Bijanzadeh, *J. Fluorine Chem.* 2009, 130, 609–614.(d) D. Zhang, L.-Z. Wu, L. Zhou, X. Han, Q.-Z. Yang, L.-P. Zhang, C.-H. Tung, *J. Am. Chem. Soc.* 2004, 126, 3440–3441. (e) X. Wei, L. Wang, W. Jia, S. Du, L. Wu, Q. Liu, *Chin. J. Chem.* 2014, 32, 1245–1250. (f) S. Fukuzumi, T. Suenobu, M. Patz, T. Hirasaka, S. Itoh, M. Fujitsuka, O. Ito, *J. Am. Chem. Soc.* 1998, 120, 8060–8068. (g) D. Wang, Q. Liu, B. Chen, L. Zhang, C. Tung, L. Wu, *Chin. Sci. Bull.* 2010, 55, 2855–2858.

<sup>27</sup> (a) K. Nakajima, S. Nojima, K. Sakata, Y. Nishibayashi, *ChemCatChem* **2016**, *8*, 1028–1032. (b) W. Chen, Z. Liu, J. Tian, J. Li, J. Ma, X. Cheng, G. Li, *J. Am. Chem. Soc.* **2016**, *138*, 12312–12315. (c) L. Buzzetti, A. Prieto, S. R. Roy, P. Melchiorre *Angew. Chem. Int. Ed.* **2017**, *56*, 15039–15043. (d) C. Verrier, N. Alandini, C. Pezzetta, M. Moliterno, L. Buzzetti, H. B. Hepburn, A. Vega-Penaloza, M. Silvi, P. Melchiorre, *ACS Catal.* **2018**, *8*, 1062–1066. For a recent review, see: W. Huang, X. Cheng, *Synlett* **2017**, *28*, 148–158.

<sup>28</sup> (a) S. O. Badir, A. Dumoulin, J. K. Matsui, G. A. Molander, *Angew. Chem. Int. Ed.* **2018**, 57, 6610–6613. (b) A. Dumoulin, J. K. Matsui, A. Gutierrez-Bonet, G. A. Molander, *Angew. Chem. Int. Ed.* **2018**, 57, 6614–6618.

<sup>29</sup> N. A. Butt, W. Zhang, Chem. Soc. Rev. 2015, 44, 7929–7967.

<sup>30</sup> Z. G. Khalil, A. A. Salim, D. Vuong, A. Crombie, E. Lacey, A. Blumental, R. J. Capon, *J. Antibiot.* 2017, 70, 1097–1103. (b) Bayliss, M.; M. I. Donaldson, S. A. Nepogodiev, G. Pergolizzi, A. E. Scott, N. J. Harmer, R. A. Field, J. L. Prior, *Carbohydr. Res.* 2017, 452, 17–24. Synthetic efforts: (c) S. Zhao, N. P. Mankad, *Angew. Chem. Int. Ed.* 2018, 130, 5969–5972. (d) A. Szekreyi, X. Garrabou, T. Parella, J. Joglar, J. Bujons, P. Clapes, *Nat. Chem.* 2015, 7, 724–729.

<sup>31</sup> For all of the calculations, dispersion-corrected, broken-spin (U)DFT functionals (UM06/6-311+G(d,p)-SMD(water)//UB3LYP/6-31G(d) and UB3LYP-D3/6-311+G(d,p)-SMD(water)//UB3LYP/6-31G(d)) were used. Further, the energy profile using dynamic correlation, open-shell domain-based local pair natural orbital coupled-cluster calculations (DLPNO-CCSD(T)/def2-TZVPP-SMD(water)//UB3LYP/6-31G(d)) were compared, which are known to provide accurate energies (within 3 kJ mol<sup>-1</sup>) with the computational cost comparable to DFT calculations. Overall, all methods provided similar conclusions (See Supporting Information for further details). For simplicity, only UM06/6-311+G(d,p)-SMD(water)//UB3LYP/6-31G(d) energies will be discussed in the manuscript.

<sup>32</sup> O. Gutierrez, J. C. Tellis, D. N. Primer, G. A. Molander, M. C. Kozlowski, *J. Am. Chem. Soc.* **2015**, *137*, 4 896-4899

<sup>33</sup> We also considered the protonation of **B**" followed by: (i) *outer-sphere*  $C(sp^2)-C(sp^3)$  bond formation and (ii) radical addition, with subsequent *inner sphere*  $C(sp^2)-C(sp^3)$  bond formation. However, the barriers for both protonated *inner* sphere and *outer* sphere pathways were higher in energy than the anionic pathway (see Supporting Information).

## COMMUNICATION



Dr. Jennifer K. Matsui, Dr. Álvaro Gutiérrez-Bonet, Madeline Rotella, Dr. Rauful Alam, Professor Osvaldo Gutierrez and Professor Gary A. Molander

*Shining New Light.* Herein, we report a highly regioselective, intermolecular, nickel-catalyzed photoredox allylic substitution that expands both the radical and electrophile scope of dual photoredox/Ni-catalyzed reactions. Quantum mechanical calculations shed light on the disclosed mechanistic pathway, supporting a Ni(0) to Ni(II) oxidative addition followed by an inner-sphere radical addition.