


# ChemComm

Accepted Manuscript



This article can be cited before page numbers have been issued, to do this please use: Z. Tan, X. Wan, Z. Zang, Q. qian, W. Deng and H. Gong, *Chem. Commun.*, 2014, DOI: 10.1039/C3CC49859J.



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

*Accepted Manuscripts* are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

Cite this: DOI: 10.1039/c0xx00000x

www.rsc.org/xxxxxx

## ARTICLE TYPE

## Ni-Catalyzed Asymmetric Reductive Allylation of Aldehydes with Allylic Carbonates

Zhouzhen Tan, Xiaolong Wan, Zhenhua Zang, Qun Qian,\* Wei Deng\* and Hegui Gong,\*<sup>a</sup>

Received (in XXX, XXX) Xth XXXXXXXXX 20XX, Accepted Xth XXXXXXXXX 20XX

DOI: 10.1039/b000000x

This work features first asymmetric Ni-catalyzed reductive coupling of allylic carbonates with aldehydes, which may proceed via allyl-Ni intermediates although Zn was used as the terminal reductant. Moderate to excellent enantiomeric excess were obtained with excellent functional group tolerance.

Asymmetric carbonyl allylation represents one of the most important reaction types in the current organic synthesis.<sup>1</sup> In particular, the reductive protocols that employ allylic and carbonyl electrophiles without pre-preparation of allylic nucleophiles has received significant advances.<sup>2-9</sup> In addition to the well-established Ni-catalyzed Nozaki-Hiyama-Kishi (NHK) method,<sup>2</sup> In-, Zn-, Et<sub>3</sub>B-, and SnCl<sub>2</sub>-mediated Barbier allylation of aldehydes with allyl halides have also been well studied.<sup>20</sup> However, the catalytic asymmetric Barbier method appears to be difficult, which generally requires stoichiometric amount of chiral auxiliaries.<sup>3</sup> The employment of more accessible and stable allylic alcohols and their derivatives such as allylic acetates and carbonates has drawn increasing attention in realizing highly enantioselective allylation of carbonyl compounds.<sup>4-9</sup> However, the transition metals involved in these reductive umpolung catalytic processes are primarily limited to palladium and iridium.<sup>5-8</sup> For instance, Krische has developed an elegant Ir-catalyzed transfer hydrogenation method allowing alcohols serving as the substrates and reducing reagents, which sets a high bar in the sense of green carbonyl allylation.<sup>6,7</sup> Zanoni and Zhou have demonstrated that catalytic Pd/phosphines in combination with Et<sub>2</sub>Zn and Et<sub>3</sub>B generate homoallylic alcohols with high enantioselectivities, respectively.<sup>8</sup> Therefore, the development of asymmetric reductive umpolung carbonyl allylation using less expensive transition metals, e.g. nickel is still in need,<sup>9</sup> although it has been widely explored in the asymmetric carbonylation of alkynes, dienes and allenes.<sup>10</sup>

In the course of our studies of Ni-catalyzed reductive coupling of alkyl electrophiles with other electrophiles,<sup>11</sup> we have noticed that allylic acetates and carbonates react with aldehydes and ketones in the presence of zinc powder. Herein we disclosed our discovery of Ni-catalyzed asymmetric reductive coupling of allylic carbonates with aldehydes using zinc as the terminal reductant. Moderate to high levels of enantioselectivities were observed, which appears to be significantly effected by the substitution patterns on the allylic carbonates. To the best of


knowledge, this work features the first Ni-catalyzed asymmetric reductive allylation of aldehydes with allylic carbonates. Our preliminary mechanistic studies favored that the enantioselectivities should arise from addition of the weak nucleophilic allyl-Ni to aldehydes.<sup>12</sup>

We first examined the coupling of methyl 2-phenyl allylic carbonate with 4-anisaldehyde (Table 1). After extensive screening of the reaction conditions, we identified that use of NiI<sub>2</sub>, tridentate Pybox ligands and CuI in DMF was superior to other nickel sources, ligands, additives and solvents (entries 1–8). With a combination of NiI<sub>2</sub>/2c/CuI in the presence of zinc powder at 25 °C, **1** was generated in 92% yield and 73% ee (entry 4). The use of Ni(COD)<sub>2</sub> further boosted the ee to 77% (entry 9). Lowering the temperature to 0 °C increased the ee to 86% (entry 10). Ni(ClO<sub>4</sub>)<sub>2</sub>•6H<sub>2</sub>O proved to be more effective than Ni(COD)<sub>2</sub>, which generated **1** in 91% ee even at 25 °C; slight increase of the ee value was observed at 0 °C (entries 11–12). Interestingly, replacement of CuI with CsI delivered equivalent yield and ee (entry 13); without additives, 91% ee could still be attained (entry 14). Moreover, 2-phenyl allylic acetate is equally effective.


Table 1. Optimization of the reaction conditions for **1**.<sup>a</sup>

entry	Ni	ligand	additive	yield (%) <sup>b</sup>	ee (%)
1	NiI <sub>2</sub>	<b>2a</b>	none	90	25
2	NiI <sub>2</sub>	<b>2a</b>	CuI (50%)	90	40
3	NiI <sub>2</sub>	<b>2b</b>	CuI (50%)	95	35
4	NiI <sub>2</sub>	<b>2c</b>	CuI (50%)	92	73
5	NiI <sub>2</sub>	<b>2d</b>	CuI (50%)	95	20
6	NiI <sub>2</sub>	<b>2e</b>	CuI (50%)	76	0
7	NiI <sub>2</sub>	<b>3</b>	CuI (50%)	trace	NA <sup>c</sup>
8	NiI <sub>2</sub>	<b>4</b>	CuI (50%)	38	0
9	Ni(COD) <sub>2</sub>	<b>2c</b>	CuI (50%)	98	77
10	Ni(COD) <sub>2</sub>	<b>2c</b>	CuI (50%)	95	86 <sup>d</sup>
11	Ni(ClO <sub>4</sub> ) <sub>2</sub> •6H <sub>2</sub> O	<b>2c</b>	CuI (50%)	95	92 <sup>d</sup>
12	Ni(ClO <sub>4</sub> ) <sub>2</sub> •6H <sub>2</sub> O	<b>2c</b>	CuI (50%)	95	91
13	Ni(ClO <sub>4</sub> ) <sub>2</sub> •6H <sub>2</sub> O	<b>2c</b>	CsI (50%)	95	91
14	Ni(ClO <sub>4</sub> ) <sub>2</sub> •6H <sub>2</sub> O	<b>2c</b>	none	95	<b>91</b>

<sup>a</sup> Reaction Conditions: aldehyde (100 mol%), 0.15 M in DMF, allylic carbonate (150 mol %), Zn (300 mol %), Ni (10 mol %), Ligand (15 mol %), 25 °C, 12 h. <sup>b</sup> Isolated yields. <sup>c</sup> Not available. <sup>d</sup> 0 °C.



Under the optimized reaction conditions (Table 1, entry 14), the coupling of unsubstituted allylic carbonate with 4-anisaldehyde delivered **5** in 49% ee albeit in high yield. Further optimization indicated that the conditions in Table 1, entry 10 promoted the ee to 66% (Scheme 1). This result suggests that optimal enantioselectivities for the allylic carbonates bearing different substitution patterns may be achieved by modification of the optimized conditions.



**Table 2.** Coupling of 2-aryl allylic carbonates with aldehydes.<sup>a,b</sup>

<b>6</b> , yield: 96% ee: 88%	<b>7</b> , yield: 95% ee: 85%	<b>8</b> , yield: 95% ee: 88%	<b>9</b> , yield: 85% ee: 86%
<b>10</b> , yield: 48% ee: 85%	<b>11</b> , yield: 95% ee: 80%	<b>12</b> , yield: 94% ee: 87%	<b>13</b> , yield: 95% ee: 73%
<b>14</b> , yield: 95% (77%) <sup>c</sup> ee: 76% (85%) <sup>c</sup>	<b>15</b> , yield: 94% ee: 77%	<b>16</b> , yield: 95% ee: 71%	<b>17</b> , yield: 95% ee: 69%
<b>18</b> , yield: 91% ee: 85%	<b>19</b> (R = H), yield: 95% ee: 85%	<b>21</b> , yield: 95% (52%) <sup>c</sup> ee: 68% (80%) <sup>c</sup>	<b>22</b> , yield: 95% (60%) <sup>d</sup> ee: 72% (81%) <sup>d</sup>
<b>20</b> (R = Me), yield: 87% ee: 91%			
<b>23</b> (R = F), yield: 60%, ee: 81%	<b>25</b> , yield: 96% ee: 60%	<b>26</b> , yield: 53% ee: 41%	<b>27</b> , yield: 45% ee: 51%
<b>24</b> (R = CF <sub>3</sub> ), yield: 90%, ee: 82%			

<sup>a</sup> Reaction Conditions: as in Table 1, entry 14. <sup>b</sup> Isolated yields. <sup>c</sup> With CsI (50 mol%) at -25 °C. <sup>d</sup> With CsI (50 mol%) at -15 °C.

Application of the optimized conditions (Table 1, entry 14) to the coupling of other 2-aryl allylic acetates with a variety of aldehydes was performed next (Table 2). The aromatic aldehydes that do not bear electron-withdrawing groups generally produced high ees as in **6–12**; this includes sterically hindered 2-*i*Pr-benzaldehyde. The naphthyl aldehyde produced **13** in 73% ee. The Cl- and Br-substituted benzaldehydes gave similar ees regardless of the substitution patterns as evident in **14** and **15**. On the other hand, the electron-withdrawing groups appeared to reduce the values of ees as evident in **16–17**. Cinnamaldehyde and (*E*)-but-2-enal also gave rise to the homoallylic alcohols **18–20** in high ees. Aliphatic 3-phenylpropanal delivered **21** in 68% ee. The coupling of benzaldehyde with 2-(4-methoxy), 2-(4-fluoro) and 2-(4-trifluoromethyl)phenyl allylic carbonates and 2-methyl allylic carbonate produced **22–25** in 72%, 81%, 82% and 60% ees, suggesting that electronic nature of the allylic partners is important in control of enantioselectivities. The sterically more hindered 2-(2-methyl)phenyl and 2-isopropyl allylic carbonates

diminished ees as evident in **26** and **27**. The addition of CsI at -25 °C boosted the ee for **14** and **21** to 85% and 80%, respectively. Likewise, the ee for **22** was enhanced to 81% at -15 °C. However, lowering the temperatures for **6**, **8**, **11**, **19**, **21**, **22** and **25** did not result in better ees.

Using the optimized conditions (Table 1, entry 14) for the reductive coupling of aromatic aldehydes with 1- and 3-substituted carbonates delivered the homoallylic products **28–30** in excellent yields (Table 3). Excellent anti/syn selectivities were observed for aryl-substituted allylic carbonates (Table 3, entries 1–3), whereas poor anti/syn selectivities were obtained for carbonates bearing methyl substituents (Table 3, entries 4–5). In general, the ees were moderate even when the temperature was lowered to -25 °C (Table 3, entries 2 and 3).

**Table 3.** Carbonyl allylation of other allylic carbonates.<sup>a,b,c,d</sup>

entry	substrate	product	result
1	Ph-CH=CH-OCO <sub>2</sub> Me		yield: 95% anti/syn: 99/1 ee: 64%
2	Ph-CH=CH-OCO <sub>2</sub> Me		yield: 95 (67%) anti/syn: 97/3 (97/3) ee: 50 (64%)
3	Ph-CH=CH-OCO <sub>2</sub> Me		yield: 85 (67%) anti/syn: 97/3 (97/3) ee: 53 (64%)
4	Me-CH=CH-OCO <sub>2</sub> Me ( <i>E</i> : <i>Z</i> = 1:1)		yield: 95% anti/syn: 59/41 ee: 60 (anti), 60 (syn)
5	Me-CH=CH-OCO <sub>2</sub> Me		yield: 95% anti/syn: 59/41 ee: 66 (anti), 66 (syn)

<sup>a</sup> Reaction Conditions: as in Table 1, entry 14. <sup>b</sup> Isolated yields. <sup>c</sup> The anti/syn ratios were determined by <sup>1</sup>H NMR analysis. <sup>d</sup> The absolute stereochemistry is not determined. <sup>e</sup> The reaction was run at -25 °C.

It was interesting to note that coupling of 1-phenyl allylic carbonate with benzaldehyde gave **29** with equivalent results as the 3-phenyl analog (Table 3, entries 2–3). Likewise, similar yields, drs and ees were observed for the coupling of 3-methyl- and 1-methyl allylic carbonates (Table 3, entries 4–5), supporting that the formation of Ni- $\pi$ -allyl complexes is one of the key steps in the catalytic process. Low diastereoselectivities were observed for 1- or 3-methyl-substituted allylic carbonates (Table 3, entries 4–5), indicating that a possible equilibrium between  $\eta^1$ -(*E*)-allyl-Ni and (*Z*)-allyl-Ni when alkyl substituents are present at C1 or C3 positions of allylic carbonates.<sup>13</sup>


**Table 4.** Coupling of 4-anisaldehyde with methyl 2-phenyl allylic carbonate generating **1** without Zn powder.

entry	Ni(COD) <sub>2</sub>	ligand <b>2c</b>	additive	yield/ee
1	150 mol %	none	none	90/0
2	150 mol %	150 mol %	none	90/35
3	150 mol %	15 mol %	Ni(ClO <sub>4</sub> ) <sub>2</sub> (10%)	93/91
4	150 mol %	15 mol %	Zn(ClO <sub>4</sub> ) <sub>2</sub> (10%)	90/91
5	150 mol %	15 mol %	CsI (100%)	90/30

Transformation of the weak nucleophilic allyl-Ni intermediate into more reactive allyl-Zn is possible by reductive transmetalation of allyl-Ni with Zn or by transmetalation of allyl-Ni with Zn<sup>2+</sup>.<sup>9a,14</sup> It is therefore important to identify whether chiral allyl-Ni(II) was capable of adding to the aldehydes.<sup>12</sup> The coupling of 4-anisaldehyde with 2-phenyl allylcarbonate using 1.5 equiv of Ni(COD)<sub>2</sub> in the absence of Zn

and ligand generated **1** in 90% yield (Table 4, entry 1). With 1.5 equiv of ligand **2c**, 35% ee was obtained without eroding the yield. Interestingly, addition of 10% of  $\text{Ni}(\text{ClO}_4)_2$  or  $\text{Zn}(\text{ClO}_4)_2$  drastically increased the ees to 91%, indicating the important role of  $\text{ClO}_4^-$  (entries 3–4). The presence of 100% CsI also provided **1** with 30% ee (entry 5). These results suggest that transformation of allyl-Ni to allyl-Zn is not necessary for this coupling event.<sup>14</sup>

In addition, treatment of 4-anisaldehyde with allylbromide in the presence or absence of  $\text{Ni}(\text{ClO}_4)_2 \cdot 6\text{H}_2\text{O}$  led to **5** in excellent yields (Scheme 2). No enantioselectivities were observed in both cases, owing to in situ formation of allyl-zinc reagents that react with aldehyde through a Barbier mechanism.<sup>15</sup> As a result, we reason that in our Ni-catalyzed reductive coupling process, the enantioselectivity should not arise from allyl-zinc reagents, although their in situ formation cannot be excluded.



Conditions A:  $\text{Ni}(\text{ClO}_4)_2 \cdot 6\text{H}_2\text{O}$  (10 mol %), **2c** (15 mol %), Zn (300 mol %), DMF, rt. Conditions B: **2c** (15 mol %), Zn (300 mol %), DMF, rt.

## Scheme 2. Coupling of allylbromide with 4-anisaldehyde

In summary, we have disclosed asymmetric Ni-catalyzed reductive coupling of allylic carbonates with aldehydes utilizing zinc powder as the terminal reductant. The reaction conditions are particularly effective for the 2-aryl-allylic carbonates which generate the homoallylic alcohols in good to excellent ees for both aromatic and aliphatic aldehydes. The preliminary studies suggest that the enantioselectivity arises from addition of allyl-Ni to aldehydes rather than the more reactive allyl-Zn that may be produced in the reactions.

We thank Ms. Yijing Dai for initial discovery of this reaction, Prof. Yongjian Zhang (Shanghai Jiaotong University) for helpful advices, and Dr Hongmei Deng (Shanghai University) for helping use of the NMR facilities. Financial support was provided by the Chinese NSF (Nos.21172140, 21372151, 21174081 and 21102088), and the Program for Professor of Special Appointment at Shanghai Institutions of Higher Learning Shanghai Education Committee.

## Notes and references

<sup>a</sup> Department of Chemistry, Innovative Drug Research Center, and School of Materials Science and Engineering, Shanghai University, 99 Shang-Da Road, Shanghai 200444, China. E-mail: Hegui\_gong@shu.edu.cn  
† Electronic Supplementary Information (ESI) available: Characterization of all new compounds and HPLC data for ees. See DOI: 10.1039/b000000x/

- For reviews on enantioselective carbonyl allylation, see: (a) M. Yus, J. C. González-Gómez and F. Foubelo, *Chem. Rev.* 2011, **111**, 7774. (b) I. Marek and G. Sklute, *Chem. Commun.* 2007, **43**, 1683. (c) D. G. Hall, *Synlett* 2007, 1644. (d) S. E. Denmark and J. Fu, *Chem. Rev.* 2003, **103**, 2763. (e) J. W. J. Kennedy and D. G. Hall, *Angew. Chem., Int. Ed.* 2003, **42**, 4732. (f) P. V. Ramachandran, *Aldrichimica Acta* 2002, **35**, 23.
- For selected examples of asymmetric NHK reactions, see: (a) G. Xia and H. Yamamoto, *J. Am. Chem. Soc.* 2006, **128**, 2554. (b) Q.-H. Deng, H. Wadepohl, L. H. Gade, *Chem. Eur. J.* 2011, **17**, 14922. (c) K. C. Harper, M. S. Sigman, *Science*, 2011, **333**, 1875. (d) M. Bandini, P. G. Cozzi, P. Melchiorre and A. Umani-Ronchi, *Angew. Chem. Int. Ed.* 1999, **38**, 3357. (d) Z. Zhang, S. Aubry, Y. Kishi, *Org.*

- Lett.* 2011, **13**, 3077. (e) H. Guo, C.-G. Dong, D.-S. Kim, D. Urabe, J. Wang, J. T. Kim, X. Liu, T. Sasaki and Y. Kishi, *J. Am. Chem. Soc.* 2009, **131**, 15387. (f) A. Berkessel, D. Menche, C. A. Sklorz, M. Schroder, I. Paterson, *Angew. Chem., Int. Ed.* 2003, **42**, 1032.
- For selected examples of asymmetric Barbier reactions, see: (a) T. D. Haddad, L. C. Hirayama, and B. Singaram *J. Org. Chem.* 2010, **75**, 642. (b) L. C. Hirayama, S. Gamsey, D. Kneuppel, D. Steiner, D. Torre and K.; Singaram, B. *Tetrahedron Lett.* 2005, **46**, 2315.
- Non-asymmetric coupling of allylic alcohol/thioether with carbonyl compounds, using Rh/B: (a) F. J. Williams, R. E. Grote and E. R. Jarvo, *Chem. Commun.*, 2012, **48**, 1496. Co/Zn: (b) P. Gomes, C. Gosmini and J. Périchon, *Synthesis* 2003, 1909. Fe-electrochemical reduction: (c) M. Durandetti, C. Meignein and J. Périchon, *J. Org. Chem.* 2003, **68**, 3121. Fe/Mn: (d) M. Durandetti, and J. Périchon, *Tetrahedron Lett.* 2006, **47**, 6255. Ti-mediated modulated by Ni or Pd: (e) A. G. Campaa, B. Bazdi, N. Fuentes, R. Robles, J. M. Cuerva, J. E. Oltra, S. Porcel and A. M. Echavarren, *Angew. Chem., Int. Ed.* 2008, **47**, 7515. Rh/CO: (f) M. Vasylyev and H. Alper, *J. Org. Chem.* 2010, **75**, 2710. Ru/CO: (g) S. E. Denmark and S. T. Nguyen, *Org. Lett.* 2009, **11**, 781. Ti: (h) T. Takeda, M. Yamamoto, S. Yoshida and M. Tsubouchi, *Angew. Chem., Int. Ed.* 2012, **51**, 7263. [Ir(cod)Cl]<sub>2</sub>/SnCl<sub>2</sub>: (i) S. Roy and M. Banerjee, *J. Mol. Catal. A* 2006, **246**, 231.
- For selected examples of Pd-catalyzed non-asymmetric carbonyl allylation with allylic alcohols and acetates, Pd/Et<sub>2</sub>Zn: (a) A. Flahaut, K. Toutah, P. Mangeney and S. Roland, *Eur. J. Inorg. Chem.* 2009, 5422. (b) M. Kimura, M. Shimizu, K. Shibata, M. Tazoe, Y. Tamaru, *Angew. Chem., Int. Ed.* 2003, **42**, 3392. Pd(OAc)<sub>2</sub>/Et<sub>2</sub>B: (c) M. Kimura, T. Tomizawa, Y. Horino, S. Tanaka, and Y. Tamaru, *Tetrahedron Lett.* 2000, **41**, 3627. Pd/diboron via allyl-B: (d) N. Selander, A. Kipke, S. Sebelius and K. Szabó, *J. Am. Chem. Soc.* 2007, **129**, 13723.
- For reviews of asymmetric umpolung reactions, see: (a) J. F. Bower, I.-S. Kim, R. L. Patman and M. J. Krische, *Angew. Chem., Int. Ed.* 2009, **48**, 34–46. (b) N. T. Patil and Y. Yamamoto, *Chemtracts* 2004, **17**, 600.
- For examples, see: (a) I.-S. Kim, M.-Y. Ngai and M. J. Krische, *J. Am. Chem. Soc.* 2008, **130**, 6340. (b) Kim, I. S.; M.-Y. Ngai and M. J. Krische, *J. Am. Chem. Soc.* 2008, **130**, 14891–14899. Ir-allylene/alcohol: (c) J. F. Bower, E. Skucas, R. L. Patman and M. J. Krische, *J. Am. Chem. Soc.* 2007, **129**, 15134.
- Pd-catalyzed asymmetric umpolung, Pd/Et<sub>2</sub>Zn: (a) G. Zanoni, S. Gladiali, A. Marchetti, P. Piccinini, I. Tredici and G. Vidari, *Angew. Chem., Int. Ed.* 2004, **43**, 846. Pd/B: (b) S.-F. Zhu, X.-C. Qiao, Y.-Z. Zhang, L.-X. Wang, Q.-L. Zhou, *Chem. Sci.* 2011, **2**, 1135. (c) S. F. Zhu, Y. Yang, L.-X. Wang, B. Liu, and Q.-L. Zhou, *Org. Lett.* 2005, **7**, 2333.
- Non-asymmetric carbonyl allylation with allyl-OR under Ni/InI: (a) T. Hirashita, S. Kambe, H. Tsuji, H. Omori, and S. Araki, *J. Org. Chem.* 2004, **69**, 5054. Intramolecular coupling of allyl ether with aldehyde: (b) D. Franco, K. Wenger, S. Antonczak, D. Cabrol-Bass, E. Dunach, M. Rocamora, M. Gomez, and G. Muller, *Chem.—Eur. J.* 2002, **8**, 664.
- J. Montgomery, *Angew. Chem. Int. Ed.* 2004, **43**, 3890
- (a) Y. Dai, F. Wu, Z. Zang, H. You and H. Gong, *Chem.—Eur. J.* 2012, **16**, 808. (b) X. Yu, T. Yang, S. Wang, H. Xu and Gong, H. *Org. Lett.* 2011, **13**, 2138. (c) H. Xu, C. Zhao, Q. Qian, W. Deng and H. Gong, *Chem. Sci.* 2013, **4**, 4022.
- For stoichiometric addition of allyl-Ni to carbonyl, see: (a) E. J. Corey and M. F. Semmelhack, *J. Am. Chem. Soc.* 1967, **89**, 2755. (b) L. S. Hegedus and R. K. Stivers, *J. Am. Chem. Soc.* 1974, **96**, 3250.
- I. S. Kim, S. B. Han, and M. J. Krische, *J. Am. Chem. Soc.* 2009, **131**, 2514.
- (a) For reductive transmetalation of allyl-Pd with In, see: G. Fontana, A. Lubineau and M.-C. Scherrmann, *Org. Biomol. Chem.* 2005, **3**, 1375. (b) S. Durandetti, S. Sibille and J. Perichon, *J. Org. Chem.* 1989, **54**, 2198.
- B. W. Berton, J. H. Shugart, C. A. Hughey, B. P., Conrad and S. M. Perala, *Molecules* 2001, **6**, 655.