This article was downloaded by: [UNSW Library] On: 26 August 2015, At: 00:10 Publisher: Taylor & Francis Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: 5 Howick Place, London, SW1P 1WG



## Journal of Carbohydrate Chemistry

Publication details, including instructions for authors and subscription information: http://www.tandfonline.com/loi/lcar20

# Novel Synthesis and Structures of Amines and Triazole-Derived Glycoside and Nucleoside Derivatives of Phosphanyl Sugar Analogs

Mitsuji Yamashita , Kazumitsu Suzuki , Yukihiro Kato , Akihito Iida , Koichi Ikai , Putta Mallikarjuna Reddy & Tatsuo Oshikawa

<sup>a</sup> Department of Materials Chemistry, Faculty of Engineering, Shizuoka University, Hamamatsu 432-8561, Japan Published online: 27 Feb 2008.

To cite this article: Mitsuji Yamashita , Kazumitsu Suzuki , Yukihiro Kato , Akihito Iida , Koichi Ikai , Putta Mallikarjuna Reddy & Tatsuo Oshikawa (1999) Novel Synthesis and Structures of Amines and Triazole-Derived Glycoside and Nucleoside Derivatives of Phosphanyl Sugar Analogs, Journal of Carbohydrate Chemistry, 18:8, 915-935, DOI: <u>10.1080/07328309908544044</u>

To link to this article: <u>http://dx.doi.org/10.1080/07328309908544044</u>

### PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing,

systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <u>http://www.tandfonline.com/</u>page/terms-and-conditions

# NOVEL SYNTHESIS AND STRUCTURES OF AMINES AND TRIAZOLE-DERIVED GLYCOSIDE AND NUCLEOSIDE DERIVATIVES OF PHOSPHANYL SUGAR ANALOGS<sup>1</sup>

Mitsuji Yamashita,\* Kazumitsu Suzuki, Yukihiro Kato, Akihito Iida, Koichi Ikai, Putta Mallikarjuna Reddy, and Tatsuo Oshikawa

Department of Materials Chemistry, Faculty of Engineering, Shizuoka University, Hamamatsu 432-8561, Japan

Received January 20, 1999 - Final Form June 29, 1999

#### ABSTRACT

3-Methyl-1-phenyl-2-phospholene and 1-phenyl-2-phospholene 1-oxides were converted into 2-bromo-3-hydroxy-3-methyl-1-phenylphospholane and 2-bromo-3-hydroxy-1-phenylphospholane 1-oxide (1-bromo-1,3,4-trideoxy-1,4-C-[(R, S)-phenylphosphi-nylidene]-glycero-tetrofuranose) by the action of bromine in aqueous medium. The bromo substituent of the phospholane was substituted by treatment with amines or an azide anion to afford novel glycoside derivatives of phosphanyl sugar analogs such as 2-amino-3-hydroxy-1-phenylphospholane (3,4-dideoxy-1,4-C-[(R, S)-phenylphosphinylidene]-glycero-tetrofuranosylamine) and 2-azido-3-hydroxy-3-methyl-1-phenylphospholane 1-oxides with retention of the configuration. The 1,3-dipolar cycloaddition of the 2-azido derivative of the phospholane with alkynes gave 3-hydroxy-3-methyl-1-phenyl-2-(triazol-1'-yl)phospholane 1-oxides as a novel triazole-derived nucleoside of phosphanyl sugar analogs. The structure of the glycoside and nucleoside derivatives of the phosphonyl sugar analogs prepared was deterimined from IR, NMR, and X-ray crystallography analysis.

#### INTRODUCTION

Normal sugar derivatives, represented by Haworth structures, have a heterocyclic structure with an oxygen atom in the hemiacetal ring. Replacement of the oxygen atom in

the hemiacetal ring of normal sugars by a hetero atom or a carbon atom leads to pseudo sugars, some of which have been widely investigated in the fields of synthetic, biological, and medicinal chemistry. In particular, hetero sugars in which the ring oxygen has been replaced by a nitrogen, sulfur, or selenium atom have been extensively studied and widely developed.<sup>2</sup> Synthesis of phosphanyl sugars, which belong to a category of pseudo sugar derivatives having a phosphorus atom instead of the oxygen atom in the hemiacetal ring of sugars, have been investigated by conventional methodology using sugar starting materials. However, the multistep synthesis methods required to prepare a wide variety of phosphanyl sugar derivatives such as glycosides and nucleosides in order to elucidate structure-activity relationships has limited their development. They have been mainly prepared from sugars as starting materials with suitable reaction sequences of OH group protections, functional group interconversions, C-P bond formation, cyclization with the P atom, deprotection, etc.<sup>3</sup> The fact that amino and thio sugars are known to exist in nature, while phosphanyl sugar derivatives have not yet been found in naturally occurring products may also have delayed the development of phosphanyl sugar chemistry.

In previous papers,<sup>4,5</sup> we reported the synthesis of phosphanyl sugar derivatives starting from 2- and 3-phospholene derivatives, having the unsaturated five membered phosphorus heterocycles. We are further interested in the synthesis of glycosides and/or nucleosides of phosphanyl sugars from phospholenes since O- and N-glycosides of normal sugars exist widely and play important biological roles in nature.<sup>2</sup> Moreover nucleosides such as azidothimidine (AZT), which is an anti-HIV agent with an N<sub>1</sub> substituent on the sugar moiety of the nucleoside,<sup>6</sup> are also biologically important substances. Thus, preparation of sugar derivatives such as glycosides and nucleosides for biological study represents an important research area of carbohydrate chemistry.<sup>7</sup> As bioactive nucleosides. ribavirin, deoxyfluridine, cytosine arabinoside, and 5-fluorodeoxyuridine as well as AZT are known as anti-influenza, or anti-tumor reagents and these nucleosides have normal sugar moieties.<sup>8-11</sup> In addition, some pseudo sugar nucleosides exist in nature and show interesting bioactivities such as anti-bacterial, anti-tumor, and anti-virus activities. Some examples are aristeromycin, cyclaradine, thiothymidine, thioddC, and dioxolane T:12 however, no phosphanyl sugar nucleoside has been reported. The present paper deals with the synthesis and the structure determination of glycoside and nucleoside derivatives of phosphanyl sugar analogs using 2-phospholenes as the starting materials.

### **RESULTS AND DISCUSSION**

Reaction of 3-methyl-1-phenyl-2-phospholene 1-oxide (1A) and 1-phenyl-2-phospholene 1-oxide  $(1B)^{13}$  with bromine or N-bromoacetamide (NBA) in chloroform-water or

acetone-water at room temperature afforded a mixture of *threo* and *erythro* bromohydrin derivatives **2A** and **2B**. Fractional recrystallization of **2B** from chloroform-carbon tetrachloride afforded *threo* 2-bromo-3-hydroxy-1-phenylphospholane 1-oxide (*threo* **2B**; 1bromo-1,3,4-trideoxy-1,4-*C*-[(*R*)-phenylphosphinylidene]- $\beta$ -D-glycero-tetrofuranose and the enantiomer; mp 180-183 °C; yield 43%) and *erythro* 2-bromo-3-hydroxy-1-phenylphospholane 1-oxide (*erythro* **2B**; 1-bromo-1,3,4-trideoxy-1,4-*C*-[(*R*)-phenylphosphinylidene]- $\alpha$ -L-glycero-tetrofuranose and the enantiomer; mp 136-139 °C; yield 24%). Fractional recrystallization of a mixture of *erythro* and *threo* products **2A** (ca. 1 : 3), gave *threo* 2-bromo-3-hydroxy-3-methyl-1-phenylphospholane 1-oxide (*threo* **2A**) in 46% yield.<sup>14</sup>

Reaction of bromohydrin *erythro* **2B** with some primary and secondary amines at 40 °C afforded novel phosphanyl sugar *N*-glycosides, 2-amino-3-hydroxy-1-phenylphospholane 1-oxides (**3Ba-d**; 3,4-dideoxy-1,4-*C*-[(*R*)-phenylphosphinylidene]- $\alpha$ -L-glycerotetrofuranosylamines and the enantiomers; Table 1). Treatment of bromohydrin *erythro* **2B** with triethylamine or primary and secondary amines at room temperature gave almost quantitatively *threo* 2,3-epoxy-1-phenylphospholane 1-oxide (*threo* **4B**; 1,2-anhydro-3,4-dideoxy-1,4-*C*-[(*R*)-phenylphosphinylidene]- $\beta$ -L-glycero-tetrofuranose and the enantiomer). Epoxide *threo* **4B** was smoothly converted into *erythro N*-glycosides **3Ba-d** by the action of primary or secondary amines at 40 °C. At 40 °C, the reaction of **2B** with primary or secondary amines at 40 °C. At 40 °C, the reaction of **2B** with primary or secondary amines at forded *N*-glycosides of phosphanyl sugar analogs **3Ba-d** in one pot through intermediary 1,2-anhydro phosphanyl sugar analogs, *N*-glycosides **3Ba-d**, from **2B** are observed through different reaction conditions, with and without isolation of intermediate epoxide **4B**.

Acyl derivatives of  $\alpha$ -D-glucopyranosyl bromide are known to be often converted by treatment of amines into 2-hydroxyglycals with elimination of hydrogen halide, instead of formation of *N*-glycosides by substitution of the halogen with amines.<sup>15</sup> An S<sub>N</sub>1 type substitution reaction of 2,3,4,6-tetra-*O*-acetyl- $\alpha$ -D-glucopyranosyl bromide with sodium iodide or silver fluoride is reported to give thermally more stable  $\alpha$  or  $\beta$  anomer of the corresponding glucopyranosyl halide when neighbouring group participation of the 2-substituent is available.<sup>16</sup> On the other hand,  $\beta$ -D-glucopyranosyl fluoride was converted into the corresoponding methyl glucoside by treatment with methoxide via epoxide formation by participation of the *trans* 2-hydroxy group in the elimination of the fluoro group to give the 1,2-epoxide derivative, this method being called the Micheel synthesis.<sup>17,18</sup> Preparation of 1,2-anhydro- $\alpha$ -D-glucose or the corresponding tri-*O*-acetyl derivative ("Brigl's anhydride")<sup>18</sup> was carried out using careful ammonolysis of 3,4,6-tri-*O*-acetyl- $\alpha$ -D-glucopyra-



Figure 1. Structure of phospholenes 1A and 1B.

Table 1. Phosphanyl sugar N-glycosides prepared.

Starting materia	$NR^{1}R^{2}R^{3}$		Reaction	condition	Product			
	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Solvent	Temp(°C)	No.	Mp(°C)	Yield(%)
erythro 2B	Me	Н	Η	MeOH-H	<sub>2</sub> O 40	erythro 3Ba	Syrup	56
erythro 2B	i-Pr	Н	Η	MeOH	40	erythro 3Bb	138.5-140	75
erythro 2B	t-Bu	Н	Η	MeOH	40	erythro 3Bc	184-185	55
erythro 2B	Et	Et	Н	MeOH	40	erythro 3Bd	Syrup	84
erythro 2B	Et	Et	Et	Et₃N	rt <sup>a)</sup>	threo 4B	Syrup	100
threo 2B	Et	Et	Et	Et₃N	rt <sup>a)</sup>	erythro 4B	116-118	91
threo 4B	<i>i</i> -Pr	H	Η	MeOH	40	erythro 3Bb	138.5-140	100
erythro 4B	Et	Et	Et	MeOH	40			NR <sup>s)</sup>

a. rt, room temperature; NR, No reaction.

nosyl chloride. Therefore, facile epoxide formation followed by substitution with a nucleophile to give an N-glycoside from phosphanyl sugar derivative 2B by an amine base treatment may be a unique feature in carbohydrate chemistry, because substitution of a halogen of acetylated glycosyl halides most generally occurs by participation of a 2-acetoxy group.



Scheme 1. Preparation of N-glycosides of phosphanyl sugars 3Ba-d.

Chemical shift $\delta$ (ppm)												
H1	H2	H3	H3'	H4	H4'	t-Bu	OH	NH	o-Ph	m-Ph	p-Ph	
2.82	3.97	2.47	1.78	2.33	2.05	0.92	1.86	2.98	7.75	7.50	7.53	
				Cou	pling c	onstant (	Hz) <sup>b)</sup>					
$J_{1,2} = 8.3$ $J_{1,p} = 4$			$_{1,p} = 4.9$	)								
$J_{2,3} = 9.6$ J		$J_{2,3} = 5.0$			$J_{2,P} = 5.0$							
$J_{3,P} = 25.0$ .		$J_3$	$J_{3,3'} = 13.2$			$J_{3,4} = 8.4$			$J_{3,4} = 3.6$			
$J_{3',4} = 7.8$			$J_{z}$	J <sub>3',P</sub> ≒ 8			$J_{3',4'} = 11.0$			$J_{4,4} \doteq 16$		
										$J_{4,\mathrm{P}} = 8$	.0	
									J <sub>4',P</sub> ≒26	5		

Table 2. Observed <sup>1</sup>H NMR (500 MHz) parameters for compound 3Bc in CDCl<sub>3</sub>.<sup>a)</sup>

a. Measured on a VXR-500 instrument (Okayama University) at 21 °C.

b. Coupling constants for aromatic protons are omitted.



Figure 2. Favorable conformation  $({}^{2}E)$  of *N*-glycoside 3Bc based on  ${}^{1}H$  NMR (CDCl<sub>3</sub>) analysis. (Numbers outside the parenthesis correspond to carbohydrate nomenclature numbering while those within the parenthesis correspond to heterocycle nomenclature numbering, respectively.)

The structure of 1-*N*-*t*-butyl-3,4-dideoxy-1,4-*C*-[(*S*)-phenylphosphinylidene]- $\beta$ -Dglycero-tetrofuranosylamine and its enantiomer was established for phosphanyl sugar **3Bc** by analysis of its 500 MHz <sup>1</sup>H NMR spectrum (Table 2). The  $J_{1,2}$  value of 8.3 Hz shows that C-1-H-1 and C-2-H-2 bonds are diaxial, whereas the small  $J_{1,P}$  value of 4.9 Hz indicates a *trans* relationship of C-1-H-1 and P=O bonds.<sup>19,20</sup> The <sup>1</sup>H NMR analysis reveals that *N*-glycoside **3Bc** exisits predominantly in the <sup>2</sup>E conformation in the solution (Figure 2). Reaction of the *threo* 2-bromo derivative, *threo* 2A, with sodium azide in DMF for 24 h at 70 °C afforded *threo* 2-azido-3-hydroxy-3-methyl-1-phenylphospholane 1-oxide (*threo* 3Aa; 1-azido-1,3,4-trideoxy-2-methyl-1,4-C-[(R)-phenylphosphinylidene]- $\beta$ -D-glycero-tetrofuranose and its enantiomer) in 87% yield. Hydrogenolysis of the azido derivative, *threo* 3Aa, with an atmospheric pressure of H<sub>2</sub> in the presence of Pd/C catalyst for 24 h at room temperature gave the corresponding *threo* 2-amino derivative, 2-amino-3-hydroxy-3-methyl-1-phenylphospholane 1-oxide (*threo* 3Ab; 3,4-dideoxy-2-methyl-1,4-C-[(R)-phenylphosphinylidene]- $\alpha$ -L-glycero-tetrofuranosylamine and the enantiomer), in 99% yield (Scheme 2). These products 3Aa, 3Ab, and 3Ba-d are N-glycosides of phosphanyl sugar analogs. The structure of *threo* 2-bromo- and *threo* 2-azido-3-methyl-1-phenylphospholane 1-oxides (2A and 3Aa) are shown in Figures 3 and 4, respectively. The retention of the configuration for the reaction of 2A to afford 3Aa may be explained by the formation of the intermediate epoxide 4A.



Scheme 2. Preparation of N-glycosides 3Aa and 3Ab.

As mentioned in Micheel's method<sup>17,18</sup> for the preparation of glycosides of normal sugars, sodium methoxide in methanol converts  $\beta$ -D-glucopyranosyl fluoride into methyl  $\beta$ -D-glucopyranoside via a 1,2-epoxide ring-opening. 1,6-Anhydro- $\beta$ -D-glucopyranose is usually obtained when phenolic glycosides undergo alkaline hydrolysis. The formation of 1,6-anhydro glucose derivatives is explained by a mechanism where an attack by the 6-hydroxyl group occurs at the C1 position of intermediary 1,2-anhydride formed instead of the attack at the C2 position of the epoxide.<sup>21-23</sup> The epoxide ring opening of 1,2-anhydro phosphanyl sugar **4A** or **4B** to afford 1-azido or 1-amino phosphanyl sugar *N*-glycoside may be attributable to a strong electron-withdrawal property of the P=O group, by which



Figure 3. ORTEP drawing of 2-bromo compound 2A.<sup>24</sup>



Figure 4. ORTEP drawing of 2-azido compound 3Aa.<sup>25</sup>

the C1 position of the 1,2-anhydride becomes more electrophilic than the C2 position for the attack of amines or azide ion, although the C1 position is sterically more hindered by the phenyl group than is the C2 position.

For the preparation of nucleosides, the following two major methodologies are usually applied: (a) the substitution reaction of a leaving substituent on an anomeric carbon atom by an activated nucleic acid base; (b) the cyclization of glycosyl ureas or glycosyl amines of sugar derivatives by a condensation reaction with acrylamides or the dipolar cycloaddition with dipolarophiles such as alkynes to form nucleic acid bases or nitrogen heterocycles.<sup>26-32</sup>



Scheme 3. Formation of epoxide 4A from 2A by treatment with TMS-pyrimidine.

Reaction of 2-bromo-3-hydroxy-3-methyl-1-phenylphospholane 1-oxide (2A) with trimethylsilylated pyrimidine was carried out according to the method (a); however, the reaction afforded only intramolecular substitution reaction product, epoxide 4A, instead of an intermolecular substitution reaction product, a pyrimidine nucleoside.

1,3-Dipolar cycloaddition of azido derivative, *threo* 3Aa, with alkynes 5a-h in 1,2-dimethoxyethane (DME) under reflux afforded regioisomeric phosphanyl sugar nucleoside analogs 6Aa-h and 7Ab,e,f having a triazole ring as the nitrogen heterocyclic nucleus depending on the substituents on the alkyne derivatives (Scheme 4 and Table 3).



Scheme 4. Preparation of nucleosides of phosphanyl sugar analogs 6A and 7A.

Disubstituted alkynes 5 c, 5 d, 5 g, and 5 h afforded triazole derivatives 6Ac, 6Ad, 6Ag, and 6Ah, respectively, whose structures were determined by <sup>1</sup>H, <sup>13</sup>C, and <sup>31</sup>P NMR

Alkyne			Reaction	Phosphanyl sugar nucleoside analogs					
No.	$R^1$	R <sup>2</sup>	time(h)	No.	Yield(%)	Ratio of 6A : 7A	mp(°C)		
5a	Н	SiMe <sub>3</sub>	12	6Aa	64.9	6Aa only <sup>2)</sup>	222		
5 b	Н	COOMe	12	6Ab+7Ai	66.6	1:1 <sup>b)</sup>	221-223 <sup>d)</sup>		
5 c	COOMe	COOMe	18	бAс	88.2		205-207		
5d	COOEt	COOEt	24	6Ad	79.4		175-176		
5 e	Н	CH <sub>2</sub> OH	48	6Ae+7Ae	66.0	$1:1^{c}$	224 <b>-</b> 226 <sup>e)</sup>		
5 f	н	CMe <sub>2</sub> OH	96	6Af+7Af	50.9	3:1°	198-202 <sup>0</sup>		
5 g	CH <sub>2</sub> OH	CH <sub>2</sub> OH	72	6Ag	66.0		206		
5h	COOH	COOH	24	6Ah	55.2		175		

**Table 3.** Phosphanyl sugar nucleoside analogs having a triazole ring as the nitrogen heterocyclic nucleus.

a. Isolated by crystallization.

b. Based on the ratios of 3-Me group and the olefin protons by <sup>1</sup>H NMR.

c. Isolated by column chromatography on silica gel.

d. Melting point for 7Ab.

e. Melting point for 7Ae.

f. Melting point for 6Af.

spectral analysis. Typical observations for triazole derivatives 6Ad compared with azido derivative 3Ad in NMR are as follows: down field chemical shift of N-CH for triazole 6Ad (5.57 ppm) was observed compared with that for azido 3Ad (4.00 ppm) by <sup>1</sup>H NMR (CDCl<sub>3</sub>); triazole  $sp^2$  carbon atoms were observed for 6Ad at 131.79 and 138.34 ppm by <sup>13</sup>C NMR (CDCl<sub>3</sub>); and only one signal was observed for 6Ad at 70.49 ppm by <sup>31</sup>P NMR (CDCl<sub>3</sub>).

<sup>1</sup>H NMR (CDCl<sub>3</sub>) spectra for 6Ae and 7Ae showed the olefinic proton signals at 7.86 and 7.36 ppm, respectively, suggesting that the more polarized isomer resonates at lower field than the less polarized isomer does. Trimethylsilylacetylene (5a) gave nucleo- side 6Aa (70.39 ppm by <sup>31</sup>P NMR, CDCl<sub>3</sub>) selectively as the sole regioisomer,<sup>33</sup> whereas methyl propiolate (5b) produced nucleoside 6Ab and 7Ab in a 1:1 ratio. Frontier  $\pi$ electron densities were obtained for 5a, 5b, and 2-azidophospholane derivative 3Aa by an MO calculation using MOPAC PM3 (Stuart)<sup>34</sup> as shown in Figure 5. For TMS derivative **5a** the unsubstituted *sp* carbon atom has almost no  $\pi$ -electron density while the substituted one exclusively has  $\pi$ -electron density. For propiolate **5b** the unsubstituted *sp* carbon



Figure 5. Comparisons of  $\pi$ -electron densities of trimethylsilylacetylene (5a), methyl propiolate (5b), and 2-azidophospholane derivative 3A a based on MO calculations using MOPAC PM3 (Stuart).

atom has a larger  $\pi$ -electron density than that of substituted *sp* carbon atom. Both effects of steric hindrance and electron density caused by the TMS group of **5a** may introduce the predominant formation of nucleoside **6Aa**. In contrast to **5a**, methyl propiolate (**5b**) has a larger electron density on the unsubstituted and less hindered *sp* carbon, hence 1:1 regio-isomeric products **6Ab** and **7Ab** are formed. Triazolyl phosphanyl sugar nucleosides **6A** and **7A** are the first ribavirin type pseudo sugar nucleosides reported.

Phosphanyl sugar nucleoside analog **6Ad** was purified by recrystallization from chloroform, affording a good single crystal. The X-ray analysis of **6Ad** revealed the structure of the novel nucleoside of the phosphanyl sugar as shown by the ORTEP drawing in Figure 6 and afforded bond lengths, bond angles, and torsion angles as shown in Table 4.

#### EXPERIMENTAL

General methods. Melting points were determined by a Yanagimoto MP-S2 micro-melting point apparatus and are reported uncorrected. <sup>1</sup>H NMR spectra were recorded on Japan Electron Optics Laboratory (JEOL) JNM-EX90 (at 90 MHz), JEOL JNM-EX270 (at 270 MHz), JEOL JNM-EX400 (at 400 MHz), and Varian VXR-500 (at 500 MHz) spectrometers using CDCl<sub>3</sub> and TMS as the solvent and the internal standard, respectively. <sup>13</sup>C NMR spectra were recorded on a JEOL EX90 (at 22.40 MHz)



Figure 6. ORTEP drawing of the nucleoside analog of phosphanyl sugar 6Ad having a triazole nucleus.<sup>35</sup>

spectrometer using CDCl<sub>3</sub> and TMS as the solvent and the internal standard, respectively. <sup>31</sup>P NMR spectra were measured by JEOL JNM-EX90 (at 36.18 MHz) and Varian VXR-500 spectrometers using CDCl<sub>3</sub> and  $H_3PO_4$  as the solvent and the external standard, respectively. IR were recorded on Japan Spectroscopic Co. Ltd. (JASCO) FT/IR-8000 and A-3 spectrophotometers. MS spectra were measured using a Hitachi RMU7M GC-MS mass spectrometer. HPLC were carried out using JASCO UNIFLOW-211 with UVIDEC-100-H, FINEPAC SIL, and MeOH-CHCl<sub>3</sub> (1:20) as the detector, column, and solvent, respectively. X-ray crystallographies for single crystals were performed using Rigaku AFC7R and AFC5S diffractometers.

2-Phospholene 1-oxides 1A and 1B were prepared according to reported methods via cycloaddition reaction of 2-methyl-1,3-butadiene and 1,3-butadiene, respectively, with phosphorus trichloride.<sup>36-40</sup>

Selected bond length		Selected bor	nd angle	Selected torsion angle		
Bond	Length (Å)	Bond A	Angle (°)	Bond	Angle (°)	
P(1)-O(2)	1.496	O(2)-P(1)-C(1)	112.1	P(1)-C(1)-C(2)-C(3)	39.2	
P(1)-C(1)	1.857	O(2)-P(1)-C(6)	111.0	C(1)-C(2)-C(3)-C(4)	-48.9	
P(1)-C(4)	1.806	C(1)-P(1)-C(6)	108.7	C(2)-C(3)-C(4)-P(1)	34.8	
C(1)-C(2)	1.566	P(1)-C(1)-C(2)	103.7	C(3)-C(4)-P(1)-C(1)	-9.5	
C(2)-C(3)	1.548	C(1)-C(2)-C(3)	105.4	C(4)-P(1)-C(1)-C(2)	-17.4	
P(1)-C(6)	1.799	C(2)-C(3)-C(4)	107.8	P(1)-C(1)-N(1)-N(2)	-77.4	
C(1)-N(1)	1.467	P(1)-C(1)-N(1)	114.2	C(2)-C(1)-N(1)-N(2)	42.7	
N(1)-N(2)	1.367	C(2)-C(1)-N(1)	115.6	C(2)-C(1)-N(1)-C(12)	-142.7	
N(2)-N(3)	1.305	N(1)-N(2)-N(3)	107.2	N(1)-N(2)-N(3)-C(13)	-0.9	
N(3)-C(13)	1.359	C(1)-N(1)-N(2)	121.8	N(2)-N(3)-C(13)-C(12	2) 0.3	
C(12)-C(13)	1.359	C(1)-N(1)-C(12)	127.8	N(3)-C(13)-C(12)-N(1	l) 0.3	
N(1)-C(12)	1.338	N(2)-N(3)-C(13)	108.5	C(13)-C(12)-N(1)-N(2	2) -0.9	
C(6)-C(7)	1.392	N(3)-C(13)-C(12)	109.2	C(12)-N(1)-N(2)-N(3)	1.1	
C(7)-C(8)	1.385	C(13)-C(12)-N(1)	105.0	P(1)-C(6)-C(7)-C(8)	179.7	
C(8)-C(9)	1.355	C(4)-P(1)-C(6)	111.0	P(1)-C(6)-C(11)-C(10	) -179.7	
C(9)C(10)	1.371	P(1)-C(6)-C(7)	122.8	C(6)-C(7)-C(8)-C(9)	1.4	
C(10)-C(11)	1.382	P(1)-C(6)-C(11)	119.4	C(7)-C(8)-C(9)-C(10)	-2.0	
C(11)-C(6)	1.377	C(6)-C(7)-C(8)	119.7	C(8)-C(9)-C(10)-C(11	) 1.9	
C(13)-C(17)	1.477	C(7)-C(8)-C(9)	120.5	C(9)-C(10)-C(11)-C(6	i) -1.3	
C(12)-C(14)	1.496	C(8)-C(9)-C(10)	121.5	C(10)-C(11)-C(6)-C(7	7) 0.7	
C(17)-O(4)	1.194	C(9)-C(10)-C(11)	119.5	C(11)-C(6)-C(7)-C(8)	-0.8	
C(2)-O(3)	1.423	C(10)-C(11)-C(6)	120.2	N(3)-C(13)-C(17)-O(	5) 2.4	
C(2)-C(5)	1.519	C(11)-C(6)-C(7)	119.4	C(17)-C(13)-C(12)-C	(14) 0.5	

Table 4. Selected bond lengths, bond angles, and torsion angles for compound 6Ad.<sup>35</sup>

erythro And threo 2-bromo-3-hydroxy-1-phenylphospholane 1-oxides (2B). To 1-phenyl-2-phospholene 1-oxide 1B (3.68 g, 20.7 mmol) in acetone (10 mL)-water (20 mL) was added NBA (4.28 g, 1.5 eq) and stirred at room temperature until the brownish color of the solution disappeared, and the completion of the reaction was confirmed by TLC. Removal of the solvent and addition of carbon tetrachloride to the residue afforded a solid product, whose recrystallization from chloroform-carbon tetrachloride afforded 2.43 g (8.84 mmol) of threo 2B (yield 43%). Fractional crystallization of the

1

mother liquid from chloroform-carbon tetrachloride gave 1.38 g (5.02 mmol) of *erythro* 2B (yield 24%).

For three **2B**: mp 180-183 °C; <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  2.0-2.8 (m, 4H, CH<sub>2</sub>-CH<sub>2</sub>), 4.1-4.8 (m, 2H, C<u>H</u>Br-C<u>H</u>OH), 5.7-6.0 (brs, 1H, OH), 7.3-8.0 (m, 5H, Ph); IR  $\nu$ (KBr) 3200 (OH), 1440 (P-Ph), 1180 (P=O), 540 (C-Br); 750 (P-C); MS (*m/z*) 275(M<sup>+</sup>), 277 (M<sup>+</sup> +2).

Anal. Calcd for C<sub>10</sub>H<sub>12</sub>BrO<sub>2</sub>P (275.1): C, 43.66; H, 4.40; P, 11.26. Found: C, 43.78; H, 4.29; P, 11.20.

For erythro 2B: mp 136-139 °C; <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  1.4-2.8 (m, 4H, CH<sub>2</sub>-CH<sub>2</sub>), 3.9-4.3 (m, 2H, CHBr), 4.7 (dm, 1H, <sup>2</sup>J<sub>HP</sub>=16 Hz, C<u>H</u>OH), 5.8-6.1 (brs, 1H, OH), 7.3-8.0 (m, 5H, Ph); IR  $\nu$  (KBr) 3250 (OH), 1440 (P-Ph), 1180 (P=O), 540 (C-Br).

*threo* Bromo-3-methyl-3-hydroxy-1-phenylphospholane 1-oxide (*threo* 2A). To 3-methyl-1-phenyl-2-phospholene 1-oxide 1A (5.53 g, 28.8 mmol) in chloroform-water (1/4 v/v, 25 mL) was added bromine (3.0 mL, 2 eq) and the reaction mixture stirred for 3 d at room temperature. Work-up of the reaction mixture and fractional recrystallization from ethyl acetate gave 3.83 g (13.2 mmol) of *threo* 2A in 46% yield; mp 150-152 °C. <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  1.67 (s, 3H, Me), 1.9-2.8 (m, 4H, CH<sub>2</sub>CH<sub>2</sub>), 4.24 (d, 1H, <sup>2</sup>J<sub>HP</sub>=5.10 Hz, 5.56 (brs, 1H, OH), 7.3-8.0 (m, 5H, Ph); <sup>13</sup>C NMR (CDCl<sub>3</sub>)  $\delta$  25.46 (d,  $J_{CP}$ =64.83 Hz, C-5), 26.01 (d, <sup>3</sup> $J_{CP}$ =6.68 Hz, Me), 35.72 (d, <sup>2</sup> $J_{CP}$ =3.34 Hz, C-4), 50.12 (d,  $J_{CP}$ =65.50 Hz, C-2), 79.43 (d, <sup>3</sup> $J_{CP}$ =12.70 Hz, C-3), 128.02 (d, <sup>3</sup> $J_{CP}$ =12.03 Hz, *m*-Ph), 131.88 (d, <sup>2</sup> $J_{CP}$ =8.70 Hz, *o*-Ph), 132.43 (d, <sup>4</sup> $J_{CP}$ =2.69 Hz, *p*-Ph); <sup>31</sup>P NMR (CDCl<sub>3</sub>)  $\delta$  68.937; IR  $\nu$  (KBr) 3150 (OH).

erythro 2-(N, N'-Diethylamino)-3-hydroxy-1-phenylphospholane 1oxide (erythro 3Bd). Reaction of bromohydrin 2B (89.4 mg, 0.325 mmol) with excess N, N'-diethylamine (2 mL) in methanol (3 mL) for 2 d at 40 °C gave N-glycoside product erythro 3Bd. Removal of the volatile materials from the reaction mixture followed by washing the chloroform solution of the resulting residue with diluted sodium hydroxide (3 mL), further extraction of the water layer with chloroform, and chromatography of the residue on silica gel gave pure product 3Bd (72.9 g, 0.273 mmol) in 84% yield. <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  0.95 (t, 6H, J=7.0 Hz, 2 x CH<sub>2</sub>CH<sub>3</sub>), 1.5-2.6 (m, 4H, CH<sub>2</sub>CH<sub>2</sub>), 2.7-3.3 (m, 5H, C<u>HNEt<sub>2</sub></u>, 2 x C<u>H<sub>2</sub>CH<sub>3</sub></u>), 3.6-4.0 (brs, 1H, OH), 4.3-4.8 (m, 1H, C<u>H</u>OH), 7.4-8.1 (m, 5H, Ph); IR  $\nu$  (neat) 3300 (OH), 1440 (P-Ph), 1185 (P=O); MS (m/z) 267 (M<sup>+</sup>).

The same procedure gave amino derivatives of phosphanyl sugar glycosides.

For methylamine derivative **3Ba**: <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  1.3-2.6 (m, 4H, CH<sub>2</sub>CH<sub>2</sub>), 2.32 (s, 3H, NMe), 2.76 (d, 1H, *J*=7.0 Hz, CH), 3.5-4.0 (m, 3H, NH, CHOH), 7.1-8.0 (m, 5H, Ph); IR  $\nu$  (neat) 3300 (OH, NH), 1445 (P-Ph), 1180 (P=O), 770 and 700 (Ph); MS (*m/z*) 226 (M<sup>+</sup>). For isopropylamine derivative **3B**b: <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  0.87 (2d, 5H, J=6.0 Hz, C<u>Me<sub>2</sub>H</u>), 1.5-3.1 (m, 5H, CH<sub>2</sub>CH<sub>2</sub>, CMe<sub>2</sub>H), 3.4-4.3 (brs, 2H, NH, OH), 7.4-8.0 (m, 5H, Ph); IR  $\nu$  (KBr) 3250, 3450 (OH, NH), 1440 (P-Ph), 1160 (P=O), 770 and 700 (Ph); MS (*m*/z) 254 (M<sup>+</sup>).

For tert-butylamine derivative **3Bc**: <sup>1</sup>H NMR (CDCl<sub>3</sub>) data are shown in Table 2. IR  $\nu$  (KBr) 3280, 3250 (OH, NH), 1440 (P-Ph), 1160 (P=O), 770 and 700 (Ph); MS (*m/z*) 268 (M<sup>+</sup>).

erythro 2,3-Epoxy-1-phenylphospholane 1-oxide (erythro 4B). threo Bromohydrin 2B (1.31 g, 4.75 mmol) was treated with triethylamine (4 mL) in methanol (10 mL) for 2 d at room temperature. The reaction mixture was concentrated *in vacuo* and the residue was dissolved into chloroform (40 mL), whose solution was washed with aqueous sodium hydrogenearbonate solution (10 mL) and water (2 x 10 mL) and dried and concentrated under reduced pressure to give crystalline product erythro 4B (recrystallized from carbon tetrachloride; 0.897 g, 4.31 mmol) in 91% yield; mp 116-118 °C. <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  1.6-3.0 (m, 4H, CH<sub>2</sub>CH<sub>2</sub>), 3.39 (dd, <sup>2</sup>J<sub>HP</sub>=30 Hz, P(O)CH), 3.7-4.0 (m, 1H, CH<sub>2</sub>C<u>H</u>), 7.2-8.0 (m, 5H, Ph); IR  $\nu$  (KBr) 1440 (P-Ph), 1240 and 835 (epoxide), 1200 (P=O).

Anal. Calcd for C<sub>10</sub>H<sub>11</sub>O<sub>2</sub>P (194.2): C, 61.86; H, 5.71; P, 15.95 Found: C, 61.73; H, 5.64; P, 15.82.

*threo* 2-Azido-3-hydroxy-3-methyl-1-phenylphospholane 1-oxide (*threo* 3Aa). *threo* 2-Bromo-3-hydroxy-3-methyl-1-phenylphosphlane 1-oxide (*threo* 2A, 1.36 g, 4.71 mmol) was stirred for 1 d at 70 °C with sodium azide in DMF (30 mL). Removal of volatile materials from the reaction mixture followed by washing of the chloroform solution (50 mL) of the residue with water (3 x 20 mL) and removal of the solvent afforded crystalline azido derivative *threo* 3Aa (1.03 g, 4.10 mmol) in 87% yield; mp 174-176 °C. <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  1.53 (s, 3H, Me), 1.6-2.6 (m, 4H, CH<sub>2</sub>CH<sub>2</sub>), 4.00 (d, 1H, <sup>2</sup>J<sub>HP</sub>=1.98 Hz, CHN<sub>3</sub>), 5.80 (brs, 1H, OH), 7.4-8.0 (m, 5H, Ph); <sup>13</sup>C NMR (CDCl<sub>3</sub>)  $\delta$  24.24 (d, <sup>3</sup>J<sub>CP</sub>=7.35 Hz, Me), 26.39 (d, J<sub>CP</sub>=63.48 Hz, C-5), 36.23 (d, <sup>2</sup>J<sub>CP</sub>=4.68 Hz, C-4), 67.89 (d, J<sub>CP</sub>=72.84 Hz, C-2), 78.45 (d, <sup>2</sup>J<sub>CP</sub>=12.03 Hz, C-3), 128.60 (d, <sup>3</sup>J<sub>CP</sub>=11.37 Hz, *m*-Ph), 129.24 (d, J<sub>CP</sub>=93.54 Hz, *x*-Ph), 131.54 (d, <sup>2</sup>J<sub>CP</sub>=9.36 Hz, *o*-Ph), 132.61 (d, <sup>4</sup>J<sub>CP</sub>=2.67 Hz, *p*-Ph); <sup>31</sup>P NMR (CDCl<sub>3</sub>)  $\delta$  67.286; IR  $\nu$  (KBr) 3150 (OH), 2120 (N<sub>3</sub>).

threo (1R,2S, 3R)-2-(4',5'-Diethoxycarbonyl-1'H-1',2',3'-triazol-1'yl)-3-hydroxy-3-methyl-1-phenylphospholane 1-oxide (6Ad). A mixture of 2-azido-3-hydroxy-3-methyl-1-phenylphospholane 1-oxide (threo 3Aa, 0.202 g, 0.804 mmol) and diethyl acetylenedicarboxylate (0.203 g, 1.19 mmol) in DME (3 mL) was refluxed for 1 d. The solvent was then removed *in vacuo* and chromatography on silica gel afforded 0.269 g (0.638 mmol) of triazole derivarive of nucleoside of phosphanyl sugar 6Ad in 79% yield; mp 175-176 °C. <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  1.31 (s, 3H, Me), 1.33 and 1.36 (2t, 6H, *J*=6.6 Hz, 2 x OCH<sub>2</sub>CH<sub>3</sub>), 2.4-3.0 (m, 4H, CH<sub>2</sub>CH<sub>2</sub>), 4.32 and 4.41 (2q, 4H *J*=6.6 Hz, 2 x OCH<sub>2</sub>CH<sub>3</sub>), 5.57 (d, 1H, <sup>2</sup>J<sub>HP</sub>=9.99 Hz, CH), 6.15 (brs, 1H, OH), 7.3-7.8 (m, 5H, Ph); <sup>13</sup>C NMR (CDCl<sub>3</sub>)  $\delta$  13.69 and 13.90 (2s, 2 x OCH<sub>2</sub>CH<sub>3</sub>), 23.45 (d, <sup>3</sup>J<sub>CP</sub>=6.68 Hz, Me), 24.82 (d, *J*<sub>CP</sub>=68.16 Hz, C-5, 37.56 (d, <sup>2</sup>J<sub>CP</sub>=4.68 Hz, C-4), 61.55 and 63.07 (2s, 2 x OCH<sub>2</sub>CH<sub>3</sub>), 68.05 (d, *J*<sub>CP</sub>=66.82 Hz, C-2), 79.89 (d, <sup>2</sup>J<sub>CP</sub>=16.04 Hz, C-3), 127.84 (d, *J*<sub>CP</sub>=93.56 Hz, *x*-Ph), 128.02 (d, <sup>3</sup>J<sub>CP</sub>=12.03 Hz, *m*-Ph), 131.14 (d, <sup>2</sup>J<sub>CP</sub>=10.01 Hz, *o*-Ph), 131.79 (d, <sup>3</sup>J<sub>CP</sub>=2.02 Hz, triazole C-5'), 132.52 (d, <sup>4</sup>J<sub>CP</sub>=2.69 Hz, *p*-Ph), 138.34 (s, triazole C-4'), 158.00 and 159.40 (2s, 2 x COCH<sub>2</sub>-CH<sub>3</sub>),  $\delta$  70.490.

Anal. Calcd for C<sub>19</sub>H<sub>24</sub>N<sub>3</sub>O<sub>6</sub>P (421.4): C, 54.15; H, 5.74; N, 9.97, P, 7.35. Found: C, 54.02; H, 5.66; N, 5.87; P, 9.83.

The same procedure gave triazole derivatives of phosphanyl sugar nucleosides 6A and 7A.

For (trimethylsilyl)acetylene adduct 6Aa: <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  0.12 (s, 9H, TMS), 1.46 (s, 3H, Me), 2.3-3.3 (m, 4H, CH<sub>2</sub>CH<sub>2</sub>), 5.31 (d, 1H, <sup>2</sup>J<sub>HP</sub>=9.48 Hz, CH), 6.36 (brs, 1H, OH), 7.2-7.7 (m, 5H, Ph), 7.34 (s, 1H, triazole H); <sup>13</sup>C NMR (CDCl<sub>3</sub>)  $\delta$  0.00 (s, TMS), 23.98 (d, J=6.68 Hz, Me), 25.24 (d, J=62.83 Hz, C-5), 37.98 (d, J=4.00 Hz, C-4), 68.93 (d, J=68.84 Hz, C-2), 79.73 (d, J=16.71 Hz, C-3), 127.88 (d, J=92.89 Hz, x-Ph), 128.00 (d, J=11.86 Hz, m-Ph), 130.85 and 145.74 (2s, triazole C-4' and C-5'), 130.93 (d, J=10.01 Hz, o-Ph), 132.37 (d, J=2.67 Hz, p-Ph); <sup>31</sup>P NMR (CDCl<sub>3</sub>)  $\delta$ 68.257.

Anal. Calcd for C<sub>16</sub>H<sub>24</sub>N<sub>3</sub>O<sub>2</sub>PSi (349.4): C, 54.99; H, 6.92; N, 12.02. Found: C, 54.69; H, 6.86; N, 11.90.

For methyl propiolate adducts 6Ab and 7Ab: <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  1.31 and 1.53 (2s, 6H, 2 x Me), 2.2-3.1 (m, 8H, CH<sub>2</sub>CH<sub>2</sub>), 3.76 and 3.83 (2s, 3H, COOMe for 6Ab and 7Ab), 4.93 (brs, 2H, 2x OH), 5.44 and 6.23 (2d, 1H, *J*=90.3 and 9.12 Hz, CH for 6Ab and 7Ab), 7.2-7.9 (m, 5H, Ph), 7.71 and 8.30 (2s, 1H, triazole H for 6Ab and 7Ab).

Anal. Calcd for C<sub>15</sub>H<sub>18</sub>N<sub>3</sub>O<sub>4</sub>P (335.3): C, 53.73; H, 5.41; N, 12.53. Found: C, 53.52; H, 5.32; N, 12.38.

For dimethyl acetylenedicarboxylate adduct 6Ac: <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  1.33 (s, 3H, Me), 2.5-3.1 (m, 4H, CH<sub>2</sub>CH<sub>2</sub>), 3.89 and 3.98 (2s, 5H, 2 x COOMe), 5.64 (d, 1H,

J=10.56 Hz, CH), 6.48 (brs, 1H, PH), 7.2-7.9 (m, 5H, Ph); <sup>13</sup>C NMR (CDCl<sub>3</sub>)  $\delta$  23.40 (d, J=60.3 Hz, Me), 24.98 (d, J=64.83 Hz, C-5), 37.63 (d, J=3.34 Hz, C-4), 52.42 and 53.61 (2s, 2 x COOMe), 68.12 (d, J=66.17 Hz, C-2), 79.86 (d, J=16.04 Hz, C-3), 127.78 (d, J=93.56 Hz, x-Ph), 128.06 (d, J=11.36 Hz, m-Ph), 131.08 (d, J=10.04 Hz, o-Ph), 131.72 (d, J=1.32 Hz, triazole C-5'), 132.60 (d, J=3.34 Hz, p-Ph), 138.19 (s, triazole C-4'), 158.30 and 159.73 (2s, 2 x COOMe); <sup>31</sup>P NMR (CDCl<sub>3</sub>)  $\delta$  69.908.

Anal. Calcd for C<sub>17</sub>H<sub>20</sub>N<sub>3</sub>O<sub>6</sub>P (390.3): C, 52.31; H, 4.39; N, 10.77. Found: C, 52.18; H, 4.26; N, 10.69.

For propargyl alcohol adduct 6Ae: <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  1.26 (s, 3H, Me), 2.2-3.1 (m, 4H, CH<sub>2</sub>CH<sub>2</sub>), 4.40 (s, 2H, CH<sub>2</sub>OH), 50.1 (d, 1H, J=8.95 Hz, CH), 7.2-7.8 (m, 5H, Ph), 7.86 (s, 1H, triazole H).

Anal. Calcd for  $C_{14}H_{18}N_3O_3P$  (307.3): C, 54.72; H, 5.90; N, 13.67. Found: C, 54.44; H, 5.65; N, 13.46.

For propargyl alcohol adduct 7Ae: <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  1.27 (s, 3H, Me), 2.2-3.1 (m, 4H, CH<sub>2</sub>CH<sub>2</sub>), 3.57 (s, 2H, CH<sub>2</sub>OH), 5.03 (d, 1H, J=8.95 Hz, CH), 7.2-7.8 (m, 5H, Ph), 7.36 (s, 1H, triazole H).

Anal. Calcd for C<sub>14</sub>H<sub>18</sub>N<sub>3</sub>O<sub>3</sub>P (307.3): C, 54.72; H, 5.90; N, 13.67. Found: C, 54.53; H, 5.71; N, 13.55.

For 3-methyl-1-butyn-3-ol adduct 6A f: <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  1.28 (s, 3H, Me), 1.37 (s, 6H, 2 x C<u>Me<sub>2</sub></u>OH), 20.-3.2 (m, 4H, CH<sub>2</sub>CH<sub>2</sub>), 5024 (d, 1H, *J*=90.3 Hz, CH), 7.3-7.8 (m, 5H, Ph), 7.44 (s, 1H, triazole H); <sup>31</sup>P NMR (CDCl<sub>3</sub>)  $\delta$  70.782.

Anal. Calcd for C<sub>16</sub>H<sub>22</sub>N<sub>3</sub>O<sub>3</sub>P (335.3): C, 57.31; H, 6.61; N, 12.53. Found: C, 57.19; H, 6.56; N, 12.41.

For 3-methyl-1-butyn-3-ol adduct 7A f: <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  1.39 (s, 3H, Me), 1.46 and 1.57 (2s, 6H, 2 x CMe<sub>2</sub>OH), 2.2-3.3 (m, 4H, CH<sub>2</sub>CH<sub>2</sub>), 5.85 (d, 1H, *J*=8.95 Hz, CH), 7.05 (s, 1H, triazole H), 7.2-7.8 (m, 5H, Ph); <sup>31</sup>P NMR (CDCl<sub>3</sub>)  $\delta$  72.918.

Anal. Calcd for C<sub>16</sub>H<sub>22</sub>N<sub>3</sub>O<sub>3</sub>P (335.3): C, 57.31; H, 6.61; N, 12.53. Found: C, 57.04; H, 6.48; N, 12.34.

For 2-butyn-1,4-diol adduct 6A g: <sup>1</sup>H NMR (CD<sub>3</sub>OD)  $\delta$  1.20 (s, 3H, Me), 2.2-3.3 (m, 4H, CH<sub>2</sub>CH<sub>2</sub>), 4.42 and 4.67 (2s, 4H, 2 x CH<sub>2</sub>OH), 5.11 (d, 1H, J=11.45 Hz, CH), 7.2-7.8 (m, 5H, Ph); <sup>13</sup>C NMR (CD<sub>3</sub>OD)  $\delta$  23.89 (d, J=6.00 Hz, Me), 25.24 (d, J=66.15 Hz, c-5), 39.17 (d, J=4.68 Hz, C-4), 51.98 and 55.33 (2s, 2 x CH<sub>2</sub>OH), 68.68 (d, J=70.18 Hz, C-2), 80.83 (d, J=17.38 Hz, C-3), 129.55 (d, J=93.56 Hz, x-Ph), 132.30 (d, J=9.34 Hz, o-Ph), 133.67 (d, J=2.67 Hz, p-Ph), 137.03 (d, J=1.99 Hz, triazole C-5'), 144.92 (s, triazole C-4').

Anal. Calcd for C<sub>15</sub>H<sub>20</sub>N<sub>3</sub>O<sub>4</sub>P (337.3): C, 53.41; H, 5.98; N, 12.46. Found: C, 53.22; H, 5.74; N, 12.32.

For acetylenedicarboxylic acid adduct 6Ah: <sup>1</sup>H NMR (CD<sub>3</sub>OD)  $\delta$  1.32 (s, 3H, Me), 2.2-3.1 (m, 4H, CH<sub>2</sub>CH<sub>2</sub>), 5.76 (d, 1H, *J*=11.36 Hz, CH), 7.2-7.8 (m, 5H, Ph); <sup>13</sup>C NMR (CD<sub>3</sub>OD)  $\delta$  23.91 (d, *J*=6.68 Hz, Me), 25.43 (d, *J*=68.16 Hz, C-5), 39.30 (d, *J*=4.01 Hz, C-4), 70.15 (d, *J*=68.19 Hz, C-2), 80.80 (d, *J*=16.03 Hz, C-3), 129.42 (d, *J*=11.36 Hz, *m*-Ph), 132.14 (d, *J*=10.03 Hz, *o*-Ph), 133.91 (d, *J*=2.67 Hz, *p*-Ph).

Anal. Calcd for C<sub>15</sub>H<sub>16</sub>N<sub>3</sub>O<sub>6</sub>P (365.3): C, 49.32; H, 4.41; N, 11.50. Found: C, 49.11; H, 4.36; N, 11.42.

2-Amino-3-hydroxy-3-methyl-1-phenylphospholane 1-oxide threo Catalytic hydrogenolysis of threo 2-azido-3-hydroxy-3-methyl-1-phenyl-(3Ab). phospholane 1-oxide (threo 3Aa, 0.896 g, 3.57 mmol) in methanol (10 mL) in the presence of 10% Pd/C at atmospheric pressure of hydrogen followed by filtration of the catalyst, removal of the solvent under reduced pressure, and column chromatography on silica gel gave the N-glycoside of phosphanyl sugar, 3Ab (0.795 g, 3.53 mmol, recrystallized from benzene) in 99% yield; mp 163-164 °C. <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  1.34 (s, 3H, Me), 1.8-2.8 (m, 4H, CH<sub>2</sub>CH<sub>3</sub>), 3.39 (d, 1H,  ${}^{2}J_{\mu\nu}$ =5.72 Hz, CH), 7.3-7.9 (m, 5H, Ph); <sup>13</sup>C NMR (CDCl<sub>3</sub>)  $\delta$  23.39 (d, <sup>3</sup> $J_{cp}$ =8.02 Hz, Me), 25.97 (d,  $J_{cp}$ =62.16 Hz, C-5), 35.77 (d,  ${}^{2}J_{cp}$ =5.33 Hz, C-4), 60.58 (d,  $J_{cp}$ =75.53 Hz, C-2), 79.04 (d,  ${}^{2}J_{cp}$ =11.36 Hz, C-3), 127.84 (d,  $J_{CP}$ =93.56 Hz, x-Ph), 128.30 (d,  ${}^{3}J_{CP}$ =11.35 Hz, m-Ph), 131.58 (d,  ${}^{3}J_{CP}$ =8.69 Hz, o-Ph), 132.08 (d,  ${}^{4}J_{cp}$ =2.67 Hz, p-Ph);  ${}^{31}$ P NMR (CDCl<sub>3</sub>)  $\delta$  68.257; IR  $\nu$  (KBr) 3040-3600 (br, OH, NH).

Anal. Calcd for C<sub>11</sub>H<sub>15</sub>NO<sub>2</sub>P (224.2): C, 58.92; H, 6.74; N, 6.25; P, 13.81. Found: C, 58.78; H, 6.14; N, 6.13; P, 13.98.

### ACKNOWLEDGMENTS

This work was partially supported by a Grant-in-Aid for Scientific Research (No. 04304049) from the Japanese Ministry of Education, Science, Sports and Culture, by Shizuoka Science and Technology Foundation, by Nippon Soda Co. Ltd, and by Suzuroku Dyeing Co. Ltd. The authors also thank Professors H. Yamamoto and T. Hanaya of Okayama University for 500 MHz NMR analyses, Professor J. Kobayashi of Shizuoka Institute of Technology for 400 MHz NMR measurement, Professor A. Kakehi of Shinshu University for X-ray analyses (Rigaku AFC5S), and Sankyo Co. Ltd. for microanalyses.

#### **REFERENCES AND NOTES**

- 1. Presented at the XIXth International Carbohydrate Symposium, San Diego, CA, USA, August 9-14, 1998.
- R. Goto, S. Inokawa, A. Sera, and S. Otani in *Tantorui no Kagaku (Chemistry of Monosaccharides)*, Maruzen, Tokyo, 1988.
- a) M. Yamashita, Y. Nakatsukasa, M. Yoshikane, H. Yoshida, T. Ogata, and S. Inokawa, *Carbohydr. Res.*, **59**, C12 (1977); b) M. Yamashita, Y. Nakatsukasa, H. Yoshida, T. Ogata, S. Inokawa, K. Hirotsu, and J. Clardy, *ibid.*, **70**, 247 (1979);
  c) M. Yamada and M. Yamashita, *ibid.*, **95**, C9 (1981); d) M. Yamashita, M. Yamada, K. Tsunekawa, T. Oshikawa, K. Seo, and S. Inokawa, *ibid.*, **121**, C4 (1983); *ibid.*, **122**, C1 (1983); e) M. Yamashita, M. Yamada, M. Sugiura, H. Nomoto, and T. Oshikawa, *Nippon Kagaku Kaishi*, 1207 (1987); f) H. Yamamoto and S. Inokawa, *Adv. Carbohydr. Chem. Biochem.*, **42**, 135 (1984).
- M. Yamashita, M. Uchimura, A. Iida, L. Parkanayi, and J. Clardy, J. Chem. Soc., Chem. Commun., 569 (1988).
- M. Yamashita, A. Yabui, K. Suzuki, Y. Kato, M. Uchimura, A. Iida, H. Mizuno, K. Ikai, T. Oshikawa, L. Parkanayi, and J. Clardy, J. Carbohydr. Chem., 16, 499 (1997).
- 6. C. Kibayashi and M. Akiba in Raifu Saiensu no Yuki Kagaku (Organic Chemistry for Life Science), Sankyo Shuppan, Tokyo, 1988, p 168.
- A. F. Bochkov and G. E. Zaikov in *Chemistry of the O-Glycosidic Bond*, Pergamon Press, Oxford, 1979, Chapter 2.
- J. T. Witkowski, R. K. Robins, R. W. Sidwell, and L. N. Simon, J. Med. Chem., 15, 1150 (1972).
- a) H. Mitsuya, K. J. Weinhold, P. A. Furman, M. H. St. Clair, S. N. Lehrman, R. L. Gallo, D. Bolognesi, D. W. Barry, and S. Broder, *Proc. Natl. Acad. Sci., USA*, 82, 7096 (1985); b) R. Z. Sterzycki, I. Gazzouli, V. Brankovan, J. C. Martin, and M. W. Mansuri, *J. Med. Chem.*, 33, 2150 (1990).
- 10. S. S. Cohen, Prog. Nucleic Acids Res. Mol. Biol., 5, 22 (1966).
- 11. J. J. Jaffe, Ann. N. Y. Acad. Sci., 255, 306 (1975).
- a) R. Vince and M. Hua, J. Med. Chem., 33, 17 (1990); b) M. Yoshikawa, N. Murakami, Y. Inoue, S. Hatakeyama, and I. Kitagawa, Chem. Pharm. Bull., 41, 636 (1993); c) D. W. Norbeck, S. Spanton, S. Broder, and H. Mitsuya, Tetrahedron Lett., 30, 6263 (1989); d) M. R. Dyson, P. L. Coe, and P. L. Walker, J. Med. Chem., 34, 2782 (1991).
- 13. L. D. Quin, J. P. Gratz, and T. P. Barket, J. Org. Chem., 33, 1034 (1968).
- 14. K. Ikai, A. Iida, and M. Yamashita, Synthesis, 595 (1989).
- 15. R. U. Lemieux and D. R. Lineback, Can. J. Chem., 43, 94 (1965).
- K. Nishizawa and J. Yoshimura in *Tansui Kabutsu (Carbohydrates)*, Asakura Shoten, Tokyo, 1980, Chapter 4.
- 17. F. Micheel and A. Klearer, Adv. Carbohydr. Chem., 16, 85 (1961).
- L. Hough and A. C. Richardson, in *Rodd's Chemistry of Carbon Compounds*, Vol IF; S. Coffey Ed.; Elsevier, Amsterdam, 1967, Chapter 23.
- Y. Kita, F. Itoh, O. Tamura, K. Y. Yaah, and Y. Tamura, *Tetrahedron Lett.*, 1431 (1987).
- S. J. Danishefsky, M. P. Deninno, and S. Chen, J. Am. Chem. Soc., 110, 3929 (1988).

- 21. C. E. Ballou, Adv. Carbohydr. Chem., 9, 59 (1954).
- 22. M. P. Bardolph and G. H. Coleman, J. Org. Chem., 15, 169 (1950).
- 23. A. Dyfverman and B. Lindberg, Acta Chem. Scand., 4, 878 (1950).
- 24. X-ray data for measurement and analysis of structure of 2A:

Rigaku AFC5S X-ray diffractometer with four-axis goniometer was used. Crystal data: Empirical formula, C<sub>11</sub>H<sub>14</sub>O<sub>2</sub>BrP; Crystal color, Habit, colorless, prismatic; Crystal dimensions, 1.00 x 0.90 x 1.00 mm; Crystal system, monoclinic; Lattice type, primitive; No. of reflections used for unit cell determination (2 $\theta$ range), 25 (29.94-34.76 °); Omega scan peak width at half-height, 0.24°; Lattice parameters, a=15.02 Å, b=10.134 Å, c=20.28 Å,  $\alpha = 90.00^{\circ}$ ,  $\beta = 107.3^{\circ}$ ,  $\gamma =$ 90.00 °, V=2948.72 Å<sup>3</sup>; Space group, P2<sub>1</sub>/a (#14); Z value, 4; D<sub>cale</sub>, 1.571 g/cm<sup>3</sup>;  $F_{000}$ , 1400;  $\mu$  (MoK  $\alpha$  ), 31.29 cm<sup>-1</sup>. Intensity measurments: Diffractometer, Rigaku AFC5S; Radiation, MoK  $\alpha$  ( $\lambda$  =0.71069Å), graphite monochromated; Attenuator, Ni foil (factors=2.3, 5.2, 11.7); Take-off angle, 6.0°; Detector aperture, 6.0 mm horizontal, 6.0 mm vertical; Diameter of beam collimator, 0.5 mm; Crystal to detector distance 400 mm; Temperature 22 °C; Scan type,  $\omega$ -2  $\theta$ ; Scan rate, 32.0° /min (in  $\omega$ )-up to 2 rescans; Scan width, (1.00 +0.30 tan  $\theta$ )°;  $2\theta_{\text{min}}$ , 55.0°; No. of reflections measured, total 6150, unique 5862 corrections, Lorentz-polarization. Structure solution and refinement: Structure solution, Direct methods (TEXSAN); Refinement, full-matrix least-squares; Function minimized,  $\Sigma$  w(IFol-IFcl)<sup>2</sup>; Least squares weights,  $1/\sigma^2$ (Fo)=4Fo<sup>2</sup>/ $\sigma^2$ (Fo<sup>2</sup>); p-factor, 0.03; Anomalous dispersion, all non-hydrogen atoms; No. observations (I > 3.00  $\sigma$  (I)), 1426; No. variables, 307; Reflection/Parameter ratio, 4.64; Residuals: R; Rw, 0.083; 0.090; Goodness of fit indicator, 2.82; Max shift/Error in final cycle, 0.39; Maximum peak in final diff. map, 1.19  $e/A^3$ ; Minimum peak in final diff. map, -0.73 e7Å<sup>3</sup>.

X-ray structure data for 2A.

Selected bond distances in Å: P(1)-C(2) 1.86, P(1)-C(5) 1.80, C(2)-C(3) 1.54, C(3)-C(4) 1.55, C(4)-C(5) 1.47.

Selected bond angles in degree: C(2)-P(1)-C(5) 94, P(1)-C(2)-C(3) 103, C(2)-C(3)-C(4) 104, C(3)-C(4)-C(5) 108, P(1)-C(5)-C(4) 108.

Selected torsional angles in degree: P(1)-C(2)-C(3)-C(4) 43, P(1)-C(5)-C(4)-C(3) 31, C(2)-P(1)-C(5)-C(4) -4, C(2)-C(3)-C(4)-C(5) -49, C(3)-C(2)-P(1)-C(5) -24.

These data imply that the conformation of 2A may be in a <sup>2</sup>E form.

25. X-ray data for measurement and analysis of structure of 3Aa:

Rigaku AFC5S X-ray diffractometer with four-axis goniometer was used. Crystal data: Empirical formula,  $C_{11}H_{14}N_3O_2P$ ; Crystal color, Habit, colorless, prismatic; Crystal dimensions, 0.16 x 0.82 x 0.98 mm; Crystal system, monoclinic; Lattice type, primitive; No. of reflections used for unit cell determination (2  $\theta$ range), 23 (20.1-29.7°); Omega scan peak width at half-height, 0.21°; Lattice parameters, a=25.18 Å, b=9.834 Å, c=14.415 Å,  $\alpha$  =90.00 °,  $\beta$  =122.23 °,  $\gamma$  = 90.00°, V=3020Å<sup>3</sup>; Space group, C2/c (#15); Z value, 8; D<sub>cale</sub>, 1.630 g/cm<sup>3</sup>;  $\mu$  (MoK  $\alpha$ ), 7.19 cm<sup>-1</sup>. Intensity measurments: Diffractometer, F<sub>000</sub>, 1520; Radiation, MoK  $\alpha$  ( $\lambda = 0.71069$  Å), graphite monochromated; Rigaku AFC5S; Attenuator, Ni foil (factors=2.3, 5.2, 11.7); Take-off angle, 6.0°; Detector aperture, 6.0 mm horizontal, 6.0 mm vertical; Diameter of beam collimator, 0.5 mm; Crystal to detector distance 400 mm; Temperature 22 °C; Scan type,  $\omega$ -2 $\theta$ ; Scan rate, 32.0° /min (in  $\omega$ )-up to 2 rescans; Scan width, (1.15 + 0.30 tan  $\theta$ )°;

 $2\theta_{max}$ , 55.0°; No. of reflections measured, total 3768, unique 3684 corrections, Lorentz-polarization. Structure solution and refinement: Structure solution, Direct methods (TEXSAN); Refinement, full-matrix least-squares; Function minimized,  $\Sigma w(IFol-IFcl)^2$ ; Least squares weights,  $1/\sigma^2(Fo)=4Fo^2/\sigma^2(Fo^2)$ ; p-factor, 0.03; Anomalous dispersion, all non-hydrogen atoms; No. observations (I > 3.00  $\sigma$  (I)), 1100; No. variables, 163; Reflection/Parameter ratio, 6.75; Residuals: R; Rw, 0.215; 0.301; Goodness of fit indicator, 7.53; Max shift/Error in final cycle, 1.35; Maximum peak in final diff. map, 6.54 e/Å<sup>3</sup>; Minimum peak in final diff. map, -0.88 e/Å<sup>3</sup>.

X-ray structure data for 3Aa.

Selected bond distances in Å: P(1)-C(1) 1.85(3), C(1)-C(2) 1.56(5), C(2)-C(3) 1.54(4), C(3)-C(4) 1.43(5), C(4)-P(1) 1.84(4).

Selected bond angles in degree: P(1)-C(1)-C(2) 101(2), C(1)-C(2)-C(3) 105(3), C(2)-C(3)-C(4) 111(3), C(3)-C(4)-P(1) 105(3), C(4)-P(1)-C(1) 96(2).

Selected torsional angles in degree: P(1)-C(2)-C(3)-C(4) -43(3), C(1)-C(2)-C(3)-C(4) -51(4), C(2)-C(3)-C(4)-P(1) -30(4), C(3)-C(4)-P(1)-C(1) -3(3), C(4)-P(1)-C(1)-C(2) -24(2).

These data imply that the conformation of 3Aa may be in a <sup>2</sup>E form.

- 26. J. Davoll and B. A. Lowy, J. Am. Chem. Soc., 53, 1650 (1951).
- 27. N. Yamaoka, K. Aso, and K. Matsuda, J. Org. Chem., 30, 145 (1965).
- 28. G. A. Howard, B. Lythgoe, and A. R. Todd, J. Chem. Soc., 1052 (1947).
- 29. M. J. Robins and R. K. Robins, J. Am. Chem. Soc., 67, 4934 (1965).
- a) U. Niedballa and H. Vorbruggen, J. Org. Chem., 39, 3654 (1974); b) K. Augustyns, J. Rozenski, A. V. Aerschoy, G. Janssen, and P. Herdewijin, *ibid.*, 58, 2977 (1993).
- a) G. Shaw, R. N. Warrener, M. H. Maguire, and R. K. Ralph, J. Chem. Soc., 2294 (1958);
  b) Y. F. Shealy and C. A. O'Dell, J. Heterocycl. Chem., 13, 1041 (1976).
- 32. I. Goodman, Federation Proceedings, Fed. Am. Soc. Exp. Biol., 15, 264 (1956).
- 33. D. J. Hlasta and J. H. Ackerman, J. Org. Chem., 59, 6184 (1994).
- 34. M. Yamashita, Y. Kato, K. Suzuki, and T. Oshikawa, *Heterocyclic Commun.*, 4, 411 (1998).
- 35. X-ray data for measurement and analysis of structure of 6d.

Rigaku AFC7R X-ray diffractometer with four-axis goniometer was used. Crystal data: Empirical formula,  $C_{19}H_{24}N_3O_6P$ ; Formula weight, 421.39; Crystal color, Habit, colorless, prismatic; Crystal dimensions, 020x0.20x0.30 mm; Crystal system, monoclinic; Lattice type, primitive; No. of reflections used for unit cell determination (2  $\theta$  range), 25 (54.5-57.0°); Omega scan peak width at half-height, 0.22°; Lattice parameters, a=14.243 Å, b=10.494 Å, c=14.486 Å,  $\beta$ =98.138°, V=1589.6 Å<sup>3</sup>; Space group, P2<sub>1</sub>/n (#9); Z value, 4; D<sub>cale</sub>, 1.306 g/cm<sup>3</sup>; F<sub>000</sub>, 888.00;  $\mu$  (CuK  $\alpha$ ), 14.86 cm<sup>-1</sup>. Intensity measurments: Diffractometer, Rigaku AFC7R; Radiation, CuK  $\alpha$  ( $\lambda$  =1.54178 Å), graphite monochromated; Attenuator, Ni foil (factors=1.00, 9.48, 9.48, 9.43); Take-off angle, 6.0°; Detector aperture, 9.0 mm horizontal, 13.0 mm vertical; Crystal to detector distance 235 mm; Temperature 20.0°C; Scan type,  $\omega$ -2 $\theta$ ; Scan rate, 16.0°/min (in  $\omega$ )-up to 3 scans; Scan width, (1.78 +0.30 tan  $\theta$ )°; 2 $\theta_{max}$ , 120.1°; No. of reflections measured, total 3537; Corrections, Lorentz-polarization. Structure solution and refinement: Structure solution, Direct methods (SHELXS86); Refinement, full-matrix least-squares; Function minimized,  $\Sigma w(|\text{Fol-IFcl})^2$ ; Least squares weights,  $1/\sigma^2(\text{Fo})=4\text{Fo}^2/\sigma^2(\text{Fo}^2)$ ; p-factor, 0.00; Anomalous dispersion, all non-hydrogen atoms; No. observations (I > 3.00  $\sigma$  (I)), 2337; No. variables, 358; Reflection/Parameter ratio, 6.53; Residuals: R; Rw, 0.056; 0.043; Goodness of fit indicator, 3.19; Max shift/Error in final cycle, 2.73; Maximum peak in final diff. map, 0.36 e'/Å<sup>3</sup>; Minimum peak in final diff. map, -0.34 e'/Å<sup>3</sup>.

Tables of atomic coordinates, bond lengths, and bond angles have been deposited with the Cambridge Crystallographic Data Centre. These tables may be obtained on request from the Director, Cambridge Crystallographic Data Centre, 12 Union Road, Cambridge CB2 IEZ, UK.

- 36. M. Yamashita, M. Uchimura, A. Iida, L. Parkanayi, and J. Clardy, J. Chem. Soc., Chem. Commun., 569 (1988).
- 37. a) J. Emsley and D. Hall in *The Chemistry of Phosphorus*, Harper & Row, London, 1976, p 157; b) L. D. Quin and T. P. Barket, *J. Chem. Soc.*, *Chem. Commun.*, 788 (1967); c) W. B. McCormack, *Org. Syn.*, 43, 73 (1963).
- a) K. Hunger, U. Hasserodt, and F. Korte, *Tetrahedron*, 20, 1563 (1964); b) G. Kresze, J. Firl, H. Zimmer, and U. Wollnic, *Tetrahedron*, 20, 1593 (1964).
- 39. L. D. Quin, J. P. Gratz, and T. P. Barket, J. Org. Chem., 33, 1034 (1968).
- 40. a) L. D. Quin in 1,4-Cycloaddition Reactions: The Diels-Alder Reaction in Heterocyclic Syntheses, J. Hamer Ed.; Academic Press, New York, 1967, Chapter 3; b) L. D. Quin in The Heterocyclic Chemistry of Phosphorus, Wiley-Interscience, New York, 1981, Chapter 2; c) J. Emsley and D. Hall in The Chemistry of Phosphorus, Harper & Row Publishers, London, 1976, Chapter 4.