

Communication

Extension of Pd-Mediated One-Pot Ketone Synthesis to Macrocyclization: Application to a New Convergent Synthesis of Eribulin

Jung Hwa Lee, Zhanjie Li, Ayumi Osawa, and Yoshito Kishi

J. Am. Chem. Soc., Just Accepted Manuscript • DOI: 10.1021/jacs.6b11663 • Publication Date (Web): 08 Dec 2016

Downloaded from http://pubs.acs.org on December 8, 2016

Just Accepted

"Just Accepted" manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides "Just Accepted" as a free service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. "Just Accepted" manuscripts appear in full in PDF format accompanied by an HTML abstract. "Just Accepted" manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are accessible to all readers and citable by the Digital Object Identifier (DOI®). "Just Accepted" is an optional service offered to authors. Therefore, the "Just Accepted" Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the "Just Accepted" Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these "Just Accepted" manuscripts.



Journal of the American Chemical Society is published by the American Chemical Society. 1155 Sixteenth Street N.W., Washington, DC 20036 Published by American Chemical Society. Copyright © American Chemical Society. However, no copyright claim is made to original U.S. Government works, or works produced by employees of any Commonwealth realm Crown government in the course of their duties.

Extension of Pd-Mediated One-Pot Ketone Synthesis to Macrocyclization: Application to a New Convergent Synthesis of Eribulin

Jung Hwa Lee, Zhanjie Li,[†] Ayumi Osawa, and Yoshito Kishi*

Department of Chemistry and Chemical Biology, Harvard University, Cambridge, Massachusetts 02138

Supporting Information Placeholder

ABSTRACT: Recently reported Pd-mediated one-pot ketone synthesis from an unactivated alkyl bromide and a thioester has been extended to a macrocyclic ketone synthesis. In-situ generation of alkylzinc halide via single electron transfer (SET), using NbCpCl₄ and CrCl₃, was the key for the success of macrocyclization. A new convergent synthesis of eribulin has been achieved, using: (1) catalytic asymmetric Ni/Cr-mediated coupling to form the C19-C20 bond, (2) base-induced cyclization to form the methylenetetrahydrofuran ring, and (3) Pd-mediated one-pot ketone synthesis to form the macrocyclic ketone.

Halichondrins are polyether macrolides, originally isolated from the marine sponge Halichondria okadai by Uemura, Hirata, and co-workers.¹ This class of natural products displays interesting structure diversities on the oxidation state at C12 and C13, cf., halichondrin A-C in Scheme 1. We chose halichondrin B as a synthetic target and began the experimental work, which led us to the first total synthesis of halichondrin B in 1992. On completion of the synthesis,² we asked (the late) Dr. Suffness at the National Cancer Institute (NCI) and Dr. Littlefield at Eisai Research Institute (ERI) to test antitumor activities of the totally synthetic halichondrins, along with several synthetic intermediates. The results were sensational: their experiments clearly demonstrated that the antitumor activities of halichondrin B resided in the right portion of the molecule, which served as the foundation for successful development of the antitumor drug Halaven (eribulin) by Eisai.^{3,4,5}

Recently, we reported a unified, convergent synthesis of this class of marine natural products, using: (1) Ni/Crmediated coupling to form the C19-C20 bond, (2) THF S_N2 cyclization between C17-Cl and C20-OH, and (3) macrolactonization (Scheme 2).^{2b,6} We were interested in extending this synthetic strategy to the synthesis of eribulin, in which the first two key synthetic transformations could be achieved by use of the chemistry developed for the unified synthesis of halichondrins. The third key synthetic transformation is the cyclization to form the macrolactone in the halichondrin series, whereas it is the cyclization to form the macrocyclic ketone in the eribulin series. Macrolactonization is a wellprecedented synthetic transformation. On the contrary, except for a few limited cases,⁷ cyclization to form a macrocyclic ketone, referred to as macroketocyclization in this paper, is an unexploited synthetic transformation.⁸ In this communication, we report a macroketocyclization between a non-activated alkylbromide with a thioester and its application to a synthesis of eribulin.





Related to this work, we recently developed a Pd-mediated one-pot ketone synthesis from non-activated alkyl bromides and thioesters, with an intention of extending it to a macro-ketocyclization.⁹ In this connection, we should note that the efficiency of this method is excellent, even with use of a \sim 1:1 molar ratio of two coupling partners. However, in order to translate the one-pot ketone synthesis to a macroketocyclization, we need to address one additional question, that is how to eliminate, or suppress, an intermolecular coupling. For the case of macrolactonization, a high-dilution technique is commonly employed to achieve this goal.

Pd-Mediated ketone synthesis is generally considered to involve three distinct steps: (1) oxidative addition of a Pd(0)catalyst to a thioester to form RCO-Pd(II)X, (2) transmetallation from an alkylzinc halide to the resultant Pd(II) species, and (3) reductive elimination, leading to a ketone and regenerating the Pd(0)-catalyst. Among these steps, we speculated the second step (transmetallation) to be most critical to effectively achieve the macroketocyclization under a highdilution condition. Upon dilution, an *intra*-molecular transmetallation would be favored over the *inter*-molecular transmetallation, but would be disfavored over undesired sidereactions due to a higher probability of wasting radical and/or organometallic species.¹⁰

Experimentally, it was found that catalytic *inter*-molecular ketone synthesis proceeded well even at 25 mM. Among three conditions, Condition C [(Pd₂dba₃ (10 mol%), PCy₃ (20 mol%), CrCl₂ (0.5 equiv), NbCpCl₄ (10 mol%), LiI (1 equiv), TESCl (1.5 equiv), Zn (xs) in DMI)] gave the best conversion.^{11,12}

Scheme 2. Three Key Transformations employed in the Unified Convergent Synthesis of Halichondrins

Halichondrin-B Series



Being encouraged with this observation, we chose substrate 4a to study the feasibility of macroketocyclization (Table 1). In this study, 4a was subjected to a specified condition, and a yield of 5a was estimated from a ¹H-NMR analysis of crude product.¹² At 50 mM concentration, which was effective for inter-molecular ketone synthesis (vide ante), 4a gave the debrominated product and dimer as major products (entry 1). Considering that the activity of reagents might diminish by dilution, we then tested the macroketocyclization in the presence of a stoichiometric amount of metals, thereby demonstrating that the desired ketone 5a was indeed formed as a major product at 10 mM concentration with only a small amount of debrominated product, although the dimer was still detected in more than 10% (entry 2). Under the stoichiometric conditions, $CrCl_2$ and $NbCpCl_4$ were essential (entries 4 & 5), but LiI and TESCl were not (entry 3). Also, reducing the amount of NbCpCl₄ resulted in a lower yield (entry 6). These observations implicate that SET activation and the early transition metals (TM) are critical for macroketocyclization. Interestingly, this coupling condition corresponds to Condition C for *inter*-molecular one-pot ketone synthesis. At present, we do not have experimental supports to suggest a specific role(s) of early transition metals. However, we would speculate that both metals play the same role(s) in both intra- and intermolecular couplings.¹³ Lastly, it was found that Cr(III)Cl₃ was more effective than Cr(II)Cl₂ to lower the concentration further (entries 9 - 11). Under these conditions, the desired product was formed, accompanied with only a trace amount of debrominated product.¹²

Table 1. Macroketocyclization: Optimization^a



^aConditions: To Pd₂dba₃ (0.04 mmol) and PCy₃ (0.08 mmol) in DMI (2 mL) were added Zn (0) (xs), CrCl₃ (0.2 mmol), and NbCpCl₄ (0.04 mmol) at rt in a glove box. Then, if needed, LiI (0.4 mmol) and TESCI (0.06 mmol) were added to the reaction mixture followed by S.M. in THF (2 mL). ^bRoughly estimated yield based on a ratio of 5a to side products (debrominated RH and dimer) in a crude ¹H NMR.¹² ^cPd₂dba₃ (0.1 equiv), PCy₃ (0.2 equiv) used. ^dRH was a major product. ^eReduction of Zn (20-40 equiv) provided slightly lower yield. ^fTrace amount of RH. ^gLower yield mainly due to dimer. ^hLower yield mainly due to debromination, yet dimer-formation was not noticeably reduced. Abbreviation: DMI = 1,3-dimethyl-2-imidazolidinone; $NbCpCl_4$ tetrachloro(cyclopeantadienyl)niobium; Pd₂dba₃ tris(dibenzyli-deneacetone)dipalladium(0); $PCy_3 = tricyclohex$ ylphosphine.

We then carried out the macroketocyclization of 4a under the condition of entry 10 in a preparative scale (0.2 mmol) (Scheme 3). In order to achieve the macroketocyclization effectively, it has become evident that two conditions must be met: (1) to maintain Pd-, Nb- and Cr-reagents in a stoichiometric amount and (2) to maintain the substrate-concentration above 5 mmol. From a practical point of view, it would be more attractive if an amount of Pd-, Nb- and Cr-reagents could be reduced.¹⁴ To address this issue, we tested the possibility of recycling the reagents mixture in one-pot. Specifically, one-half of substrate 4a was added into one-half amount of the reagent mixture used for the stoichiometric conditions and, after 7 hours, the remaining half of 4a was added to the same reaction mixture. Under this setting, the cyclization completed to give 5a in 58% isolated yield after chromatographic purification. Thus, the macroketocyclization was effective with use of one-half of the reagents mixture in the price of time, i.e., 7 vs. 14 hours. Similarly, the macroketocyclization was tested by adding 1/4- and 1/8-amounts of 4a into a 1/4- and 1/8-amount of the reagents mixture, respectively, every 7 hours, to give 5a in 55%. Overall, under these conditions, the cyclization was achieved with use of ~30% and $\sim 15\%$ of the reagents mixture, in the price of time, i.e., 7 hours vs. 28 and 56 hours.¹⁵ This procedure was also found effective for 16-membered ketone 5b to give 57% yield from **4b**. In both cases, dimers and debrominated products were detected, but only in insignificant amounts (<10%).





^{*a*}Conditions: To Pd₂dba₃ (0.1 mmol) and PCy₃ (0.2 mmol) in DMI/THF (10 mL/5 mL) were added CrCl₃ (0.5 mmol), Zn (xs), and NbCpCl₄ (0.5 mmol) at rt in a glove box. SM (0.2 mmol) was added to the reagent mixture in two portions. First, one half (0.1 mmol) of SM in THF (2.5 mL) was added to the reagent mixture and stirred at rt. After 7 h, the remaining half (0.1 mmol) of SM in THF (2.5 mL) was added and stirred overnight. ^{*b*}See Supporting Information for details.

Having demonstrated the feasibility of one-pot macroketocyclization, we shifted our focus onto its application to a synthesis of eribulin (Scheme 2). The synthesis of aldehyde 1 was started from the known sulfone 6.¹⁶ Protecting group manipulation, hydroxylation of sulfone 7 to alcohol $\mathbf{8}$, ¹⁶ followed by tosylation and bromide substitution proceeded uneventfully (Scheme 4). However, deprotection of 4methoxytrityl (MMTr) ether required an optimization, because of a concomitant deprotection of the primary TBS group; 1,1,1,3,3,3-hexafluoro-2-propanol (HFIP), which was known effective for selective deprotection of 4,4dimethoxytrityl (DMTr),¹⁷ resulted in only partial deprotection of MMTr at 40 °C. Assuming that a MMTr-cation acceptor might enhance the deprotection, we eventually found that an addition of water (HFIP/H₂O = 40/1) allowed us selectively to complete the required deprotection at rt. Then, the resulting alcohol was oxidized to aldehyde 1. On the other hand, thioester 2 was straightforwardly prepared from the known methyl ester 9 in two-steps:⁶ hydrolysis by Me₃SnOH¹⁸ and coupling with EtSH by DCC.

With both aldehyde 1 and vinyl iodide 2 in hand, we studied the C19-20 Ni/Cr-mediated coupling. Initially, the condition optimized for the synthesis of halichondrin A was applied for coupling of 1 and 2 with Ni-complex I,^{2b} but gave the desired product 10 only in a modest yield (~40%). We speculated that the low yield might be attributed to a poor selectivity in activation of the C19-vinyl iodide – note the presence of an alkyl bromide as well as a thioester, which might potentially be activated with low-valent Ni. With this speculation, we searched for a Ni-catalyst, which would allow us selectively to activate the C19-vinyl iodide and consequently improve the efficiency of Ni/Cr-mediated coupling of 2 with 1. Through this search, it was found that a combination of Ni-complex II, prepared from electron-rich 2,3,4,7,8,9-hexamethyl-1,10phenanthroline, and Cr-catalyst, prepared from unnat-i-Pr/Me/OMe sulfonamide I, gave a satisfactorily high coupling yield (86% yield; $dr = \sim 10.1 (^{1}H-NMR)$).¹⁹

Scheme 4. A New Convergent Synthesis of Eribulin



^aReagents and Conditions: a. 1. MMTrCl, *i*-Pr₂NEt, CH₂Cl₂, 93%. 2. K₂CO₃, MeOH. 3. TBSCl, imidazole, 88% for 2 steps. b. n-BuLi, THF, -78 °C; HBSia₂, -10 °C to rt, >12 h; H₂O₂, 3M NaOH, 0 °C, 81%. c. 1. TsCl, DMAP (cat.), Et₃N, CH₂Cl₂, 88%. NaBr, Bu₄NBr (cat.), acetone, reflux, 90%. 3. 2 $(CF_3)_2CHOH/H_2O = 40/1$, 3 h, 90%. 4. Dess-Martin Ox., 90%. d. 1. Me₃SnOH, 80 ~ 85 °C, DCE; 0.1 N HCl. 2. EtSH, DCC, DMAP, 94%. e. CrCl₂ (20 mol%), Cr-Ligand I (24 mol%), proton sponge (24 mol%), Ni-complex II (5 mol%), LiCl, Mn, ZrCp₂Cl₂, CH₃CN/EtOAc= 3/1 (0.15 M), 86%. f. SrCO₃ (xs), t-BuOH/H₂O = 20/1 (4 mM), 95°C, open to air, 87%. g. Pd₂dba₃ (1 equiv), PCyp₃ (2 equiv),²⁰ CrCl₃ (5 equiv), NbCpCl₄ (1 equiv), Zn (0) (xs), DMI/THF (1/1, 27 mM), 64%. h. see reference 24. Abbreviation: DCC = dicyclohexylcarbodiimide; $PCyp_3 = tricy$ clopentylphosphine.

The next task was to cyclize **10** to **11**, which had been done with AgOTf/Ag₂O in the synthesis of halichondrin A. Apparenly, this condition was not suitable to the substrate **10**, because of the presence of thioester- and bromide-groups. Thus, we tested the cyclization conditions reported by Britton (100 °C in water),²¹ which gave the desired product **11**, although accompanied with a large amount of unidentified decomposition products. We speculated that the liberated HCl might have caused the decomposition, and began an extensive search for a suitable base, leading us to a satisfactory condition: $SrCO_3(s)$ at 95 °C.²² Under the optimized conditions, **11** was isolated in 87% yield and fully characterized.

Finally, 11 was subjected to macroketocyclization under the stoichiometric conditions. It is worthwhile noting that, contrary to model compounds 4a,b, the major side-reaction in this series was the reductive quentching of -CH₂Br to CH₃ rather than the dimerization, thereby suggesting the possibility of using a higher concentration. We assume that the difference in behavior might be attributed to the difference in conformational property of 11, compared to 4; namely, 11 might have adopted a favorable conformation required for the macroketocyclization. Consistent with this assumption, the macroketocyclization was achieved, without noticeable dimerization, even at 27 mM concentration, to furnish ketone 3 in 64% yield (52 mg scale).²³ Spectroscopic comparisons (¹Hand ¹³C-NMR, HR-MS) firmly established that **3** thus obtained was identical with the authentic sample.3,5 Lastly, macrocyclic ketone **3** was converted into eribulin in three steps.²⁴

In summary, a method has been developed for macroketocyclization between an alkyl bromide and a thioester under mild conditions. NbCpCl₄ and CrCl₃ are key components not only for in-situ activation of alkyl bromide to alkylzinc halide via a SET process but also for acceleration of Pd-mediated coupling. Notably, this unique macroketocyclization does not require any special template or functional group to be removed after cyclization. Overall, the newly-developed macroketocyclization has allowed us to synthesize eribulin with the same synthetic strategy as the one used in the halichondrins.

ASSOCIATED CONTENT

Supporting Information

Additional data for Table 1, experimental procedures, characterization data, and copies of spectra. The Supporting Information is available free of charge on the ACS Publications website.

AUTHOR INFORMATION

Corresponding Author

*kishi@chemistry.harvard.edu

Present Addresses

†The Dow Chemical Company, 2200 W. Salzburg Road, Midland, MI 48686

Notes

No competing financial interests have been declared.

ACKNOWLEDGMENT

Financial support from the Eisai USA Foundation is gratefully acknowledged.

REFERENCES

¹ (a) Uemura, D.; Takahashi, K.; Yamamoto, T.; Katayama, C.; Tanaka, J.; Okumura, Y.; Hirata, Y. *J. Am. Chem. Soc.* **1985**, *107*, 4796. (b) Hirata, Y.; Uemura, D. *Pure Appl. Chem.* **1986**, *58*, 701. ² (a) Aicher, T. D.; Buszek, K. R.; Fang, F. G.; Forsyth, C. J.; Jung, S. H.; Kishi, Y.; Matelich, M. C.; Scola, P. M.; Spero, D. M.; Yoon, S. K. *J. Am. Chem. Soc.* **1992**, *114*, 3162. (b) Ueda, A.; Yamamoto, A.; Kato, D.; Kishi, Y. *J. Am. Chem. Soc.* **2014**, *136*, 5171 and references cited therein.

³ For discovery of eribulin, see: Zheng, W.; Seletsky, B. M.; Palme, M. H.; Lydon, P. J.; Singer, L. A.; Chase, C. E.; Lemelin, C. A.; Shen, Y.; Davis, H.; Tremblay, L.; Towle, M. J.; Salvato, K. A.; Wels, B. F.; Aalfs, K. K.; Kishi, Y.; Littlefield, B. A.; Yu, M. J. *Bioorg. Med. Chem. Lett.* **2004**, *14*, 5551.

⁴ For reviews on discovery of eribulin, see: (a) Yu, M. J.; M.; Kishi, Y.; Littlefield, B. A. *Anticancer Agents from Natural Products*; CRC Press: 2005, p 241. (b) Yu, M. J.; Zheng, W.; Seletsky, B. M.; Littlefield, B. A.; Kishi, Y. *Annu. Rep. Med. Chem.*; John, E. M., Ed.; Academic Press: 2011, Vol. 46, p 227.

⁵ For process development of eribulin, see: Austad, B. C.; Calkins, T. L.; Chase, C. E.; Fang, F. G.; Horstmann, T. E.; Hu, Y.; Lewis, B. M.; Niu, X.; Noland, T. A.; Orr, J. D.; Schnaderbeck, M. J.; Zhang, H.; Asakawa, N.; Asai, N.; Chiba, H.; Hasebe, T.; Hoshino, Y.; Ishizuka, H.; Kajima, T.; Kayano, A.; Komatsu, Y.; Kubota, M.; Kuroda, H.; Miyazawa, M.; Tagami, K.; Watanabe, T. *Synlett* **2013**, *24*, 333 and preceding two papers.

⁶ (a) ref. 2b. (b) Li, J.; Yan, W.; Kishi, Y. J. Am. Chem. Soc. **2015**, 137, 6226 and the preceding paper.

⁷ For selected examples for macroketocyclization, see: (a) Nishikawa, K.; Yoshimi, Y.; Maeda, K.; Morita, T.; Takahashi, I.; Itou, T.; Inagaki, S.; Hatanaka, M. J. Org. Chem. **2013**, 78, 582. (b) Tsuna, K.; Noguchi, N.; Nakada, M. Tetrahedron Lett. **2011**, 52, 7202. (c) Porter, N. A.; Chang, V. H. T.; Magnin, D. R.; Wright, B. T. J. Am. Chem. Soc. **1988**, 110, 3554. (d) Kong, K.; Romo, D.; Lee, C. Angew. Chem., Int. Ed. **2009**, 48, 7402. (e) Boger, D. L.; Mathvink, R. J. J. Am. Chem. Soc. **1990**, 112, 4008. (f) Kende, A. S.; Liu, K.; Kaldor, I.; Dorey, G.; Koch, K. J. Am. Chem. Soc. **1995**, 117, 8258. (g) Wu, J.; Panek, J. S. J. Org. Chem. **2011**, 76, 9900. (h) Lowe, J. T.; Panek, J. S. Org. Lett. **2008**, 10, 3813.

⁸ In the previous synthesis, the macrocyclization was achieved by an intramolecular Ni/Cr-mediated coupling at C13-C14, before construction of the C8-C14 polycycle: see, (a) ref. 3. (b) Namba, K.;

Kishi, Y. J. Am. Chem. Soc. **2005**, *127*, 15382, (c) ref. 5. (d) Inanaga, K.; Fukuyama, T.; Kubota, M.; Komatsu, Y.; Chiba, H.; Kayano, A.; Tagami, K. Org. Lett. **2015**, *17*, 3158.

Lee, J. H.; Kishi, Y. J. Am. Chem. Soc. 2016, 138, 7178.

¹⁰ We hoped to suppress undesired side-reactions by either slow activation of RX and/or acceleration of transmetallation to avoid accumulation of generated alkylzinc halides.

¹¹ The conversion was 58%, 71%, and 80% under Conditions A, B, and C, respectively.

¹² See Supporting Information for details.

¹³ For the case of intermolecular ketone coupling, we have suggested a possibility that the early transition metals might shift equilibrium from stable RZnX to higher-order orgnozincates and/or might break Pd-Zn to restore Pd reactivity, respectively.

¹⁴ There was one additional benefit in reducing an amount of the reagent, i.e., isolation of the product was much easier with a lesser amount of the reagent.

¹⁵ These experimental data should allow us to identify a proper setting for use of a syringe-pump.

¹⁶ Liu, L.; Henderson, J. A.; Yamamoto, A.; Brémond, P.; Kishi, Y. *Org. Lett.* **2012**, *14*, 2262.

¹⁷ Leonard, N. J.; Neelima Tetrahedron Lett. 1995, 36, 7833.

¹⁸ Nicolaou, K. C.; Estrada, A. A.; Zak, M.; Lee, S. H.; Safina, B. S. *Angew. Chem., Int. Ed.* **2005**, *44*, 1378.

¹⁹ With the Cr-catalyst derived from *unnat-i*-Pr/PhCl₂/OCy(Me)₂ sulfonamide, the dr observed for this Ni/Cr-mediated coupling was around 20:1 in the halichondrin series: see ref. 2b and Guo, H.; Dong,

60

CG.; Kim, DS.; Urabe, D.; Wang, J.; Kim, J. T.; Liu, X.; Sasaki, T.; Kishi, Y. J. Am. Chem. Soc. 2009 , 131, 15387. ²⁰ PCyp ₃ is better than PCy ₃ for this type of substrates, see ref. 9 ²¹ Kang, B.; Chang, S.; Decker, S.; Britton, R. Org. Lett. 2010 , 12, 1716. ²² Interestingly, soluble amine bases gave complicated side reactions such as halide exchange. Insert Table of Content $MeQ \bigoplus_{i=1}^{B} \bigoplus_{j=1}^{EIS} \bigoplus_{j=1}^{U} \bigoplus_$	 ²³ Reductive debromination was observed at 18 mM concentrat t not at 25 mM concentration. On the other hand, dimerization t observed even at 27 mM concentration. ²⁴ See (a) ref. 3. (b) ref. 5. (c) Kaburagi, Y.; Kishi, Y. <i>Tetrahe tt.</i> 2007, <i>48</i>, 8967. s artwork here
Insert Table of Content MeQ, Br EtS + O + H + H + H + H + H + H + H + H + H	s artwork here
MeQ, Br EtS, O H H H H H H H H H H H H H H H H H H H	
64% isolated	TBSO MeO H H H H H H H H H H H H H H H H H H H
64% isolated	F (27 mM) , yield