

Communication

Ni-Catalyzed Enantioselective Conjunctive Coupling with C(sp³) Electrophiles: A Radical-Ionic Mechanistic Dichotomy

Gabriel J. Lovinger, and James P. Morken

J. Am. Chem. Soc., **Just Accepted Manuscript** • DOI: 10.1021/jacs.7b10519 • Publication Date (Web): 08 Nov 2017

Downloaded from <http://pubs.acs.org> on November 8, 2017

Just Accepted

“Just Accepted” manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides “Just Accepted” as a free service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. “Just Accepted” manuscripts appear in full in PDF format accompanied by an HTML abstract. “Just Accepted” manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are accessible to all readers and citable by the Digital Object Identifier (DOI®). “Just Accepted” is an optional service offered to authors. Therefore, the “Just Accepted” Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the “Just Accepted” Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these “Just Accepted” manuscripts.

Ni-Catalyzed Enantioselective Conjunctive Coupling with C(sp³) Electrophiles: A Radical-Ionic Mechanistic Dichotomy

Gabriel J. Lovinger and James P. Morken*


Department of Chemistry, Merkert Chemistry Center, Boston College, Chestnut Hill, Massachusetts 02467, United States

Supporting Information Placeholder

ABSTRACT: The catalytic enantioselective conjunctive coupling of C(sp³) electrophiles can be accomplished with Ni catalysis. The enantioselectivity of the reaction is dependent on reaction mechanism with many substrates able to engage in an asymmetric process with Pybox-Ni complexes, whereas other substrates provide racemic product mixtures. The link between substrate structure and selectivity is addressed.

Chiral alkylboronate esters are important building blocks in contemporary organic chemistry and have been adopted for both academic and industrial asymmetric synthesis.¹ New methods for constructing these motifs from untapped classes of starting materials can therefore have a significant impact in a number of venues.² In this connection, we have been developing transition-metal catalyzed conjunctive cross-coupling reactions as a means to produce alkylboronate esters from vinylboron "ate" complexes and organic electrophiles.^{3,4} Preliminary studies have indicated that conjunctive coupling can be conducted catalytically and enantioselectively with either Ni^{3d} or Pd^{3a-c} complexes and, mechanistically, appears to proceed by way of a metal-induced metallate rearrangement. While these processes operate efficiently and enantioselectively with C(sp²) electrophiles, an important additional dimension in conjunctive coupling would arise by the introduction of reactions that employ C(sp³) electrophiles. In this report, we describe the utility of chiral Ni complexes in efficient catalytic conjunctive coupling reactions with alkyl electrophiles and describe how the reaction path – proceeding either through an enantioselective metal-induced metallate rearrangement, or through a non-selective radical-polar cross over manifold as

Scheme 1. Ni-Catalyzed Conjunctive Coupling of C(sp³) Electrophiles.




elaborated by Studer⁵ and Aggarwal^{6,7} – is intimately linked to substrate structure and therefore controllable by judicious substrate selection.

While most research in the field of catalytic cross-coupling⁸ has focused on the construction of C(sp²)-C(sp²) bonds, recent advances have expanded the scope of cross-coupling methods to include C(sp²)-C(sp³) and C(sp³)-C(sp³) couplings.⁹ The latter class of C-C bonds are often difficult to forge by cross-coupling due to mechanistic considerations involving challenging oxidative addition, slow transmetalation, and facile β-hydride elimination. In this light, use of alkyl halide electrophiles in conjunctive cross-coupling might appear to be fraught with complication. However, it was considered that the use of Ni complexes¹⁰ might ameliorate challenges associated with β-hydrogen elimination from Ni(alkyl) catalytic intermediates. Moreover, barriers to cross-coupling that arise from transmetalation of alkyl-M reagents¹¹ to catalytic centers might be avoided in conjunctive coupling since the transmetalation step is replaced with a metal-induced metallate rearrangement. We therefore began our investigations by subjecting boron "ate" complex **1**, generated from PhBpin and vinyl-lithium, to 3-phenyl-iodopropane (**2**) and either Ni(acac)₂ alone, or in combination with diamine **L1**¹², the optimal ligand for Ni-catalyzed conjunctive coupling of aryl electrophiles.^{3d} As depicted in Table 1, neither reaction furnished detectable quantities of conjunctive coupling product (**4**). A survey of other ligands indicated that bis(oxazoline), phosphine-oxazoline, and pyridyl-oxazoline are ineffective ligand scaffolds; however, tri-coordinate Pybox ligands furnished conjunctive coupling products with good levels of asymmetric induction. Modification of the ligand and nickel source revealed **L7** and [(methallyl)NiCl]₂ as a particularly competent combination, delivering 70% yield of conjunctive coupling product **4** in 96% ee (entry 10). In contrast to the enantioselective reaction with substrate **2**, when α-bromo-methyl acetate **3** was employed, the reaction furnished racemic product (entries 11 and 12). Unlike reaction of unactivated substrate **2**, reaction of compound **3** is efficient in the absence of ligand and, with addition of NaI, can furnish the racemic product **5** in outstanding yield.

Table 1. Ni-Catalyzed Conjunctive Cross-Coupling with C(sp³) Electrophiles

| entry | E-X | ligand | Ni salt | % yield ^a | er |
|-------------------|-----|--------|--------------------------------|----------------------|-------|
| 1 | 2 | none | Ni(acac) ₂ | <10% | nd |
| 2 | 2 | L1 | Ni(acac) ₂ | <10% | nd |
| 3 | 2 | L2 | Ni(acac) ₂ | <10% | nd |
| 4 | 2 | L3 | Ni(acac) ₂ | <10% | nd |
| 5 | 2 | L4 | Ni(acac) ₂ | <5% | nd |
| 6 ^b | 2 | L5 | Ni(acac) ₂ | 17 | 89:11 |
| 7 ^b | 2 | L6 | Ni(acac) ₂ | 36 | 99:1 |
| 8 ^{b,c} | 2 | L6 | Ni(acac) ₂ | 49 | 98:2 |
| 9 ^{b,c} | 2 | L6 | [(methallyl)NiCl] ₂ | 51 | 98:2 |
| 10 ^{b,c} | 2 | L7 | [(methallyl)NiCl] ₂ | 70 | 98:2 |
| 11 ^{b,c} | 3 | L7 | Ni(acac) ₂ | 34 | 50:50 |
| 12 ^d | 3 | | Ni(acac) ₂ | 92 | nd |


(a) Determined by NMR versus an internal standard. (b) Determined by chiral SFC analysis. (c) Catalyst solution pre-complexed for 1.5 h. (d) Reaction conducted at 22 °C without ligand and with one equivalent NaI added.




The marked difference in reaction selectivity when using substrate **2** versus **3** suggested that the operative mechanisms for these reactions are likely different. In light of recent studies by Studer⁵ and Aggarwal⁶ on radical-polar crossover reactions between organic halides and vinyl boron "ate" complexes, it was suspected that radical processes might intervene in the Ni-catalyzed conjunctive coupling reactions of particular substrates and alter their course. Three mechanistic scenarios were considered as plausible reaction pathways (Scheme 2), with all involving initial halogen atom abstraction¹³ to convert the C(sp³) halide to a carbon-centered radical and nickel complex **A**. In one direction, following the precedent of Studer and Aggarwal, reaction of the carbon radical with the olefinic "ate" complex might deliver α-boryl radical **B**, which undergoes single-electron-transfer with the organohalide substrate thereby regenerating the carbon radical and delivering transient carbocation **C**; cation **C** would be expected to undergo metallate shift and necessarily deliver racemic reaction product. Formation of enantiomerically-enriched reaction product could occur by one of two routes: first, recombination of the initially-generated carbon radical with Ni complex **A** might deliver complex **D** – a compound that could engage in selective metal-induced metallate shift similar to previously described Pd and Ni complexes. Alternatively, aligned with recent observations by Fu¹⁴ on cross-coupling of α-haloboronic esters, α-boryl radical **B** might reengage Ni complex **A** to furnish **E**; subsequent intramolecular transmetallation and reductive elimination would furnish nonracemic product. Following the

above-described mechanistic proposals, substrates that would furnish an electrophilic radical would be more prone to react with the electron-rich vinyl boron "ate" complex¹⁵ and undergo the radical chain process described by the Studer-Aggarwal cycle. Substrates leading to non-stabilized radicals could favor the metallate shift pathway since **A** would be expected to rapidly recombine with the initially-formed radical, or these substrates might operate the Fu-type cycle since single-electron transfer from **B** to an alkyl halide might be impeded by the redox potential of the nonactivated electrophile.

Scheme 2. Prospective Mechanisms for Ni-catalyzed Conjunctive Coupling of C(sp³) Electrophiles.



To disentangle the operative pathways for these conjunctive coupling reactions, a series of mechanistic experiments were undertaken. First, we examined reactions of α-bromo ester **3** in the presence of TEMPO. As depicted in equation 1 (Scheme 3), addition of one equivalent of the radical scavenger is sufficient to inhibit formation of **5**, thereby suggesting that radicals are indeed involved in the reaction of this substrate. Consistent with the non-stereospecific nature of the radical-polar crossover mechanism, conjunctive coupling between **3** and isotopically labeled "ate" complex **6** furnished the reaction product as a mixture of diastereomers (eq. 2). Of note, radical intermediates still appear to intervene in the enantioselective reactions of non-activated haloalkanes as indicated by the ring-opening and ring-closing reactions in equations 3 and 4 (Scheme 3)¹⁶ as well as the observation that TEMPO inhibits reaction of 1-iodobutane (eq. 5). However, the "Fu-type" cycle is ruled out as a mechanistic possibility by the observation that the enantioselective conjunctive coupling of iodobutane is a stereospecific process (eq. 6).

Table 3. Ni-Catalyzed Enantioselective Conjunctive Coupling with C(sp³) Electrophiles.^a

(a) Unless otherwise noted, the "ate" complexes were prepared by addition of vinylolithium to RB(pin). All yields represent isolated product obtained after chromatographic purification. (b) Compounds **4**, **9**, **23**, **24**, and **25** were also prepared by addition of PhLi to vinylB(pin) and provide products in 67, 65, 57, 47 and 44% yield, respectively. (c) Product isolated as the derived alcohol after peroxide oxidation. (d) (*S,S*)-L7 employed as ligand.

(1) Reviews, see: (a) Brown, H. S.; Singaram, B. *Acc. Chem. Res.* **1988**, *21*, 287. (b) *Organoboranes for Synthesis*, Ramachandran, P. V.; Brown, H. C., Eds. ACS Symposium Series 783; American Chemical Society: Washington, DC 2001. (c) *Stereodirected Synthesis with Organoboranes* Matteson, D. S. Springer: New York, 1995. (d) Leonori, D.; Aggarwal, V. K. *Accs. Chem. Res.* **2014**, *47*, 3174.

(2) For a recent review: Collins, B. S. L.; Wilson, C. M.; Myers, E. L.; Aggarwal, V. K. *Angew. Chem Int. Ed.* **2017**, *56*, 11700.

(3) (a) Zhang, L.; Lovinger, G. J.; Edelstein, E. K.; Szymaniak, A. A.; Chierchia, M. P.; Morken, J. P. *Science*. **2016**, *351*, 70. (b) Lovinger, G. J.; Aparece, M. D.; Morken, J. P. *J. Am. Chem. Soc.* **2017**, *139*, 3153. (c) Edelstein, E. K.; Namirembe, S.; Morken, J. P. *J. Am. Chem. Soc.* **2017**, *139*, 5027. (d) Chierchia, M.; Law, C.; Morken, J. P. *Angew. Chem. Int. Ed.* **2017**, *56*, 11870.

(4) For aligned processes, see: (a) Ishida, N.; Shimamoto, Y.; Murakami, M. *Org. Lett.* **2009**, *11*, 5434. (b) Ishida, N.; Shinmoto, T.; Sawano, S.; Miura, T.; Murakami, M. *Bull. Chem. Soc. Jpn.* **2010**, *83*, 1380. (c) Ishida, N.; Narumi, M.; Murakami, M. *Org. Lett.* **2008**, *10*, 1279. (d) Ishida, N.; Shimamoto, Y.; Murakami, M. *Org. Lett.* **2010**, *12*, 3179. (e) Ishida, N.; Ikemoto, W.; Narumi, M.; Murakami, M. *Org. Lett.* **2011**, *13*, 3008.

Lastly, for electrophiles with labile β-substituents (**30**), elimination is not observed. Current challenges to this catalytic cross-coupling include methyl-, *t*-butyl-, and benzyl-based electrophiles as well as alkyl migrating groups (inset, Table 3).

In conclusion, we have developed a highly enantioselective nickel-catalyzed conjunctive cross-coupling that can engage primary and secondary C(sp³) electrophiles and can tolerate a range of functional groups in the production of chiral alkylboronate ester products. Mechanistic experiments have elucidated the origin of the divergent reactivity between activated and unactivated electrophilic substrates and we anticipate that this knowledge will be useful in the design of new processes.

■ ASSOCIATED CONTENT

Supporting Information. Procedures, characterization and spectral data. This material is available free of charge via the Internet at <http://pubs.acs.org>.

■ AUTHOR INFORMATION

Corresponding Author

*morken@bc.edu

ORCID

James P. Morken 0000-0002-9123-9791

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGEMENTS

This work was supported by the NIH (GM-118641).

■ REFERENCES

- (5) Kischkewitz, M.; Okamoto, K.; Mück-Lichtenfeld, C.; Studer, A. *Science*, **2017**, *355*, 936.
- (6) Silvi, M.; Sandford, C.; Aggarwal, V. K. *J. Am. Chem. Soc.* **2017**, *139*, 5736.
- (7) For an overview, see: Hidasová, D.; Jahn, U. *Angew. Chem. Int. Ed.* **2017**, *56*, 9656.
- (8) For a comprehensive overview of catalytic cross-coupling, see: (a) de Meijere, A., Diederich, F., Eds. *Metal-Catalyzed Cross-Coupling Reactions*, 2nd Completely Revised and Enlarged ed., Vol. 1; Wiley-VCH: Weinheim, Germany, 2004. (b) de Meijere, A., Diederich, F., Eds. *Metal-Catalyzed Cross-Coupling Reactions*, 2nd Completely Revised and Enlarged ed., Vol. 2; Wiley-VCH: Weinheim, Germany, 2004.
- (9) For a recent review, see: Choi, J.; Fu, G. C. *Science*, **2017**, *356*, 152.
- (10) Overviews of mechanistic features in cross-coupling with Ni catalysts, see: (a) Hu, X. *Chem. Soc. Rev.* **2011**, *2*, 1867. (b) Tasker, S. Z.; Standley, E. A.; Jamison, T. F. *Nature* **2014**, *509*, 299.
- (11) Overview: Jana, R.; Pathak, T. P.; Sigman, M. S. *Chem. Rev.* **2011**, *111*, 1417.
- (12) For selected enantioselective Ni-catalyzed cross-coupling with Ni/diamine catalysts, see: (a) Schmidt, J.; Choi, J.; Liu, A. T.; Slusarczyk, M.; Fu, G. C. *Science*, **2016**, *354*, 1265. (b) Wilsily, A.; Tramutola, F.; Owston, N. A.; Fu, G. C. *J. Am. Chem. Soc.* **2012**, *134*, 5794. (c) Lu, Z.; Wilsily, A.; Fu, G. C. *J. Am. Chem. Soc.* **2011**, *133*, 8154. (d) Saito, B.; Fu, G. C. *J. Am. Chem. Soc.* **2007**, *129*, 9602.
- (13) For lead references, see: (a) Fu, G. C. *ACS Cent. Sci.* **2017**, *3*, 692. (b) Jahn, U. *Top. Curr. Chem.* **2012**, *320*, 323.
- (14) Schmidt, J.; Choi, J.; Liu, A. T.; Slusarczyk, M.; Fu, G. C. *Science*, **2016**, *354*, 1265.
- (15) For polar effects in radical addition reactions, see: Giese, B. *Angew. Chem. Int. Ed. Engl.* **1983**, *22*, 753.
- (16) For a review of cyclopropylcarbinyl radical chemistry, see: Nonhebel, D. C. *Chem. Soc. Rev.* **1993**, 347.

