


Nickel-Catalyzed Cross-Coupling of
Potassium Aryl- and
Heteroaryltrifluoroborates with
Unactivated Alkyl HalidesGary A. Molander,^{*,†} O. Andreea Argintaru,[†] Ioana Aron,[†] and Spencer D. Dreher[§]

Roy and Diana Vagelos Laboratories and Penn/Merck Laboratory for High
Throughput Experimentation, Department of Chemistry, University of Pennsylvania,
231 South 34th Street, Philadelphia, Pennsylvania 19104-6323, and
Department of Process Chemistry, Merck Research Laboratories,
P.O. Box 2000, Rahway, New Jersey 07065

gmolandr@sas.upenn.edu

Received November 9, 2010

ABSTRACT



A method for the cross-coupling of alkyl electrophiles with various potassium aryl- and heteroaryltrifluoroborates has been developed. Nearly stoichiometric amounts of organoboron species could be employed to cross-couple a large variety of challenging heteroaryl nucleophiles. Several functional groups were tolerated on both the electrophilic and the nucleophilic partners. Chemoselective reactivity of C(sp³)-Br bonds in the presence of C(sp²)-Br bonds was achieved.

Several recent studies have focused on the development of cross-coupling strategies to unite alkyl electrophiles with aryl nucleophilic partners.¹ Among these, the Suzuki–Miyaura reaction has emerged as one of the most powerful methods because of the low toxicity, air and water stability, functional group compatibility, and commercial availability of the organoboron compounds.^{2,3} Nickel catalysts were reported to be among the most successful for C(sp²)-C(sp³) bond formation via the Suzuki–Miyaura reaction.¹ Nevertheless, limitations to the developed method remain:^{4–6} depending on the

nucleophile, a significant excess of boronic acid is usually required, and *ortho*-substituted arylboronic acids only cross-couple to a limited extent. Perhaps most critically, only a few isolated examples have been reported to partner alkyl halides with heteroarylboron nucleophiles (e.g., indole-5-boronic acid, thiophene-3-boronic acid), and virtually all protocols explicitly have failed for other important heterocyclic systems.⁵

[†] University of Pennsylvania.[§] Merck Research Laboratories.

(1) (a) Rudolph, A.; Lautens, M. *Angew. Chem., Int. Ed.* **2009**, *48*, 2656–2670, and references cited therein. (b) Ejiri, S.; Odo, S.; Takahashi, H.; Nishimura, Y.; Gotoh, K.; Nishihara, Y.; Takagi, K. *Org. Lett.* **2010**, *12*, 1692–1695. (c) Hatakeyama, T.; Hashimoto, T.; Kondo, Y.; Fujiwara, Y.; Seike, H.; Takaya, H.; Tamada, Y.; Ono, T.; Nakamura, M. *J. Am. Chem. Soc.* **2010**, *132*, 10674–10676. (d) Owson, N. A.; Fu, G. C. *J. Am. Chem. Soc.* **2010**, *132*, 11908–11909. (e) Lundin, P. M.; Fu, G. C. *J. Am. Chem. Soc.* **2010**, *132*, 11027–11029. (f) Lou, S.; Fu, G. C. *J. Am. Chem. Soc.* **2010**, *132*, 1264–1266.

(2) For reviews, see: (a) Miyaura, N.; Suzuki, A. *Chem. Rev.* **1995**, *95*, 2457–2483. (b) Little, A. F.; Fu, G. C. *Angew. Chem., Int. Ed.* **2002**, *41*, 4177–4211.

(3) For studies and examples of palladium-catalyzed cross-coupling reactions of alkyl electrophiles, see: (a) Ariafard, A.; Lin, Z. *Organometallics* **2006**, *25*, 4030–4033. (b) Zhou, J.; Fu, G. C. *J. Am. Chem. Soc.* **2003**, *125*, 12527–12530. (c) Kirchhoff, J. H.; Netherton, M. R.; Hills, I. D.; Fu, G. C. *J. Am. Chem. Soc.* **2002**, *124*, 13662–13663. (d) He, A.; Falck, J. R. *J. Am. Chem. Soc.* **2010**, *132*, 2524–2525.


(4) Zhou, J.; Fu, G. C. *J. Am. Chem. Soc.* **2004**, *126*, 1340–1341.

(5) Gonzalez-Bobes, F.; Fu, G. C. *J. Am. Chem. Soc.* **2006**, *128*, 5360–5361.

(6) Duncton, M. A. J.; Estiarte, M. A.; Tan, D.; Kaub, C.; O'Mahony, D. J. R.; Johnson, R. J.; Cox, M.; Edwards, W. T.; Wan, M.; Kincaid, J.; Kelly, M. G. *Org. Lett.* **2008**, *10*, 3259–3262.

We envisioned that these limitations could be overcome through the use of the more robust potassium aryl- and heteroaryltrifluoroborates as nucleophilic coupling partners in the Ni-catalyzed cross-coupling with various alkyl electrophiles.⁷ Using the reaction between 1-bromo-3-phenylpropane and a stoichiometric amount of potassium (4-methoxyphenyl)-trifluoroborate (Scheme 1) as a template, a

Scheme 1. Optimization of the Cross-Coupling Conditions



wide variety of catalyst/ligand combinations, solvents, bases, and temperatures were screened.⁸ The best coupling conditions were determined to be 10 mol % of NiBr₂·glyme, 10 mol % of 4,7-diphenyl-1,10-phenanthroline (bathophenanthroline), and 3 equiv of LiHMDS in *sec*-butanol.⁸

With these results in hand, we investigated the cross-coupling of 2-(bromomethyl)tetrahydro-2H-pyran with different potassium aryltrifluoroborates (Table 1).

Both electron-rich and electron-poor substituted trifluoroborates successfully underwent cross-coupling using nearly stoichiometric amounts of the nucleophile (1.02 equiv). Sterically hindered *ortho*-substituted trifluoroborates also underwent the reaction in good yields (Table 1, entries 2 and 9). Additionally, substrates containing a secondary alcohol (entry 7), a ketone (entries 8 and 9), and an alkene (entry 5) were tolerated. Importantly, the reactions could be performed on gram scale using only a 1 mol % catalyst loading with little effect on the yield (entry 2).

To expand the scope of this method, we examined the cross-coupling of the same alkyl bromide with a large variety of heteroaryltrifluoroborates including furans, benzo- and dibenzofuran, pyridines, pyrimidines, thiophenes, quinolines, indole, and imidazole, again using virtually stoichiometric organotrifluoroborates (Table 2).

The corresponding cross-coupled products were obtained in moderate to good yields. Given the extraordinarily high propensity for heteroarylboron reagents to undergo protodeboronation,⁹ these transformations represent a significant advance in the formation of core substructures of greatest

Table 1. Cross-Coupling of 2-(Bromomethyl)tetrahydro-2H-pyran with Potassium Aryltrifluoroborates

entry	R-BF ₃ K	yield %	entry	R-BF ₃ K	yield %
1	MeO-C ₆ H ₄ -BF ₃ K R = H, 86 R = OMe, 72		6	2-naphthyl-BF ₃ K	68
2	Ph-BF ₃ K R = <i>o</i> -Me, 76 R = <i>m</i> -Me, 81 (76) ^b R = <i>p</i> -Me, 70		7	4-(1-hydroxyethyl)phenyl-BF ₃ K	81
3	F ₃ C-C ₆ H ₄ -BF ₃ K	64	8 ^a	4-(1-oxoethyl)phenyl-BF ₃ K	66
4	F-C ₆ H ₄ -BF ₃ K	71	9 ^a	2-(1-oxoethyl)phenyl-BF ₃ K	68
5	CH ₂ =C ₆ H ₄ -BF ₃ K	53			

^a Reactions run at 80 °C. ^b Reaction performed on 5 mmol scale using 1 mol % catalyst loading.

Table 2. Cross-Coupling of 2-(Bromomethyl)tetrahydro-2H-pyran with Potassium Heteroaryltrifluoroborates

entry	R-BF ₃ K	yield %	entry	R-BF ₃ K	yield %
1	X-BF ₃ K X = O, 71 X = S, 63		7	2-benzofuranyl-BF ₃ K	78
2	2-thienyl-BF ₃ K	66	8	2-pyridyl-BF ₃ K	82
3	2-imidazolyl-BF ₃ K	35	9	2-quinolyl-BF ₃ K	70
4	X-BF ₃ K X = O, 68 X = S, 81 (74) ^a		10	2-pyridyl-BF ₃ K	71
5	2-indolyl-BF ₃ K X = O, 80 X = S, 58		11	2-pyridyl-BF ₃ K	59
6	2-pyrimidyl-BF ₃ K R = H, 77 R = OMe, 71		12	2-pyridyl-BF ₃ K	51
			13	2-(4-bromophenyl)pyridyl-BF ₃ K	63

^a Reaction performed on 5 mmol scale using 1 mol % catalyst loading.

(7) For reviews on potassium organotrifluoroborates in cross-coupling reactions, see: (a) Molander, G. A.; Ellis, N. M. *Acc. Chem. Res.* **2007**, *40*, 275–286. (b) Darses, S.; Genet, J. P. *Chem. Rev.* **2008**, *108*, 288–325. (c) Molander, G. A.; Canturk, B. *Angew. Chem., Int. Ed.* **2009**, *48*, 9240–9286. (d) Stefani, H. A.; Cella, R.; Vieira, A. S. *Tetrahedron* **2007**, *63*, 3623–3658.

(8) The screening was performed through microscale High Throughput Experimentation (HTE). Dreher, S. D.; Dormer, P. G.; Sandrock, D. L.; Molander, G. A. *J. Am. Chem. Soc.* **2008**, *130*, 9257–9259. See Supporting Information.

(9) (a) Molander, G. A.; Canturk, B.; Kennedy, L. E. *J. Org. Chem.* **2009**, *74*, 973–980, and references therein. (b) Knapp, D. M.; Gillis, E. P.; Burke, M. D. *J. Am. Chem. Soc.* **2009**, *131*, 6961–6963.

interest in the pharmaceutical, agrochemical, and materials science sectors. As in the case of the aryltrifluoroborates, a heteroaryl cross-coupling could be carried out on gram scale using only 1 mol % of the nickel catalyst (entry 4). Of additional interest, the C(sp²)–Br bond in 5-bromo-3-pyridyltrifluoroborate remained intact during the cross-coupling reactions, maintaining the potential for further functionalization of the pyridine (entry 13).

To test the general applicability of our method to other electrophiles, potassium 2-benzofuranyl- and 4-pyridinyltrifluoroborate were reacted with various functionalized alkyl halides (Table 3).

Table 3. Cross-Coupling of Diverse Alkyl Halides with Potassium 2-Benzofuranyl- and 4-Pyridinyltrifluoroborates

entry	Alkyl-X	yield %	entry	Alkyl-X	yield %
1		1, 79 2, 81	7		1, 70 2, 61
2		2, 61	8		1, 84 2, 63
3		1, 80 2, 76	9		1, 60 2, 71
4		1, 76 2, 78	10		1, 67 2, 68
5		2, 72 ^a	11		1, 71 2, 69
6		1, 63 2, 68	12		1, 62 2, 63

^a Isolated as the *s*-Bu ester.

Electrophiles containing an acetal, a benzyl ether, and a distal olefin were tolerated (Table 3). *p*-Bromo-(3-bromopropyl)-benzene smoothly underwent chemoselective cross-coupling at the C(sp³)–Br bond, leaving the C(sp²)–Br bond available for further functionalization (entry 2). Another bidirectional functionalization opportunity arises from the selective reactivity of alkyl bromides and iodides over alkyl chlorides (entries 7 and 8). The same set of conditions could be applied to the successful cross-coupling of secondary bromides and iodides (entries 9–12). Although stereochemical studies have not been conducted, on the basis of previously reported investigations⁵ we would anticipate that these processes would transpire via radical intermediates, and thereby the stereochemical fidelity of enantioenriched starting materials would not be transferred to the products.

The conditions employed for the cross-coupling of alkyl iodides and bromides were not successful when applied to the more challenging alkyl chlorides. As previously reported in the literature,⁵ the use of L-prolinol as a ligand was required for the reaction of alkyl chlorides (Table 4). Although the *ortho*-substituted boronic acids have been problematic in the Ni-catalyzed cross-coupling with alkyl chlorides,⁵ use of the more stable potassium trifluoroborates led to the formation of the desired product in 60% yield (entry 5).¹⁰

In conclusion, we have developed an efficient method for the Ni-catalyzed cross-coupling of unactivated halides. Bathophenanthroline was used as a ligand for the Suzuki–Miyaura

Table 4. Cross-Coupling of Various Alkyl Chlorides with Potassium Aryl- and Heteroaryltrifluoroborates

entry	Alkyl-Cl	Ar/HetAr	yield %
1			60
2			54
3			47
4			42
5			60
6			49 ^a

^a Reaction using KHMDS (3 equiv).

cross-coupling of alkyl bromides and iodides with aryl and heteroaryl nucleophiles, while L-prolinol was required for the cross-coupling of alkyl chlorides. Several major advances have derived from these studies: (1) Only 1 equiv of the organoboron reagent is necessary, as opposed to 1.2 to 2 equiv reported in previous studies; (2) both heteroaryl- and sterically hindered *ortho*-substituted organoborons, previously reported to be problematic substrates, can be coupled in good yields; and (3) on a larger scale, low (1 mol %) catalyst loadings can be used to effect efficient cross-coupling. The use of organotrifluoroborates thus represents a significant advance in cross-coupling with alkyl halide electrophiles.

Acknowledgment. We thank Frontier Scientific for boronic acids. Deidre L. Sandrock (University of Pennsylvania) is acknowledged for performing HTE experiments. Financial support has been provided by the NIH General Medical Sciences (R01 GM035249) and the NSF GOALI program (CHE-0848460). I.A. thanks CURF (University of Pennsylvania) for a fellowship. Dr. Rakesh Kohli (University of Pennsylvania) is acknowledged for obtaining HRMS data.

Supporting Information Available: Experimental procedures and compound characterization data. This material is available free of charge via the Internet at <http://pubs.acs.org>.

OL102717X

(10) Under these conditions, reactions of other heteroaryltrifluoroborates (furans, thiophenes, pyrazoles) proceeded in relatively low yield (<40%, data not shown). The cross-coupling of alkyl chlorides with aryl- and heteroaryltrifluoroborates is currently under study.