Synthesis of Bis(phosphinomethyl)amines via Bis(hydroxymethyl)phosphonium Salts. Isolation of 9,9-Bis(hydroxymethyl)-9-phosphoniabicyclo[3.3.1]nonane Hydrogensulfate and Chloride Salts, and the Crystal Structures of $[PPh_2(CH_2OH)_2]^+CI^-$ and $[(C_6H_{11})_2PCH_2]_2^-$ NCHMePh[†]

John Fawcett,^a Peter A. T. Hoye,^b Raymond D. W. Kemmitt,^{*,a} David J. Law^a and David R. Russell^a

^a Department of Chemistry, The University of Leicester, Leicester LE1 7RH, UK ^b I.T.C., Albright & Wilson Ltd., West Midlands B694 4LN, UK

Aminomethylphosphines have been known for some 30 years since the discovery by Coates and Hoye¹ that hydroxymethylphosphines react with primary and secondary amines to give aminomethylphosphines. Since then aminomethylphosphines have been used in various commercial applications such as flame retardants^{2,3} and metal sequestrants.^{4,5} Various aminomethylphosphines have been prepared to date,6-8 including compounds with cage structures⁹ and with chiral centres at phosphorus.¹⁰ Despite this interest almost all published aminomethylphosphine chemistry concentrates on phenyl- and diphenyl-substituted phosphines, and characterisation details have been minimal with almost no elemental analysis or phosphorus NMR data. The most favoured route to aminomethylphosphines has been the modified Mannich reaction using hydroxymethylphosphines or hydroxymethylphosphonium salts as the key starting materials. The reaction of new hydroxymethylphosphonium salts, having alkyl or aryl substituents on phosphorus, with a variety of primary amines leads to new functionalised phosphines.

Results and Discussion

Bis(hydroxymethyl)phosphonium chloride salts 1 were prepared in quantitative yields (95-99%) by treatment of a secondary phosphine, PPh₂H or P(C₆H₁₁)₂H, with aqueous formaldehyde and concentrated hydrochloric acid, Scheme 1. The compounds are highly crystalline white solids which are



Scheme 1

soluble only in polar solvents such as water or methanol. The phenyl derivative 1a has been reported previously¹¹ although only the melting point and phosphorus analysis were given for characterisation. The melting point obtained for 1a is in agreement with that previously obtained; $162 \degree C$ (lit., ¹¹ 165 °C). The bis[bis(hydroxymethyl)phosphonium] sulfate and [bis-(hydroxymethyl)phosphonium] hydrogensulfate salts 3a and 3b were also prepared by treating the mixed isomers of $C_8H_{14}PH 2$ with aqueous formaldehyde and concentrated sulfuric acid, Scheme 2. Despite the presence of both HSO_4^- and SO_4^{2-} salts, the ³¹P NMR spectrum shows only two signals corresponding to the [3.3.1] and [4.2.1] isomers. Interestingly, the chemical shifts following quaternisation are significantly different for each isomer, $\delta - 54.6$ (2a), -49.2 (2b), +23.4 (3a) and +55.7 (3b). The chemical shift differences between the isomers 3a and 3b were taken to indicate a difference in the properties of the two isomeric species. Work by Hoye¹² has shown that phosphonium sulfate/hydrogen sulfate salts can be decomposed to give methylphosphine oxide compounds by refluxing in sulfuric acid and removal of formaldehyde. Using this method the phosphonium salts 3a and 3b were refluxed at 150 °C in 36% sulfuric acid, Scheme 2. The reaction mixture was monitored by ³¹P NMR spectroscopy, revealing the formation of the [3.3.1] and [4.2.1] isomers of the decomposition product $(C_8H_{14})P(O)Me$ at δ 56.0 and 88.0

[†] Supplementary data available: see Instructions for Authors, J. Chem. Soc., Dalton Trans., 1993, Issue 1, pp. xxiii-xxviii.

Non-SI unit employed: mmHg \approx 133 Pa.



Scheme 2 $X = HSO_4^-$ or $\frac{1}{2}SO_4^{2^-}$. (*i*) HCHO, 98% H₂SO₄; (*ii*) 36% H₂SO₄, 150 °C, 13 h

Table 1 Selected bond lengths (Å), angles (°) and non-bonded contacts (Å) with estimated standard deviations (e.s.d.s) in parentheses for the phosphonium salt $[PPh_2(CH_2OH)_2]^+Cl^-$

C(2)-P-C(1)	107.8(3)	C(21)-P-C(2)	110.6(2)
C(21)-P-C(1)	108.8(2)	C(31)-P-C(21)	111.3(2)
C(31)-P-C(1)	111.2(2)	O(1)-C(1)-P	110.6(4)
C(31)-P-C(2)	107.1(2)	O(2)-C(2)-P	112.1(4)
C(1)-P	1.822(5)	C(1)-O(1)	1.400(7)
C(2)-P	1.838(6)	C(2)-O(2)	1.409(7)
C(21)-P	1.786(3)	H(1)-O(1)	0.79(7)
C(31)-P	1.781(3)	H(2)-O(2)	0.68(8)
$\begin{array}{c} H(1) \cdots Cl \\ H(2) \cdots Cl \end{array}$	2.227 2.367	$\begin{array}{c} O(1) \cdots Cl \\ O(2) \cdots Cl \end{array}$	3.020 3.030



respectively. After a reaction time of 13 h the spectrum of the mixture showed total conversion of 3b into the corresponding methylphosphine oxide and only about 10% conversion of 3a. Compound 3a was then isolated by recrystallisation in a 79.4% yield based on the original isomer ratio of [3.3.1]:[4.2.1] =58.5:41.5. It was thought, from molecular models, that the [4.2.1] isomer 3b would be less stable than typical hydroxymethylphosphonium sulfate salts due to apparent bond strain about phosphorus. However, the present studies show that it has similar reactivity to other hydroxymethylphosphonium sulfate salts under the given conditions, whereas isomer 3a appears less reactive than was predicted. This is probably due to the stability of the two six-membered heterocyclic rings present in the symmetrical isomer of C₈H₁₄PH. Various derivatives of the [3.3.1] and [4.2.1] isomers of $C_8H_{14}PH$ have been reported over the past 25 years.¹³ In all these studies the compounds are a mixture of isomers. The present work thus provides the first example of a derivative of 9-phosphabicyclo[3.3.1]nonane being isolated from its [4.2.1]



Fig. 1 Molecular structure of $[PPh_2(CH_2OH)_2]^+Cl^-$

 $2[PR_2(CH_2OH)_2]^+CI^- \xrightarrow{(i)} 2PR_2(CH_2OH) \xrightarrow{(ii)} (R_2PCH_2)_2NR'$



Scheme 3 (i) NEt₃; (ii) NR'H₂, water-methanol, 60 °C, 1-3 h

isomer. By treating a hot aqueous solution of the salt **3a** with barium chloride the corresponding chloride salt **4** was readily obtained.

As the phosphonium salts 1a, 1b, 3a, 3b and 4 were so highly crystalline, an X-ray diffraction study was carried out on $[PPh_2(CH_2OH)_2]^+Cl^-$ 1a. The molecular structure is shown in Fig. 1 and selected bond lengths and angles in Table 1. The compound exists as a quaternary phosphonium cation with a chloride anion. Intermolecular contacts are indicated by the short distances of the $H(1) \cdots Cl$ and $H(2) \cdots Cl$ non-bonded contacts (Table 1). The O··· Cl distances are also typical for OH···Cl hydrogen bonds of 2.99-3.05 Å.14 The geometry around phosphorus is essentially tetrahedral, with the C-P-C angles in the range $107.1-111.3^{\circ}$. The P-C(1) and P-C(2) bond lengths of 1.822(5) and 1.838(6) Å are comparable to those reported for tertiary phosphonium alkyls of 1.800(15) Å.¹⁵ The P-C(21) and P-C(31) bond distances of 1.786(3) and 1.781(3) Å also compare with those typical for tertiary phosphonium aryl compounds of 1.793(11) Å.¹⁵ The C(1)–O(1) and C(2)–O(2) bond lengths of 1.400(7) and 1.409(7) Å appear slightly short when compared to typical CH₂-OH bond distances of 1.426(11) Å,¹⁵ but this is presumably a consequence of the presence of the positively charged phosphonium ion which would have the effect of shortening these bonds.

The phosphines **5a**, **6a** and **7–10** were prepared by treating the phosphonium salt 1 with triethylamine followed by addition of a primary amine NRH₂, and gentle reflux for 1 h in a methanol-water solution, Scheme 3. The amines used were (R)-(+)-1-methylbenzylamine (**5a**), L-alanine ethyl ester (**6a**), L-alanine methyl ester (**7**), (R)-(+)-bornylamine (**8**), ethanolamine (**9**) and allylamine (**10**). The phosphines **5b** and **6b** were prepared in a similar fashion by treating the phosphonium salt 1b with triethylamine and a primary amine NR'H₂. The compounds were mainly isolated as highly viscous oils with yields in the range 61-95%. Triethylamine and solvents evident in the ¹H NMR spectra of the crude oils required long periods (10–100 h)



Scheme 4 (i) NEt₃, then $H_2NCHMePh$; (ii) NEt₃, then NEt₂H

under high vacuum, with warming, for complete removal. The phosphine **6a**, however, was isolated as a white solid after several days under high vacuum. The phosphine 5b was isolated by addition of methanol to a solution of it in toluene at 15 °C to give large colourless crystals in 56% yield. Attempts to recrystallise 6a, or to induce crystallisation in 5a, 6b and 7-10 using the solvents dichloromethane-light petroleum, toluene-methanol and boiling PriOH, were unsuccessful. The mass spectrum of **5b** showed peaks at m/z 541, corresponding to the molecular ion, and at 211, assigned to the fragment $[(C_6H_{11})_2PCH_2]^+$. The phosphine 11 was prepared by treating the pure [3.3.1] isomer of the phosphonium salt **3a** with triethylamine to form the hydroxymethylphosphine (C_8H_{14})-PCH₂OH. This was extracted with toluene and refluxed for 3 h in the presence of $(R-(\pm)-1-phenylethylamine, Scheme 4.$ Significantly more forcing conditions were required to prepare 11 than for the other isostructural aminophosphines 5a-10. This may be due to the presence of bulky $C_8H_{14}P$ and NCH(Me)Ph units. In contrast the mixed-phosphine isomers 12a and 12b were readily prepared by treating the mixed phosphonium salt isomers 3a and 3b with tri- and di-ethylamine in a watermethanol solution at 60 °C, to give [3.3.1] and [4.2.1] isomers 12a and 12b in 84% yield with an isomeric ratio of 3:5, Scheme 4. The hydroxymethylphosphine intermediates PPh₂(CH₂OH), $P(C_6H_{11})_2(CH_2OH)$ and $P(C_8H_{14})(CH_2OH)$, [3.3.1] isomer were monitored in their respective reaction mixtures by ³¹P NMR spectroscopy using single peaks at $\delta - 12.0$, -0.4 and +1.0 respectively. The phosphines **5a**, **6a** and **7** have been reported previously $^{7,16-18}$ although not in any detail, **5a** only in metal complexes with no NMR or analytical data being given.¹⁷ Phosphine 7 has been reported in rhodium(1) complexes but no data for the free phosphine were cited.¹⁸ The ¹H, ¹³C-{¹H} and ³¹P-{¹H} NMR spectra of the

aminomethylphosphines are given in the Experimental section. In phenyl-substituted phosphines there is a deshielding of the PCH₂N protons as compared to the cyclohexyl derivatives, which can be attributed to the deshielding effect of the phenyl rings as has been observed previously in organophosphines.¹⁹ The ¹H NMR spectra of phosphines 5-8 also showed interesting features. In particular, the diastereotopic CH₂ groups appeared as an AB pattern²⁰ with the low-frequency side exhibiting a larger coupling to phosphorus; the phosphorus-proton coupling was confirmed by ${}^{31}P{}{}^{1}H$ heteronuclear shift correlation NMR spectra. Asymmetric ABX spin systems have been previously observed in aminophosphines,^{16,17,21} and in these and other organophosphorus compounds ${}^{2}J(PH)$ is much larger when the coupled proton lies close to the orbital of the phosphorus(III) lone pair and is small when remote.^{19,21} In view of the availability of crystals of 5b it was therefore of interest to investigate its structure to determine the orientation of the α -methylene protons with respect to phosphorus.

The molecular structure of compound **5b** is illustrated in Fig. 2, and selected bond lengths and angles are shown in Table 2. Fig. 2 gives a view looking down onto the nitrogen lone pair. The diagram clearly shows the geometry of the bulky



Fig. 2 Molecular structure of $[(C_6H_{11})_2PCH_2]_2NCHMePh$. All cyclohexyl and phenyl H atoms have been omitted

Table 2 Selected bond lengths (Å) and angles (°) with e.s.d.s in parentheses for $[(C_6H_{11})_2PCH_2]_2NCHMePh$

$\begin{array}{c} C(31) - P(1) - C(21) \\ C(51) - P(2) - C(4) \\ C(51) - P(2) - C(41) \end{array}$	104.2(2) 99.9(2) 106.8(2)	C(4)-N-C(1) C(4)-N-C(3)	112.7(4) 110.8(4)
C(3)-P(1) C(4)-P(2) C(4)-N C(3)-N C(1)-N	1.856(5) 1.861(4) 1.477(6) 1.483(5) 1.475(6)	P(1)-C(21) P(1)-C(31) P(2)-C(41) P(2)-C(51)	1.850(5) 1.862(6) 1.875(5) 1.865(5)

substituents about the central atom. The C-N-C bond angles [110.8(4)–113.7(4)°] reflect the presence of bulky substituents about nitrogen as compared to NH₃, with H-N-H bond angles (106.8°) ,²² but are far from the near planarity of the extremely bulky N(SiMe₃)₃, Si–N–Si 119.6^o.²³ The C–P–C angles between phosphorus cyclohexyl groups are large $[104.2(2)-106.8(2)^{\circ}]$, typically comparable to those in PBu'₃ (105.7°).²⁴ The other C-P-C angles between cyclohexyl groups and PCH₂N carbon atoms C(3) and C(4) are more moderate [97.8(2)-101.5(2)°], being only slightly larger than in PMe₃ (98.9°).²⁴ The distances P(1)-C(3) and P(2)-C(4) appear within the typical range for P-C single bonds of 1.87(2) Å,¹⁴ as are the phosphoruscyclohexyl bond distances. The distances N-C(3) and N-C(4) also compare favourably with the N–C bond lengths in NMe₃ (1.47 Å).^{23,25} The approximate dihedral angles between the theoretical position of the phosphorus lone pair for P(1) and the methylene protons H(31) and H(32) are 145 and 107° respectively. For the phosphorus atom P(2) and the methylene protons H(41) and H(42) the dihedral angles are 161 and 87° respectively. It is therefore apparent that in the solid state the methylene protons do occupy different positions with respect to the lone pair on phosphorus. In solution, however, the conformation of the phosphine may be different and we note that on cooling a CD_2Cl_2 solution of **5b** to -40 °C the resonances due to the PCH₂ protons broaden, but at this temperature the phosphine crystallises from solution preventing further studies. From 0 to 100 °C there is no apparent change in the PCH₂ signals.

Experimental

Melting points were measured in air on a Gallenkamp apparatus and are uncorrected. Hydrogen-1 NMR spectra were recorded on a Bruker AM 300 spectrometer at 300.13 MHz or on a JEOL EM 390 spectrometer at 90 MHz, with SiMe₄ (δ 0.0) as internal

reference, positive values being to high frequency. The ${}^{13}C-{}^{1}H$ spectra were recorded in [²H₁]chloroform, unless otherwise stated, on a Bruker AM 300 spectrometer operating at 75.47 MHz with SiMe₄ as internal reference, 26 ${}^{31}P-{}^{1}H$ spectra on a JEOL JNM-FX60 spectrometer at 24.15 MHz with $[P(OH)_4]^+$ in $[^2H_2]$ water (δ 0.0) as external reference and ${}^{31}P_{-1}H$ heteronuclear shift correlation spectra on a Bruker AM 300 spectrometer at 120.06 MHz with the same reference. Experiments were carried out under a dry, oxygen-free, nitrogen atmosphere using solvents dried and distilled under nitrogen prior to use. The compounds diphenylphosphine, diethylamine, $D(+)-\alpha$ -methylbenzylamine, L-alanine ethyl ester hydrochloride, ethanolamine and allylamine were used as supplied by Aldrich. L-Alanine methyl ester hydrochloride and endo-(1R)-1,7,7-trimethylbicyclo[2.2.1]heptan-2-ylamine were used as supplied from Fluka. Dicyclohexylphosphine and the mixed isomers of phosphabicyclononane (C8H14PH) were used as supplied by Albright & Wilson Ltd. Microanalyses were carried out by Butterworth Laboratories Ltd.

Preparation of Bis(hydroxymethyl)phosphonium Salts.—(i) [PPh₂(CH₂OH)₂]⁺Cl⁻ 1a. Solutions of formaldehyde (9 cm³, 40%) and of HCl (5 cm³, 36%) were added to stirred diphenylphosphine (10 g, 54 mmol). The mixture became hot and homogeneous. On cooling a white solid crystallised out. This was filtered off and recrystallised from boiling methanol to give the compound 1a (15.0 g, 99%) (Found: C, 59.5; H, 5.8; P, 10.6. Calc. for C₁₄H₁₆ClO₂P: C, 59.6; H, 5.8; P, 11.0%), m.p. 160–162 °C. ³¹P-{¹H} NMR (CD₃OD): δ 16.7. X-Ray quality crystals were grown slowly from methanol–diethyl ether at -15 °C in air.

(*ii*) $[P(C_6H_{11})_2(CH_2OH)_2]^+Cl^-$ **1b.** Solutions of formaldehyde (96 cm³, 40%) and of HCl (47.5 cm³, 36%) were added to stirred dicyclohexylphosphine (100.0 g, 0.5 mol). The mixture became hot, and upon cooling a white solid crystallised out. This was filtered off and recrystallised from boiling PrⁱOH to give the required compound **1b** (144.3 g, 98%) (Found: C, 57.2; H, 9.5; P, 10.2. $C_{14}H_{28}ClO_2P$ requires C, 57.0; H, 9.6; P, 10.5%), m.p. 170–173 °C. NMR (CD₃OD): ¹H(90 MHz), δ 0.9–2.3 (m, 22 H, C₆H₁₁), 4.3 [d, 4 H, CH₂, ²J(PH) 3.0 Hz] and 4.5 (br s, 2 H, OH); ³¹P-{¹H} (24 MHz), δ 28.3.

(*iii*) $2[\dot{P}(C_8H_{14})(CH_2OH)_2]^+SO_4^{2-}/[P(C_8H_{14})(CH_2-OH)_2]^+HSO_4^-$, mixed isomers **3a** and **3b**. Solutions of formaldehyde (405 cm³, 40%) and of H₂SO₄ (180 cm³, 98%) were added to a stirred sample of C₈H₁₄PH (mixture of [3.3.1] and [4.2.1] isomers, 58.5:41.5) (365.3 g, 2.51 mol). The temperature rose to 60 °C over 1 h. On cooling two layers separated. The lower aqueous phase was decanted, washed with toluene and filtered through Dicalite. The solvent was removed under reduced pressure yielding a white solid which was recrystallised from boiling methanol to give the required compounds **3a** and **3b** (969 g, 81.3%), m.p. 190–200 °C. ³¹P-{¹H</sup> NMR (D₂O, 24 MHz): δ 23.0 (s, 1P, [3.3.1] isomer) and 55.7 (s, 1P, [4.2.1] isomer).

(iv) Isolation of compound 3a from the isomeric mixture of 3a and 3b. A sample of compounds 3a and 3b (125 g, 0.26 mol) was dissolved in H_2SO_4 (70 cm³, 36%) in a flask fitted with a dropping funnel, thermometer and a Dean and Stark condenser, The solution was heated to a constant 150 °C. Distillate (about 10 cm³) was removed at regular intervals and replaced by distilled water (10 cm³). This procedure was continued for 13 h, samples being monitored by ³¹P NMR spectroscopy. The reaction mixture was then diluted with distilled water (60 cm³) and refluxed at 105-110 °C. Further distillate was collected to remove any remaining formaldehyde. The solution was then filtered hot through carbon to remove discolourations and neutralised with calcium carbonate (100 g, 1.0 mol). It was filtered and the solvent removed under reduced pressure. The resulting solid was recrystallised from boiling methanol to give isomer **3a** (58.1 g, 46.5% recovery, based on the [3.3.1] isomer), m.p. 208–210 °C. ³¹P-{¹H} NMR (D₂O): δ 21.8 (s, 1P, [3.3.1] isomer) (Found: P, 11.0. Requires 10.3-12.3%, ratio 35:65, mono-:di-anion).

(v) $[P(C_8H_{14})(CH_2OH)_2]^+Cl^- 4$. The phosphonium salt 3a (5 g) was dissolved in hot distilled water (30 cm³) and barium chloride (2.5 g, 10 mmol) in distilled water (30 cm³) was added. The solution was allowed to cool and filtered through Celite. The solvent was removed under reduced pressure and the resulting solid recrystallised from boiling methanol to yield the required compound 4 (3.56 g, 84%), m.p. 248–250 °C (Found: C, 50.5; H, 8.3; P, 12.8. C₁₀H₂₀ClO₂P requires C, 50.3; H, 8.4; P, 13.0%). ³¹P-{¹H} NMR (D₂O): δ 23.4 (s, 1P, [3.3.1] isomer).

Preparation of Bis(diphenylphosphinomethyl)amines: General Method.—A slight excess of triethylamine was added to a stirred solution of $[PPh_2(CH_2OH)_2]Cl$ 1a in water-methanol (2:1, ca. 50 cm³). To this solution was added 1 equivalent of primary amine. The mixture was heated to reflux for 1 h. On cooling two layers separated. The product was extracted with dichloromethane (40 cm³), dried over magnesium sulfate and filtered. The solvent was then removed under high vacuum over a minimum of 24 h to give the products as either a highly viscous oil or a white solid, depending on the amine.

(i) $(Ph_2PCH_2)_2NCHMePh$ **5a**. The phosphonium salt **1a** (10 g, 35 mmol) with triethylamine (5.2 cm³, 35 mmol) and D(+)- α -methylbenzylamine (2.3 cm³, 18 mmol) gave compound **5a** as a highly viscous oil (8.27 g, 91.4%) (Found: C, 79.8; H, 6.8; N, 2.3. C₃₄H₃₃NP₂ requires C, 79.4; H, 6.2; N, 2.6%). NMR: ¹H, δ 1.55 [d, 3 H, Me, ³J(HH) 6.8], 3.8 [ABX, 2 H, PCH₂, ²J(HH) 13.2, ²J(PH) 6.0], 4.05 [ABX, 2 H, PCH₂, ²J(HH) 13.2, ²J(PH) 6.0], 4.05 [ABX, 2 H, PCH₂, ²J(HH) 13.2, ²J(PH) 0], 5.0 (m, 1 H, CH) and 7.35–7.8 (m, 25 H, Ph); ¹³C-{¹H}, δ 15.9 (s, Me), 56.2 [d, PCH₂, J(PC) 5.0], and 61.3 [t, NCH, ³J(PC) 9.0 Hz]; ³¹P-{¹H}, δ -27.8 (s).

(*ii*) (Ph₂PCH₂)₂NCHMeCO₂Et **6a**. The phosphonium salt **1a** (10 g, 35 mmol) with triethylamine (5.2 cm³, 36 mmol) and Lalanine ethyl ester hydrochloride (4.44 g, 29 mmol) neutralised with triethylamine (5.2 cm³, 36 mmol) gave **6a** as a white airsensitive solid (5.58 g, 61.5%) (Found: C, 71.5; H, 6.5; N, 2.9. Calc. for C₃₁H₃₃NO₂P₂: C, 72.5; H, 6.5; N, 2.9%), m.p. 76– 77 °C. NMR: ¹H, δ [d, 3 H, Me, ³J(HH) 7.1], 1.15 [t, 3 H, CH₂Me, ³J(HH) 7.1], 3.5 [ABX, 2 H, PCH₂, ²J(HH') 13.2, ²J(PH') 9.3], 3.9 [ABX, 2 H, PCH₂, ²J(HH') 13.2, ²J(PH) 0.0], 4.05 [m, 2 H, OCH₂Me, ²J(HH') 12.0, ³J(HH) 7.1, ⁵J(HH) 3.6], 4.3 [m, 1 H, NCH, ⁵J(HH) 3.6, ³J(HH) 7.1, ⁴J(PH) 11.0] and 7.1–7.6 (m, 20 H, Ph); ¹³C-{¹H}, δ 14.0 (s, Me), 14.5 (s, Me), 54.9 [d, PCH₂, J(PC) 4.9], 58.3 [t, NCH, ³J(PC) 9.4 Hz], 59.9 (s, OCH₂) and 172.7 (s, C=O); ³¹P-{¹H}, δ -26.4 (s). (*iii*) (Ph₂PCH₂)₂NCHMeCO₂Me 7. The phosphonium salt **1a**

(*iii*) (Ph₂PCH₂)₂NCHMeCO₂Me 7. The phosphonium salt **1a** (5 g, 17.7 mmol), triethylamine (2.5 cm³, 34 mmol) and L-alanine methyl ester hydrochloride (1.24 g, 8.9 mmol) gave compound 7 as an air-sensitive sticky white solid (3.56 g, 80.5%) (Found: C, 72.0; H, 6.2; N, 2.7. $C_{30}H_{31}NO_2P_2$ requires C, 72.1; H, 6.3; N, 2.8%). NMR: ¹H, δ 1.0 [d, 3 H, CH*Me*, ³*J*(HH) 7.1], 3.5 [*A*BX, 2 H, PCH₂, ²*J*(HH) 13.2, ²*J*(PH') 9.0], 3.55 (s, 3 H, OMe), 3.85 [ABX, 2 H, PCH₂, ²*J*(HH) 13.2, ²*J*(PH') 0.], 4.3 (m, 1 H, CH) and 7.2–7.5 (m, 20 H, Ph), ¹³C-{¹H}, δ 14.7 (s, Me), 50.9 (s, OMe), 54.9 [d, PCH₂, *J*(PC) 4.7], 58.5 [t, NCH, ³*J*(PC) 9.2 Hz] and 173.0 (s, CO₂Me); ³¹P-{¹H}, δ -26.6 (s).

(*iv*) $(Ph_2PCH_2)_2NCHCH_2CH_2CH_2CH_2C(Me)CMe_2 8$. The phosphonium salt **1a** (3.7 g, 13 mmol) with triethylamine (1.8 cm³, 13 mmol) and *endo*-(1*R*)-1,7,7-trimethylbicyclo-[2.2.1]heptan-2-ylamine (1.0 g, 6.5 mmol) gave compound **8** as a highly viscous oil (3.39 g, 9.5%) (Found: C, 78.5; H, 7.4; N, 2.4. C₃₆H₄₁NP₂ requires C, 78.7; H, 7.5; N, 2.6%). NMR spectra: ¹H, δ 0.55 (s, 3 H, Me), 0.67 (s, 3 H, Me), 0.81 (s, 3 H, Me), 0.9-1.05 (m, 2 H, CH₂), 1.37 [t, 1 H, CH, ³*J*(HH) 4.2], 1.4–1.55 (m, 2 H, CH₂), 3.0m (1 H, NCH), 3.85 [AB, 4 H, PCH₂, ²*J*(HH') 13.5] and 7.2–7.5 (m, 20 H, Ph); ¹³C-{¹H}, δ 16.1 (s, Me), 18.3 (s, Me), 19.7 (s, Me), 27.0 (s, CH₂), 28.3 (s, CH₂), 44.2 (s, CH), 48.9 (s, C), 49.5 (s, C), 57.4 [t, PCH₂, *J*(PC) 11.8] and 68.3 [t, NCH, ³*J*(PC) 7.2 Hz]; ³¹P-{¹H}, δ – 28.6.

(v) $(Ph_2PCH_2)_2NCH_2CH_2OH 9$. The phosphonium salt 1a (2.0 g, 7.0 mmol) with triethylamine (1 cm³, 7.0 mmol) and ethanolamine (0.21 cm³, 3.5 mmol) gave compound 9 as a highly viscous oil (1.54 g, 96%) (Found: C, 72.4; H, 6.6; N, 3.8. $C_{28}H_{29}NOP_2$ requires C, 73.5; H, 6.4; N, 3.0%). NMR: ¹H, δ 3.0 [t, 2 H, CH₂N, ³J(HH) 5.2], 3.27 [br s, exchange D₂O, 1 H, OH], 3.45 [t, 2 H, CH₂O, ³J(HH) 5.2], 3.85 [d, 4 H, PCH₂, ²J(PH) 3.3 Hz] and 7.2–8.0 (m, 20 H, Ph); ¹³C-{¹H}, δ 56.0 (s, CH₂O) and 57.7–58.2 (m, PCH₂, NCH₂); ³¹P-{¹H}, δ – 28.2 (s). (vi) $(Ph_2PCH_2)_2NCH_2CH=CH_2$ 10. The phosphonium salt

1a (2.0 g, 7 mmol) with triethylamine (1 cm³, 7 mmol) and allylamine (0.26 cm³, 3.5 mmol) gave compound 10 as a highly viscous oil (1.49 g, 94%) (Found: C, 75.8; H, 6.4; N, 3.5. $C_{29}H_{29}NP_2$ requires C, 76.8; H, 6.4; N, 3.1%). NMR: ¹H, δ 3.48– 3.57 (m, 2 H, NCH₂), 3.6 [d, 4 H, PCH₂, ²J(PH) 2.9], 5.04–5.1 (m, 2 H, =CH₂), 5.63–5.9 (m, 1 H, =CH) and 7.1–7.5 (m, 20 H, Ph); ³¹P-{¹H}, δ –28.8 (s).

Preparation of Bis(dicyclohexylphosphinomethyl)amines.—(i) [(C₆H₁₁)₂PCH₂]₂NCHMePh **5b**. Triethylamine (8.5 cm³, 84 mmol) was added to a stirred solution of compound **1b** (24.3 g, 82.5 mmol) in water-methanol (2:1, 100 cm³) followed by D(+)-α-methylbenzylamine (5.3 cm³, 46 mmol). The mixture became highly viscous and toluene (50 cm³) was added. The reaction was heated to reflux on an oil-bath for 45 min and then allowed to cool. The organic layer was decanted, dried over magnesium sulfate and filtered. Methanol was then added to the solution until it began to cloud. The solution was placed in a freezer at -15 °C. Large colourless crystals of **5b** were collected, washed with methanol, and dried *in vacuo* (12.5 g, 56%) (Found: C, 75.2; H, 10.8; N, 2.5. C₃₄H₅₇NP₂ requires C, 75.3; H, 10.5; N, 2.6%), m.p. 87-89 °C. NMR: ¹H, δ 1.0-1.85 (m, 47 H, 4C₆H₁₁ + Me), 2.6 [*A*BX, 2 H, PCH₂, ²*J*(HH) 12.9, ²*J*(PH) 3.82], 2.8 [*ABX*, 2 H, PCH₂, ²*J*(HH) 12.9, *J*(PH)0], 4.85 [m, 1 H, CH, ³*J*(HH) 6.9] and 7.15-7.4 (m, 5 H, Ph); ¹³C-{¹H}, δ 11.1 (s, Me), 47.9 (m, PCH₂) and 57.3 [t, CH, ³*J*(PC) 9.9 Hz] (cyclohexyl region omitted); ³¹P-{¹H}, δ -18.4 (s). Mass spectrum *m*/*z* 541 (*M*⁺, 541), 211 [100, (C₆H₁₁)₂PCH₂] and 105 [28%, CH(Me)Ph].

(*ii*) $[(C_6H_{11})_2PCH_2]_2NCHMeCO_2Et$ **6b**. Triethylamine (1.5 cm³, 20 mmol) was added to a stirred solution of compound **1b** (3.15 g, 10.7 mmol) in water-methanol (2:1, 30 cm³). L-Alanine ethyl ester hydrochloride (0.82 g, 5.3 mmol) and triethylamine (4.44 cm³, 6 mmol) in water-methanol (2:1, 30 cm³) were then added. The mixture was heated to 60 °C for 3 h and allowed to cool. The product was extracted with dichloromethane (30 cm³) and dried over magnesium sulfate. The solvent was then removed under reduced pressure to yield **6b** as a highly viscous oil (2.53 g, 90%) (Found: C, 70.1; H, 10.4; N, 2.9. $C_{31}H_{57}NO_2P_2$ requires C, 70.9; H, 10.9; N, 2.7%). NMR: ¹H, δ 1.1–1.9 (m, 50 H, 4C₆H₁₁ + 2Me), 2.7 [*A*BX, 2 H, PCH₂, ²*J*(HH) 14.0, ²*J*(PH) 6.8], 3.2 [*A*BX, 2 H, PCH₂, ²*J*(HH) 14.0, ²*J*(PH) 0.0], 4.3 [q, 2 H, OCH₂, ³*J*(HH) 7.1] and 4.5 (m, 1 H, CH); ¹³C-{¹H}, δ 13.5 (s, Me), 13.8 (s, Me), 29.5 [d, PCH, *J*(PC) 12.6], 47.7 (unresolved, PCH₂), 56.7 [t, NCH, ³*J*(PC) 10.2 Hz], 59.2 (s, OCH₂) and 172 (s, CO₂Et); ³¹P-{¹H}, δ – 18.2 (s).

Preparation of $[(C_8H_{14})PCH_2]_2NCHMePh 11.$ —Triethylamine was added to a solution of the phosphonium salt **3a** ([3.3.1] isomer) in water-methanol (2:1, 40 cm³). The compound $C_8H_{14}PCH_2OH$ was then extracted from the mixture with toluene (100 cm³). $D(+)-\alpha$ -Methylbenzylamine (1.5 cm³, 13 mmol) was added to the solution and the mixture refluxed for 3 h. The solvent was removed under reduced pressure. After 3 d at < 1 mmHg a white solid **11** formed (2.6 g, 48.5%) (Found: C, 72.5; H, 9.7; N, 3.2. $C_{26}H_{41}NP_2$ requires C, 72.7; H, 9.6; N, 3.4%). NMR: ¹H, δ 1.35 [d, 3 H, Me, ³J(HH) 6.8], 1.5–2.2 (m, 28 H, C_8H_{14}), 3.0 [ABX, 2 H, PCH₂, ²J(HH) 13.8, ²J(PH) 2.9], 3.25 [ABX, 2 H, PCH₂, ²J(HH) 13.8, ²J(PH) 2.0], 4.35 (m, 1 H, CH) and 7.15–7.4 (m, 5 H, Ph); ¹³C-{¹H}, δ 16.4 (s, Me), 48.0 [dd, PCH₂, ³J(PC) 13.1, ³J(PC) 9.5] and 60.3 [t, NCH, ${}^{3}J(PC)$ 9.7] (C₈H₁₄ moiety omitted); ${}^{31}P-{}^{1}H$, δ -46.6 (s).

Preparation of $(C_8H_{14})PCH_2NEt_2$ 12 (mixed isomers).—The phosphonium salts 3a and 3b ([4.2.1] and [3.3.1] isomers) (10 g, 45 mmol) were dissolved in water—methanol (2:1, 40 cm³) and a solution of Na₂SO₃ (11 g, 87 mmol) in water (25 cm³) followed by diethylamine (5 cm³, 97 mmol) were then added. The mixture was heated to about 60 °C for 1 h. On cooling the mixture separated into two layers. The organic layer was decanted and the product distilled under high vacuum to give a mixture of the isomers 12a and 12b as a clear liquid (8.65 g, 84%) (Found: C, 68.5; H, 11.7; N, 6.4. C₁₃H₂₆NP requires C, 68.7; H, 11.5; N, 6.2%), b.p. 124–127 °C at 2.5 mmHg. NMR: ¹H, δ 0.88 [t, 6 H, Me, ³J(HH) 7.1], 0.9 [t, 6 H, ³J(HH) 7.1], 1.3–2.3 [m, 28 H, 2C₈H₁₄], 2.4 [d, 2 H, PCH₂, ²J(PH) 2.2], 2.46 [dq, 4 H, CH₂CH₃, ³J(HH) 7.1, ⁴J(PH) 0.85] and 2.8 [d, PCH₂, ²J(PH) 1.5]; ¹³C-{¹H}, δ 12.7 (s, Me), 25.2 [d, PCH₂, J(PC) 10.5], 39.7 [d, PCH, J(PC) 11.8], 49.0 (m, PCH₂N), 50.5 [d, NCH₂, ³J(PC) 15.1] and 51.9 [d, NCH₂, ³J(PC) 9.1 Hz]; ³¹P-{¹H}, δ -9.9 (s, 1P, [3.3.1] isomer) and -49.0 (s, 1P, [4.2.1] isomer).

X-Ray Crystallography.—(a) [PPh₂(CH₂OH)₂]Cl 1a. A crystal with dimensions $0.6 \times 0.16 \times 0.12$ mm was glued to the end of a thin glass fibre using epoxy resin. Unit-cell parameters were determined by least-squares refinement of ω angles from 294 centred reflections with $15 < 2\theta < 30^{\circ}$.²⁷ Intensities of 1416 reflections in the range $7 < 2\theta < 50^{\circ}$ and 0 < h < 18, 0 < k < 8, 0 < l < 15 were measured on a Stoe STADI-2 Weissenberg diffractometer using an ω -scan technique and graphite-monochromated Mo-K α radiation. No crystal decay was detected from periodically measured check reflections. The data were corrected for Lorentz and polarisation effects, but not for absorption, and merged to give 1270 unique reflections with 994 having $I > 3\sigma(I)$.

Crystal data. $C_{14}H_{16}ClO_2P$, M = 282.71, orthorhombic, space group $Pna2_1, a = 14.615(10), b = 6.476(4), c = 15.126(2)$ Å, U = 1432(2) Å³, $Z = 4, \mu = 3.2$ cm⁻¹, λ (Mo-K α) = 0.7107 Å, $F(000) = 592, D_c = 1.31$ g cm⁻³, T = 293 K.

All calculations were performed on a VAX 8650 computer. The structure was solved by Patterson techniques using SHELXS 86.²⁸ All subsequent calculations were carried out using the computer program SHELX 76.29 Phenyl rings, including H atoms, were included as rigid groups with D_{6h} symmetry, C-C 1.395 and C-H 1.08 Å. The remaining hydrogen atoms were located on Fourier difference maps and refined as normal atoms. All non-hydrogen atoms were refined with anisotropic thermal parameters during the later cycles of refinement; hydrogen atoms were refined with group isotropic thermal parameters. Final cycles of least-squares refinement used a weighting scheme $w = 0.5/[\sigma^2(F_o) + gF_o^2], g = 0.0054$, and gave final residual indices of $R [=\Sigma(|F_o| - |F_c|)/\Sigma|F_o]] =$ 0.038 and $R' \{ = [\Sigma w(|F_0| - |F_c|)/\Sigma w|F_0] \} = 0.044$, the average shift/e.s.d. = 0.02 (maximum = 0.209). An analysis of the final weighting scheme over $|F_{o}|$ and $(\sin \theta)/\lambda$ was satisfactory. The final difference electron density had maximum and minimum values of +0.38 and -0.20 e Å⁻³ respectively. The atomic coordinates for the structure are given in Table 3.

(b) $[(C_6H_{11})_2PCH_2]_2NCHMePh$ **5b**. The analysis was carried out as for compound **1a**. Intensities were measured from a crystal with dimensions $0.64 \times 0.64 \times 0.20$ mm. Cell parameters were obtained by refinement of 334 centred reflections in the range $7 < 2\theta < 30^\circ$. A total of 7886 reflections was measured in the range $7 < 2\theta < 50^\circ$ with -34 < h < 34, 0 < k < 15, 0 < l < 11 which merged to 6435 unique reflections, 3597 having $I > 3\sigma(I)$.

Crystal data. $C_{34}H_{57}NP_2$, M = 541.8, orthorhombic, space group $P2_12_12_1$, a = 27.925(8), b = 12.531(6), c = 9.520(11) Å, U = 3331.3 Å³, Z = 4, $D_c = 1.08$ g cm⁻³, Mo-K α , $\lambda = 0.710$ 73 Å, $\mu = 1.2$ cm⁻¹, F(000) = 1192, T = 293 K.

Table 3 Fractional atomic coordinates for [PPh₂(CH₂OH)₂]⁺Cl⁻

Atom	x	у	z
Cl	0.280 89(9)	-0.188 66(21)	0.004 88(19)
Р	0.016 51(7)	0.108 87(17)	0.000 00(0)
O(1)	-0.15832(27)	0.116 3(8)	-0.048 7(3)
O(2)	-0.016 8(4)	-0.286 4(7)	0.047 6(4)
C(1)	-0.1026(3)	0.173 6(9)	0.022 4(4)
C(2)	0.022 6(4)	-0.170 1(8)	-0.0213(5)
C(21)	0.083 90(23)	0.175 7(5)	0.094 28(21)
C(22)	0.140 30(23)	0.349 3(5)	0.092 10(21)
C(23)	0.190 93(23)	0.404 2(5)	0.166 69(21)
C(24)	0.185 19(23)	0.285 6(5)	0.243 47(21)
C(25)	0.128 81(23)	0.111 9(5)	0.245 68(21)
C(26)	0.078 15(23)	0.057 0(5)	0.171 09(21)
C(31)	0.056 75(21)	0.236 0(5)	-0.096 98(20)
C(32)	0.147 32(21)	0.200 7(5)	-0.121 88(20)
C(33)	0.181 37(21)	0.287 3(5)	-0.199 69(20)
C(34)	0.124 85(21)	0.409 3(5)	-0.252 62(20)
C(35)	0.034 27(21)	0.444 6(5)	-0.227 71(20)
C(36)	0.000 24(21)	0.357 9(5)	-0.149 88(20)
H(1)	0.176(4)	0.006(11)	-0.033(4)
H(2)	-0.061(6)	-0.273(13)	0.036(6)
H(3)	-0.107(3)	0.314(8)	0.031(3)
H(4)	-0.108(5)	0.071(12)	0.084(6)
H(5)	0.092(4)	-0.213(8)	-0.024(3)
H(6)	-0.007(4)	-0.168(9)	-0.074(5)

Table 4 Fractional atomic coordinates for $[(C_6H_{11})_2PCH_2]_2$ -**NCHMePh**

Atom	x	у	Ζ
P(1)	0.134 30(4)	0.203 78(10)	0.238 26(15)
P(2)	0.114 15(5)	-0.12215(10)	0.038 63(15)
N	0.069 38(13)	0.058 0(3)	0.133 3(4)
C(1)	0.018 93(16)	0.024 2(4)	0.121 4(6)
C(2)	-0.01041(20)	0.087 7(5)	0.013 3(7)
C(3)	0.075 46(16)	0.174 2(4)	0.156 3(6)
C(4)	0.099 00(17)	0.021 3(3)	0.014 2(5)
C(11)	-0.003 50(12)	0.020 31(28)	0.265 3(3)
C(12)	-0.034 98(12)	0.098 55(28)	0.313 6(3)
C(13)	-0.052 96(12)	0.093 18(28)	0.450 1(3)
C(14)	-0.039 45(12)	0.009 54(28)	0.538 3(3)
C(15)	-0.007 96(12)	-0.068 70(28)	0.490 0(3)
C(16)	0.010 02(12)	-0.063 32(28)	0.353 5(3)
C(21)	0.145 93(16)	0.337 6(4)	0.163 5(6)
C(22)	0.162 45(19)	0.324 6(4)	0.013 9(6)
C(23)	0.177 59(20)	0.429 5(5)	-0.0532(7)
C(24)	0.215 28(22)	0.484 0(5)	0.028 8(9)
C(25)	0.199 47(24)	0.501 8(5)	0.179 8(9)
C(26)	0.184 13(18)	0.394 8(4)	0.251 3(7)
C(31)	0.116 60(20)	0.231 0(4)	0.423 4(6)
C(32)	0.078 90(22)	0.316 2(5)	0.449 4(6)
C(33)	0.070 58(29)	0.332 6(6)	0.607 6(8)
C(34)	0.059 37(29)	0.233 2(7)	0.678 6(8)
C(35)	0.096 0(3)	0.148 4(6)	0.651 1(7)
C(36)	0.105 11(25)	0.130 6(5)	0.497 0(6)
C(41)	0.168 36(18)	-0.103 5(5)	0.152 2(6)
C(42)	0.180 99(20)	-0.2001(5)	0.230 2(7)
C(43)	0.221 91(24)	-0.180 6(6)	0.335 0(7)
C(44)	0.265 33(21)	-0.134 1(6)	0.261 6(9)
C(45)	0.253 02(25)	-0.037 0(6)	0.187 4(8)
C(46)	0.212 33(19)	-0.0531(5)	0.079 8(7)
C(51)	0.136 34(17)	-0.154 8(4)	-0.1410(5)
C(52)	0.164 81(18)	-0.2593(4)	-0.1401(6)
C(53)	0.179 21(19)	-0.2947(5)	-0.285 3(6)
C(54)	0.136 43(21)	-0.3043(5)	-0.3818(7)
C(55)	$0.108\ 60(21)$	-0.200 8(5)	-0.386 1(6)
C(56)	0.092 94(17)	-0.1681(4)	-0.239 0(6)

All non-hydrogen atoms were refined anisotropically and all hydrogen atoms were included in calculated positions (C-H 1.08 Å), with a common fixed isotropic parameter. The final

R = 0.0611 and R' = 0.0595 (g = 0.002) for 147 parameters refined. Maximum and minimum peak heights in the final Fourier difference map were 0.47 and $-0.32 \text{ e} \text{ Å}^{-3}$. The atomic coordinates for the structure are given in Table 4.

Additional material available from the Cambridge Crystallographic Data Centre comprises H-atom coordinates, thermal parameters and remaining bond lengths and angles.

Acknowledgements

We thank the SERC and Albright and Wilson for financial support.

References

- 1 H. Coates and P. A. T. Hoye, Br. Pat., 842 593, 1960.
- 2 A. D. F. Toy and E. N. Walsh, Phosphorus Chemistry in Everyday Living, American Chemical Society, New York, 1987.
- 3 A. Granzow, J. Am. Chem. Soc., 1977, 99, 2648.
- 4 R.H. Carlson, Hooker Chemical Corporation, Br. Pat., 125 131, 1971.
- 5 J. Schapira, K. Mai Xuan and A. Askierzy, Fr. Pat., 2 157 108, 1973. 6 S. O. Grim, D. P. Shah and L. T. Matienzo, Inorg. Chem., 1980, 19,
- 2475.
- 7 K. Kellner, A. Tzshach, Z. Nagy-Magos and L. Marko, J. Organomet. Chem., 1980, 193, 307.
- 8 G. Markl and G. Yu. Jin., Tetrahedron Lett., 1980, 21, 1409.
- 9 J. Navech and J.-P. Majoral, Phosphorus Sulfur, Relat. Elem., 1983, 15, 51 and refs. therein.
- 10 G. Markl and G. Yu. Jin, Tetrahedron Lett., 1980, 21, 3461.
- 11 K. A. Petrov, V. A. Parshina and V. A. Gardamak, Zh. Obshch. Khim., 1961, **31**, 3186.
- 12 P. A. T. Hoye, Eur. Pat., 281 311, 1988.
- 13 Shell International Research, Netherlands Pat., 6 604 094, 1966; G. Elsner, G. Heymer and W. Stephan, Ger. Pat., 2 703 802, 1978; V. Winkle, J. L. Morris, R. Clarke and R. F. Mason, Ger. Pat., 1 909 620, 1969; Shell International Research, Netherlands Pat., 7 312 880, 1973; G. Elsner, G. Heymer and W. Stephan, Br. Pat., 1 561 874, 1980; Nippon Industrial Chemical Co. Ltd., Jap. Pat., 55 122 790, 1980; 55 122 791, 1980; N. Weferling, Ger. Pat., 3 235 787, 1984; N. Weferling and W. Klase, Ger Pat., 3 535 149, 1987; K. Imamura, S. Shimura and K. Takahashi, Jap. Pat., 62 022 791, 1987; 62 022 792, 1987; N. Weferling, Z. Anorg. Allg. Chem., 1987, **548**, 55; W. Klaeui and C. E. Song, Inorg. Chem., 1989, **28**, 3845; J. Svara and N. Weferling, Ger. Pat., 3 737 938, 1989.
- 14 International Tables for X-Ray Crystallography, Kynoch Press, Birmingham, 1962, vol. 3.
- 15 F. Allen, O. Kennard, D. G. Watson, L. Brammer, A. G. Orpen and R. Taylor, J. Chem. Soc., Perkin Trans. 2, 1987, S1; J. Chem. Soc., Dalton Trans., 1989, S1.
- 16 G. Markl and G. Yu. Jin., Tetrahedron Lett., 1981, 22, 229.
- 17 K. Kellner, A. Tzshach, W. Hanker, L. Marko and Z. Nagy-Magos, G.D.R. Pat., DD 1 496 692, 1981.
- 18 K. Kellner, A. Tzshach, W. Hanker, L. Marko and Z. Nagy-Magos, J. Organomet. Chem., 1984, 268, 175.
- 19 L. D. Quin, The Heterocyclic Chemistry of Phosphorus, Wiley Interscience, New York, 1981.
- 20 D. H. Williams and I. Fleming, Spectroscopic Methods in Organic Chemistry, 3rd edn., McGraw-Hill, London, 1980.
- 21 A. W. Frank and G. L. Drake, J. Org. Chem., 1973, 37, 2752.
- 22 L. N. Ferguson, The Modern Structural Theory of Organic Chemistry, Prentice-Hall, Englewood Cliffs, NJ, 1963.
- 23 K. Jones, in Comprehensive Inorganic Chemistry, eds. J. C. Bailar, H. J. Emeleus, R. S. Nyholm and A. F. Trotman-Dickenson, Pergamon, Oxford, 1973, vol. 2, p. 147.
- 24 C. A. Tolman, Chem. Rev., 1977, 313, 77. 25 A. F. Wells, Structural Inorganic Chemistry, 3rd edn., Oxford University Press, London, 1962.
- 26 T. Glonek and J. R. van Wazer, J. Magn. Reson., 1974, 13, 390.
- 27 W. Clegg and G. M. Sheldrick, Z. Kristallogr., 1984, 167, 23. 28 G. M. Sheldrick, SHELXS 86, Program for crystal structure determination, University of Göttingen, 1986.
- 29 G. M. Sheldrick, SHELX 76, Program for crystal structure determination, University of Cambridge, 1976.

Received 16th February 1993; Paper 3/00935A