

Nickel catalyzed Grob fragmentation: ω -dienyl aldehydes synthesis†

Masahiko Mori,^b Masanari Kimura,^b Yushi Takahashi^b and Yoshinao Tamaru^{*a}

Received (in Cambridge, UK) 17th July 2006, Accepted 27th July 2006

First published as an Advance Article on the web 1st September 2006

DOI: 10.1039/b610164j

With a proper choice of phosphane ligands, a Ni(cod)₂-phosphane catalyst promotes decarboxylative ring-opening reaction of a wide structural variety of cyclic carbonates **1** to give ω -dienyl aldehydes **2** in good yields.

The Grob fragmentation reaction is a very powerful tool for the construction of desired molecules.¹ However, harsh reaction conditions, either strongly basic or acidic and/or high temperatures, have limited its wide use.

In the last decade, C–C bond cleavage reactions catalyzed or mediated by transition metals have gained significant attention,^{2,3} because they can be performed under much milder reaction conditions and provide new organic transformations that are otherwise difficult to achieve.

In 1997, we revealed that many palladium species, especially Pd₂[(dba)₃]₂·CHCl₃ [dba = dibenzylideneacetone] in acetonitrile, catalytically promoted the Grob type decarboxylative ring-opening reaction of cyclic carbonates **1** into ω -dienyl aldehydes **2** (Scheme 1).⁴ The reaction proceeds at room temperature and is rationalized by a facile β -C elimination of an oxapalladacyclopentane intermediate **I** (M = Pd²⁺). On the other hand, we have demonstrated that, Ni(acac)₂ in the presence of triethylborane⁵ or diethylzinc⁶ catalytically promotes the reductive cyclization (homoallylation) of **2** and provides bis-homoallyl alcohols **3** in good yields (Scheme 1).⁷ This formally reverse reaction was supposed to proceed through oxidative addition of a nickel(0) species across the diene and aldehyde moieties of **2**, followed by the reduction of an oxanickellacyclopentane intermediate **I** (M = Ni²⁺) with Et₃B or Et₂Zn. Recently, the intermediacy of **I** (M = Ni²⁺) was supported by Ogoshi and Kurosawa, who have succeeded in

the isolation and X-ray structure characterization of **I** (without a tether) prepared by the stoichiometric reaction of Ni(cod)₂ [cod = 1,5-cyclooctadiene], a diene and an aldehyde in the presence of a monodentate phosphane ligand.⁸

These observations suggest that the equilibrium between **I** and M⁰ is strongly affected by the nature of the transition metal; nickel tends to shift the equilibrium to the side of **I**, while palladium works oppositely to drive the equilibrium to the side of **2** and Pd⁰. This contrasting behaviour between nickel and palladium might be rationalized in part by the fact that nickel forms a stronger metal–oxygen bond than palladium, e.g., $\Delta H^\circ(\text{Ni–O}) = 60.5 \text{ kcal mol}^{-1}$ and $\Delta H^\circ(\text{Pd–O}) = 58.9 \text{ kcal mol}^{-1}$ for *cis*-(PH₃)₂M(OMe)Me.⁹


Here we disclose that Ni(cod)₂, despite the above-mentioned unfavorable reaction features, nicely catalyzes the decarboxylative fragmentation reaction of a wide structural variety of **1** and furnishes ω -dienyl aldehydes **2** in comparable or better yields than palladium species. However, unlike the palladium catalysis, which worked equally well irrespective of the structure types of **1**,⁴ a nickel species required a fine tuning of the reaction conditions, especially an appropriate choice of phosphane ligands, depending on the structure types of **1**.

As summarized in Table 1, carbonates characterized by their ring strain (**1a–c**) and/or torsional strain (**1d,e**) smoothly underwent the fragmentation reaction in the presence of Ni(cod)₂ (10 mol%) and PPh₃ (40 mol%) in acetonitrile at room temperature and furnished **2** in comparable yields or even in better yields than those obtained by Pd₂(dba)₃·CHCl₃.

In sharp contrast to these, as shown in run 1, Table 2, a strain-free six-membered carbonate **1f** was robust and remained unchanged under usual conditions (at 25 °C) and even under forcing conditions undertaken at a higher temperature (81 °C). This was rather surprising in view of a smooth fragmentation of **1f** with Pd₂[(dba)₃]₂·CHCl₃ at room temperature yielding **2f** in 77% isolated yield (*E* : *Z* = 10 : 1, footnote b, Table 2). Neither the monodentate trialkyl- nor triarylphosphane ligands were effective (runs 1–3). Among bidentate phosphane ligands, only those with bite angles of ca. 95–105° effectively promoted the fragmentation reaction (runs 6 and 7, Table 2).

Similar results were obtained for other cyclohexane derivatives **1j** (runs 7 and 8, Table 3) and **1k** (runs 9 and 10); with Ni(cod)₂/PPh₃, they remained intact even after long heating at the refluxing temperature of acetonitrile, while with Ni(cod)₂/DPPF, they smoothly underwent the ring-opening reaction at room temperature and provided **2j** and **2k** in much better yields than those with Pd₂[(dba)₃]₂·CHCl₃.

The large bite angle of DPPF (99–105°) might be suited to the tetrahedral geometry of a Ni⁰ species **III** and not to a square planar geometry of a Ni²⁺ species **II** (Scheme 2).¹⁰ This destabilization of



Scheme 1 Nickel (or palladium) catalyzed fragmentation of **1** to **2** and cyclization of **2** to **3** via a common intermediate **I** (M = Ni or Pd).

^aDepartment of Applied Chemistry, Faculty of Engineering, Nagasaki University, 1-14 Bunkyo, Nagasaki, 852-8521, Japan.
E-mail: tamaru@net.nagasaki-u.ac.jp

^bGraduate School of Science and Technology, Nagasaki University, 1-14 Bunkyo, Nagasaki, 852-8521, Japan

† Electronic supplementary information (ESI) available: Experimental procedures and analytical and spectroscopic data for **1a–m**¹³ and **2a–m**. See DOI: 10.1039/b610164j

Table 1 Synthesis of ω -dienyl aldehydes **2** by Ni-catalyzed decarboxylative ring-opening reaction of **1**^a

Run	1	<i>t</i> /h	Product 2	Yield ^b (%)	
				Ni	Pd
1		2		75 (1 : 2.4)	65 (1 : 4.4) ^c
2		6		78 (2 : 1)	87 (1.7 : 1) ^c
3		0.5		96 (only <i>E</i>)	70 (only <i>E</i>) ^d
4		24		97 (1 : 1)	91 (1 : 1) ^c
5		24		85 (1 : 1)	85 (1 : 3) ^d

^a Reaction conditions: **1** (1 mmol), Ni(cod)₂ (0.1 mmol) and PPh₃ (0.4 mmol) in acetonitrile (5 mL) at room temperature under N₂. ^b *E*/*Z* Ratio of products in parenthesis. ^c Yield obtained using Pd₂(dba)₃·CHCl₃. ^d Ref. 4a. ^e A mixture of stereoisomers (see ESI).

II and stabilization of **III** by bidentate ligands with wide bite angles might cooperate to facilitate the fragmentation of **II** and shift the








Table 2 Effects of phosphanes on the fragmentation of **1f**

Reaction scheme showing the conversion of compound **1f** to compound **2f** using 10 mol% $\text{Ni}(\text{cod})_2$.

Run	Phosphane/ solvent ^c	<i>T</i> /°C, <i>t</i> /h	Yield (<i>E</i> : <i>Z</i>) ^d (%)	Bite angle/ ^e
1	PPh_3/AN	81, 24	NR	—
2	<i>n</i> - $\text{Bu}_3\text{P}/\text{AN}$	81, 24	NR	—
3	$\text{Cy}_3\text{P}/\text{AN}$	81, 24	NR	—
4	DPPE/THF	25, 48	NR	85
5	DPPP/THF	25, 48	NR	90–95
6	DPPB/THF	25, 4	76 (>20 : 1)	94–99
7	DPPF/THF	25, 4	78 (18 : 1)	99–105

^a Reaction conditions: **1f** (1 mmol, a mixture of stereoisomers, see ESI), Ni(cod)₂ (0.1 mmol) and a phosphane (0.4 mmol for a monodentate and 0.2 mmol for a bidentate ligand) in a solvent (5 mL) under N₂. ^b **2f** in 77% (*E* : *Z* = 10 : 1) using Pd₂(dba)₃·CHCl₃ as a catalyst. ^c AN = acetonitrile, THF = tetrahydrofuran, Cy = cyclohexyl, DPPE = 1,2-bis(diphenylphosphino)ethane, DPPP = 1,3-bis(diphenylphosphino)propane, DPPB = 1,4-bis(diphenylphosphino)butane, DPPF = 1,1'-bis(diphenylphosphino)ferrocene. ^d Yields refer to the isolated spectroscopically homogeneous **2f**.


Table 3 Acceleration of the Ni-catalyzed fragmentation of **1** with bidentate phosphane ligands with wide bite angles^a

			Yield (<i>E</i> / <i>Z</i>) (%)	
Run	Phosphane/solvent ^b	<i>T</i> /°C, <i>t</i> /h	Ni	Pd
				
1	PPh ₃ /AN	81, 24	NR	
2	DPPF/THF	25, 45	91 (only <i>E</i>)	82 (only <i>E</i>) ^c
				
3	PPh ₃ /AN	81, 24	NR	
4	DPPF/THF	25, 3	81 (only <i>E,E</i>)	93 (only <i>E,E</i>) ^c
				
5	PPh ₃ /AN	81, 24	24 (only <i>E</i>) ^c	
6	DPPF/ether	25, 9	98 (only <i>E</i>)	55 (only <i>E</i>) ^d
				
7	PPh ₃ /AN	81, 24	NR	
8	DPPF/THF	25, 4	84 (3 : 1)	42 (2 : 1) ^d
				
9	PPh ₃ /AN	81, 30	NR	
10	DPPF/THF	25, 3	94 (4 : 1)	60 (1 : 1) ^d
				
11	PPh ₃ /AN	81, 15	40 (1 : 5)	
12	DPPF/Et ₂ O	25, 2	94 (2 : 1)	54 (1 : 4) ^c
				
13	PPh ₃ /AN	81, 40	NR	
14	DPPF/THF	25, 72	43 (only <i>E</i>)	
15	DPPPent/THF	25, 30	100 (only <i>E</i>)	63 (only <i>E</i>) ^c


^a Reaction conditions: **1** (1 mmol), Ni(cod)₂ (0.1 mmol) and a phosphane (0.4 mmol for a monodentate and 0.2 mmol for a bidentate ligand) in a solvent (5 mL) under N₂. ^b DPPF = 1,1'-bis(diphenylphosphino)ferrocene. ^c Ref. 4a. ^d Yield obtained using Pd₂(dba)₃·CHCl₃. ^e 59% conversion (59% isolated yield based on conversion). ^f A mixture of stereoisomers (see ESI).

equilibrium to the side of a mixture of **2** and **III**. The complex **III**, however, might be reactive enough toward oxidative addition to allylic carbonates **1** and might be able to supply **II**, hence accomplishing the catalytic cycle.

In Table 3 are summarized the carbonates **1** that are either unreactive (NR) or provide **2** albeit in low yields with the catalyst Ni(cod)₂/PPh₃. Comparison of the data in Tables 1 and 3 clearly



Scheme 2 Acceleration of β -C elimination by bidentate ligands with wide bite angles (θ).



Scheme 3 Increase in steric repulsion between P1 and vinyl moiety in a transition state V, placing all the P1, P2, Ni, C5, C4, C3 and O atoms in the same plane.

indicates that both the C1'- and C2'-substituents on the vinyl group of **1** seriously retard the reaction. For example, while **1b** undergoes fragmentation at room temperature with $\text{Ni}(\text{cod})_2/\text{PPh}_3$ (run 2, Table 1), the C1'-Me and C2'-Me derivatives, **1g** and **1h**, remain unchanged under the same or even under forcing conditions (runs 1 and 3, Table 3). The fragmentation of **1g** and **1h** was successfully achieved by making use of the $\text{Ni}(\text{cod})_2/\text{DPPF}$ catalytic system (runs 2 and 4). More contrasting results were observed between the reactions of **1e** (run 5, Table 1) and **1m** (runs 13–15). For the latter substrate, even the $\text{Ni}(\text{cod})_2/\text{DPPF}$ catalytic system was not effective enough, and $\text{Ni}(\text{cod})_2/\text{DPPent}$ [1,5-bis(diphenylphosphino)pentane] turned out to work much better.¹¹

The retardation of the reaction by the C1' and C2' substituents, especially by the one on C1' (cf, the reaction times in runs 2 and 4, Table 3), might be ascribed to the increase in the steric repulsion between P1 and the vinyl group moiety that arises when **IV** approaches a transition state **V** (Scheme 3), since this process is accompanied by the decrease in the absolute value of the dihedral angle Ni–C5–C4–C3, and hence the decrease in the dihedral angle P1–Ni–C5–C1'. That is, there seems to exist an interesting dichotomy in the role of bidentate phosphane ligands with wide bite angles; electronically they accelerate the fragmentation reaction by stabilizing **III** and destabilizing **II** (Scheme 2),¹² while sterically they retard the reaction preventing the Ni–C5 bond from taking coplanar conformation with the C3–C4 bond.

Cyclic carbonates **1a–m** were prepared readily according to the following three-step procedures (see, ESI†): (1) cross-aldol of α,β -unsaturated aldehyde with lithium enolate of cyclic ketone, (2) LiAlH_4 reduction to diol, (3) cyclic carbonation with methyl chloroformate/triethylamine.

Finally, it may be worth noting that the present nickel catalyzed degradation not only records better yields, but also shows higher (*E*)-selectivity than the palladium catalyzed reaction. For example, a mixture of (*E*)- and (*Z*)-**2k** was obtained in a ratio of 1 : 1 in 60% isolated yield under the palladium catalysis, while under the nickel catalysis the same product was obtained in a ratio of 4 : 1 in 94% yield (run 10, Table 3). More contrasting results were observed for the reaction of **1l**, where the yield was doubled and the stereoselectivity was reversed from the (*Z*)-selective one to the (*E*)-selective one (run 12, Table 3).

In summary, the $\text{Ni}(\text{cod})_2/\text{PPh}_3$ or $\text{Ni}(\text{cod})_2/\text{DPPF}$ catalytic system promotes a novel decarboxylative fragmentation of a wide

structural variety of cyclic carbonates **1** to furnish ω -dienyl aldehydes **2**, where the phosphane ligands play a pivotal role to shift the equilibrium between an intermediate **I** and an ω -dienyl aldehyde **2** to the product side. The reaction shows better yields and higher (*E*)-selectivity than the original palladium-catalyzed reaction reported previously by the authors.^{4a} The ready availability of the starting materials **1** and the importance of the products **2** as strategic intermediates for natural product synthesis may augment the utility of the present reaction as a synthetic method. Furthermore, the reaction provides examples that explicitly demonstrate the salient difference in catalytic reactivity between a nickel(0) and a palladium(0) species.

Financial support from the Ministry of Education, Culture, Sports, Science and Technology, Japanese Government [Grant-in-Aid for Scientific Research (B) 16350058 and Priority Areas 17035065] is gratefully acknowledged.

Notes and references

- G. Mehta and R. S. Kumaran, *Tetrahedron Lett.*, 2005, **46**, 8831–8835; T. Constantieux and J. Rodriguez, *Sci. Synth.*, 2004, **26**, 413–462, and references therein.
- β -C Elimination of strained cyclobutanes and cyclopropanes: M. Murakami, S. Ashida and T. Matsuda, *J. Am. Chem. Soc.*, 2006, **128**, 2166–2167; M. Murakami, S. Ashida and T. Matsuda, *J. Am. Chem. Soc.*, 2005, **127**, 6932–6933; T. Nishimura and S. Uemura, *Synlett*, 2004, 201–216, and references therein.
- Norbornene-mediated reactions: C. Bressy, D. Alberico and M. Lautens, *J. Am. Chem. Soc.*, 2005, **127**, 13148–13149; M. Catellani, E. Motti, F. Faccini and R. Ferraccioli, *Pure Appl. Chem.*, 2005, **77**, 1243–1248.
- (a) H. Harayama, T. Kuroki, M. Kimura, S. Tanaka and Y. Tamaru, *Angew. Chem., Int. Ed. Engl.*, 1997, **36**, 2352–2354; see also: H. Harayama, M. Kimura, S. Tanaka and Y. Tamaru, *Tetrahedron Lett.*, 1998, **39**, 8475–8478.
- M. Kimura, A. Ezoe, K. Shibata and Y. Tamaru, *J. Am. Chem. Soc.*, 1998, **120**, 4033–4034; see also: M. Kimura, A. Ezoe, S. Tanaka and Y. Tamaru, *Angew. Chem., Int. Ed.*, 2001, **40**, 3600–3602; K. Shibata, M. Kimura, M. Shimizu and Y. Tamaru, *Org. Lett.*, 2001, **3**, 2181–2183.
- M. Kimura, A. Miyachi, K. Kojima, S. Tanaka and Y. Tamaru, *J. Am. Chem. Soc.*, 2004, **126**, 14360–14361; M. Kimura, H. Fujimatsu, A. Ezoe, K. Shibata, M. Shimizu, S. Matsumoto and Y. Tamaru, *Angew. Chem., Int. Ed.*, 1999, **38**, 397–400.
- Closely related reactions involving **I** as an intermediate: M. Kimura, A. Ezoe, M. Mori and Y. Tamaru, *J. Am. Chem. Soc.*, 2005, **127**, 201–209; A. Ezoe, M. Kimura, I. Inoue, M. Mori and Y. Tamaru, *Angew. Chem., Int. Ed.*, 2002, **41**, 2784–2786; M. Kimura, S. Matsuo, K. Shibata and Y. Tamaru, *Angew. Chem., Int. Ed.*, 1999, **38**, 3386–3388.
- S. Ogoshi, K. Tonomori, M. Oka and H. Kurosawa, *J. Am. Chem. Soc.*, 2006, **128**, 7077–7086.
- S. A. Macgregor and G. W. Neave, *Organometallics*, 2004, **23**, 891–899.
- E. Burello, P. Marion, J.-C. Galland, A. Chamard and G. Rothenberg, *Adv. Synth. Catal.*, 2005, **347**, 803–810.
- Bite angles (103.55–103.86°) of DPPent for CuI complexes determined by X-ray crystallography: Effendy, C. D. Nicola, M. Fianchini, C. Pettinari, B. W. Skelton, N. Somers and A. H. White, *Inorg. Chim. Acta*, 2005, **358**, 763–795; bite angle (104.37°) for a palladium complex obtained by DFT-optimization: R. Kuwano and M. Yokogi, *Org. Lett.*, 2005, **7**, 945–947.
- Three different fragmentation modes of nickellacyclopentanes: R. H. Grubbs, A. Miyashita, M. Liu and P. Burk, *J. Am. Chem. Soc.*, 1978, **100**, 2418–2425.
- Crystal data for **1c**: $\text{C}_{14}\text{H}_{20}\text{O}_3$, $M = 236.31$, orthorhombic, space group $P2_12_12_1$, $a = 7.5620(4)$, $b = 10.2674(8)$, $c = 16.5730(11)$ Å, $U = 1286.76(15)$ Å³, $T = 296.1$ K, $Z = 4$, $\mu(\text{Mo-K}\alpha) = 0.7107$ mm^{−1}, 1551 reflections measured, 985 unique ($R_{\text{int}} = 0.025$), $wR(F^2) = 0.1596$ (all data). For crystallographic data in CIF or other electronic format see DOI: 10.1039/b610164j.