A SIMPLE, HIGHLY VERSATILE SYNTHESIS OF CYCLOPROPYL PHENYL SULFIDES

Kazuhiko TANAKA^{*}, Hideki UNEME, Syuichi MATSUI, Rikuhei TANIKAGA, and Aritsune KAJI Department of Chemistry, Faculty of Science, Kyoto University, Sakyo-ku, Kyoto 606

1,3-Bis(phenylthio)propane reacted readily with 2-2.2 equiv. of n-butyllithium in THF at 0°C to afford 1-lithiocyclopropyl phenyl sulfide. Treatment of the substituted 1,3-bis(phenylthio)propanes with 1.1 equiv. of n-butyllithium afforded the corresponding cyclopropanes in good yields.

1-Lithiocyclopropyl phenyl sulfide is a useful reagent for the synthesis of cyclobutanone derivatives,¹⁾ γ -ketosulfides, β -bromoethyl vinyl sulfides.²⁾ Recently, Cohen found an efficient route to the lithium derivatives of cyclopropanes.³⁾ We now wish to report an extremely facile method for the preparation of l-lithiocyclopropyl phenyl sulfide. Treatment of 1,3-bis(phenylthio)propane⁴⁾ (I) with 2-2.2 equiv. of n-butyllithium in THF at 0°C followed by addition of electrophilic trapping reagents gave the cyclopropanes (II) in good yields (Table 1).

 $PhSCH_2CH_2CH_2SPh \xrightarrow{1) n-BuLi} \qquad \qquad \swarrow_{E}^{SPh} + PhSH$

Reaction of a variety of alkyl substituted 1,3-bis(phenylthio)propanes with 1.1 equiv. of n-butyllithium gave the corresponding cyclopropanes in good isolated yields.

(I) mmol	n-BuLi mmol	Electrophile ^{a)}	Product (II) ^{b)}	Yield (%)	b.p. °C/Torr
			SPh		
20	44	D ₂ 0	D	65	84-87/3.5
			SPh		
20	44	4-Heptanone	С (ОН) [(СН ₂) 2СН ₃] 2	71	140/0.9
2.0		D	SPh	67	_
20	44	Benzaldenyde	SPD	67	C
20	44	Cyclohexanone	$\mathbf{N}_{\mathbf{C}}(\mathbf{OH})$ (CH ₂)	60	с
		-1	SPh		-
20	44	3-Pentanone	С (ОН) (CH ₂ CH ₃)	69	126-128/0.9
			SPh		
20	40	3-Pentanone ^{d)}	→ с (он) (сн ₂ сн ₃) 2	85	126-128/0.9
		(b	SPh		
20	40	Heptanal"	СН (ОН) (СН ₂) ₅ СН ₃	59	151-152/0.8
20	40	2-Hexanone ^{d)}	ССН ₃ (он) (СН ₂) ₃ СН ₃	65	129/0.6

Table I. Preparation of cyclopropanes (II) from 1,3-bis(phenylthio)propane (I)

a) 10 mmol scale. b) Isolated by distillation or chromatography on silica gel and adequately characterized by analytical and spectral data. c) Oil. d) CuI (20 mmol) was added.

This approach can be extended to the preparation of cyclopropanol. Thus, reaction of 2-hydroxy-1,3-bis(phenylthio)propane with 2.2 equiv. of n-butyllithium followed by quenching with saturated aqueous ammonium chloride solution gave 2-hydroxycyclopropyl phenyl sulfide.

The experimental simplicity and use of readily available 1,3-bis(phenylthio)propanes make the present method highly advantageous for the synthesis of cyclopropane derivatives.

References

- a) B. M. Trost, Acc. Chem. Res., <u>7</u>, 85 (1974). b) B. M. Trost, D. Keeley, and M. J. Bogdanowicz, J. Am. Chem. Soc., <u>95</u>, 3068 (1973). c) B. M. Trost and D. E. Keeley, *ibid.*, <u>96</u>, 1252 (1974). d) B. M. Trost and D. E. Keeley, *ibid.*, <u>98</u>, 248 (1976). e) B. M. Trost, D. E. Keeley, H. C. Arndt, and M. J. Bogdanowicz, *ibid.*, <u>99</u>, 3088 (1977).
- 2) a) R. D. Miller and D. R. McKean, Tetrahedron Lett., <u>1979</u>, 583. b) R. D. Miller, D. R. McKean, and D. Kaufmann, *ibid.*, <u>1979</u>, 587. c) S. Halazy and A. Krief, *ibid.*, <u>1979</u>, 4233.
- 3) a) T. Cohen and W. M. Daniewski, Tetrahedron Lett., <u>1978</u>, 2991. b) T. Cohen,
 W. M. Daniewski, and R. B. Weisenfeld, *ibid.*, <u>1978</u>, 4665.
- 4) 1,3-Bis(phenylthio)propane was prepared in 93% yield from 1,3-dibromopropane and sodium benzenethiolate.⁵)
- 5) F. G. Mann and D. Purdie, J. Chem. Soc., 1935, 1549.