

## 16. Nucleosides and Nucleotides

Part 25<sup>1)</sup>

### Synthesis of a Protected 1-(2'-Deoxy- $\beta$ -D-ribofuranosyl)-1*H*-benzimidazole 3'-Phosphate

by Christos Papageorgiou and Christoph Tamm\*

Institut für Organische Chemie der Universität Basel, St. Johannis-Ring 19, CH-4056 Basel

(24.X.86)

---

1-(2'-Deoxy-5'-*O*-dimethoxytrityl- $\beta$ -D-ribofuranosyl)-1*H*-benzimidazole 3'-[(*p*-chlorophenyl) (2-cyanoethyl) phosphate] (**6**) has been synthesized from 1-( $\beta$ -D-ribofuranosyl)-1*H*-benzimidazole (**3b**) using regiospecific 2'-de-oxygenation. The latter compound was obtained by glycosylation of benzimidazole with the D-ribose derivative **2** leading exclusively of the  $\beta$ -D-anomer.

---

**Introduction.** – For our continuing efforts towards the study of the specificity of DNA polymerase [2], we required an oligonucleotide sequence with a  $\beta$ -D-deoxyribonucleoside in which an unnatural base could be incorporated. The 1-(2'-deoxy- $\beta$ -D-ribofuranosyl)-1*H*-benzimidazole (Drb) was chosen for this purpose since it has been shown that Drb are simulating purine-nucleoside analogs [3]. For example, the inhibition of influenza-B virus by 5,6-dichloro-1-(2'-deoxy- $\beta$ -D-ribofuranosyl)-1*H*-benzimidazole is reversed by adenosine [4]. It has been suggested that this compound interferes with preliminary synthesis of ribonucleic acid.


**Results.** – Because of the biochemical interest of Drb, a number of syntheses of this compound have been reported, all starting from 2-deoxy-D-ribose derivatives giving quite low yields of the  $\beta$ -D-anomer [5]. To overcome this problem, a strategy based on glycosylation of benzimidazole with D-ribose derivatives followed by regiospecific 2'-deoxygenation was adopted. It was anticipated that the presence of a suitable protected 2'-hydroxy group could lead to stereochemical control giving only the  $\beta$ -D-anomer. The resulting nucleoside could then be readily transformed to the corresponding nucleotide **6** needed for the synthesis of the sequence.

It is known that fusion between benzimidazole and tetra-*O*-acetyl-D-ribose [6] or the reaction between silylated benzimidazole and tri-*O*-benzoyl-D-ribofuranosyl bromide [7] both yielded mixtures enriched in the  $\beta$ -D-anomer. In order to avoid this problem, the condensation of chloromercurio-benzimidazole **1** with 2,3,5-tri-*O*-benzoyl-D-ribofuranosyl chloride (**2**) was studied (*Scheme*). Analogous glycosylations giving exclusively the  $\beta$ -D-anomers had already been reported [8].

---

<sup>1)</sup> Part 24: [1].

## Scheme



In our case, the reaction proceeded with good yield in refluxing toluene to give only the anomer **3a**. The  $^1\text{H}$ -NMR spectrum of this compound showed a  $d$  at 6.42 ppm ( $J = 5.4$ ) representing the anomeric proton at  $\text{C}(1')$ . This signal corresponds to the upfield resonance for  $\text{H}-\text{C}(1')$  in the spectrum of the  $\alpha$ -D/ $\beta$ -D mixture prepared according to [7]. Thus, the configuration at the glycosidic bond of **3a** is established as  $\beta$ -D.

Debenzoylation of the nucleoside **3a** with NaOMe afforded **3b**, which was then subjected to the four-step procedure for transforming a ribonucleoside to the 2'-deoxy analog [9]. Selective protection of the 5'- and 3'-hydroxy functions was achieved with 1,3-dichloro-1,1,3,3-tetraisopropyldisiloxane ( $[(i\text{-Pr})_2\text{SiCl}_2\text{O}]$ ) in pyridine yielding **4a**. The remaining 2'-hydroxy group was first transformed to a thiocarbonate leading to **4b** which was homolytically deoxygenated with  $\text{Bu}_3\text{SnH}$  to give **5a**. Removal of the protecting groups with  $\text{Bu}_4\text{NF}$  in THF afforded the 2'-deoxy compound **5b**. The deoxygenation sequence proceeded with an overall yield of 43 %. Confirmation of the  $\beta$ -D-configuration of **5b** was again obtained by its  $^1\text{H}$ -NMR. The signal for  $\text{H}-\text{C}(1')$  appeared as a pseudo- $t$  at 6.35 ppm. Two signals exchangeable with  $\text{D}_2\text{O}$ , a  $t$  and a  $d$ , assured the presence of a primary and a secondary alcohol.

The desired fully protected deoxynucleoside 3'-phosphate was obtained from **5b** in two steps with 65% overall yield. First, the 5'-hydroxy function was protected as a  $(\text{MeO})_2\text{Tr}$  ether making the 3'-OH available for phosphorylation. This latter operation was effected with the monofunctional ( $p$ -chlorophenyl) (2-cyanoethyl) phosphochloridate [10] to give the desired protected nucleoside **6**.

The investigations were supported by the Swiss National Science Foundation.

## Experimental Part

**General.** All reactions were routinely run in a flame-dried (*in vacuo*) round-bottom flask under dry Ar. Toluene was dried over Na. Other solvents and reagents were reagent-grade and stored over 4-Å molecular sieves. TLC: precoated silica-gel plates (60 GF 254 Merck). Column chromatography: silica gel C-560 from *Chemische Fabrik Uetikon*. M.p.: *Will-Wetghar* apparatus; uncorrected. Optical rotations: *Perkin-Elmer-141* polarimeter; cell length, 10 cm, 1 ml capacity; Na light.  $^1\text{H-NMR}$  spectra: *Varian EM-390*, chemical shifts in ppm downfield from tetramethylsilane (TMS) as internal standard, coupling constants  $J$  in Hz. The elemental analyses were performed by the Microanalytical Laboratory of our Institute.

**1-(2',3',5'-Tri-O-benzoyl- $\beta$ -D-ribofuranosyl)-1H-benzimidazole (3a).** To a stirred suspension of 7.7 g (22 mmol) of 1-chloromercuro-1H-benzimidazole (1) [8] in 100 ml of toluene, a soln. of 10.45 g (20 mmol) of tetra-O-benzoyl-D-ribofuranosyl chloride [8] in 30 ml of toluene was added. The heterogeneous mixture was refluxed for 2 h under continuous stirring and then allowed to cool. After filtration of the insoluble material, the filtrate was first washed with a 30% aq. KI soln., then with  $\text{H}_2\text{O}$ . The org. layer was dried over  $\text{MgSO}_4$  and concentrated to a sirup. Chromatography over silica gel using  $\text{CHCl}_3$  afforded 6.1 g (55%) of **3a** as a white foam.  $[\alpha]_D^{20} = -66.1^\circ$  ( $c = 1.0$ ,  $\text{CH}_2\text{Cl}_2$ ).  $^1\text{H-NMR}$  ( $\text{CDCl}_3$ ): 8.3–7.1 ( $m$ , 5 arom. H); 6.43 ( $d$ ,  $J(1',2') = 5.4$ ,  $\text{H-C}(1')$ ); 6.2–5.9 ( $m$ ,  $\text{H-C}(2')$ ,  $\text{H-C}(3')$ ); 5.0–4.7 ( $m$ ,  $\text{H-C}(4')$ , 2  $\text{H-C}(5')$ ).

**1-[3',5'-O-(1,1,3,3-Tetraisopropylidisiloxane-1,3-diyl)- $\beta$ -D-ribofuranosyl]-1H-benzimidazole (4a).** To a soln. of 0.50 g (0.90 mmol) of NaOMe in 150 ml of MeOH were added 5.0 g (8.9 mmol) of **3a**. After refluxing for 15 min, the soln. was allowed to cool. AcOH (5 ml) was added, and the solvents were removed under reduced pressure.  $\text{H}_2\text{O}$  (40 ml) was added to the crude product, and the resulting soln. was evaporated. This operation was repeated twice to ensure the complete removal of the methyl benzoate. TLC ( $\text{CH}_2\text{Cl}_2/\text{MeOH}$  85:15) indicated that the material thus obtained was mainly **3b**. After careful drying, it was used without further purification for the next step. To a soln. of **3b** in pyridine (30 ml), 2.8 g (2.8 ml, 8.9 mmol) of  $[(i\text{-Pr})_2\text{SiCl}]_2\text{O}$  were added dropwise, and the soln. was allowed to stand overnight at r.t. After removal of the solvent, the residue was partitioned between  $\text{Et}_2\text{O}$  and  $\text{H}_2\text{O}$ . The org. layer was dried over  $\text{MgSO}_4$ , filtered, evaporated, and the crude material obtained by chromatography over silica gel ( $\text{CH}_2\text{Cl}_2/\text{Et}_2\text{O}$  2:1) to give 2.6 g (62%) of **4a**. M.p.  $128^\circ$ .  $[\alpha]_D^{20} = -3.1^\circ$  ( $c = 5.0$ , MeOH).  $^1\text{H-NMR}$  ( $\text{CDCl}_3$ ): 8.21 ( $s$ ,  $\text{H-C}(2)$ ); 7.9–7.2 ( $m$ , 4 arom. H); 6.00 ( $d$ ,  $J(1',2') = 2$ ,  $\text{H-C}(1')$ ); 4.7–4.4 ( $m$ ,  $\text{H-C}(2')$ ); 4.3–4.1 ( $m$ ,  $\text{H-C}(3')$ ,  $\text{H-C}(4')$ , 2  $\text{H-C}(5')$ ); 3.17 ( $m$ , OH); 1.1–0.9 ( $m$ , 4  $\text{Me}_2\text{CH}$ ). Anal. calc. for  $\text{C}_{24}\text{H}_{40}\text{N}_2\text{O}_5\text{Si}$  (464.64): C 58.50, H 8.13, N 5.69; found: C 58.02, H 8.18, N 5.68.

**1-[2'-Deoxy-3',5'-O-(1,1,3,3-tetraisopropylidisiloxane-1,3-diyl)- $\beta$ -D-ribofuranosyl]-1H-benzimidazole (5a).** A soln. of 2.0 g (4.0 mmol) of **4a**, 1.0 g (80 mmol) of 4-(dimethylamino)pyridine and 0.8 g (0.7 ml, 4.4 mmol) of *p*-tolyl chlorothioformate in  $\text{CH}_3\text{CN}$  (50 ml) was stirred for 2 h at r.t. Then, TLC ( $\text{CH}_2\text{Cl}_2/\text{Et}_2\text{O}$  4:1) indicated that the reaction was complete. The solvent was evaporated and the crude material partitioned between  $\text{Et}_2\text{O}$  and  $\text{H}_2\text{O}$ . The org. layer was washed with cold 1N HCl, dried over  $\text{MgSO}_4$ , and evaporated. To the yellow oil (**4b**), degassed toluene (50 ml) was added followed by 1.2 g (1.0 ml, 4.0 mmol) of  $\text{Bu}_3\text{SnH}$  and 0.15 g of 2,2'-azobis(2-methylpropanenitrile). The soln. was refluxed for 2 h and then allowed to cool. Evaporation of toluene followed by silica-gel chromatography of the resulting oil ( $\text{CH}_2\text{Cl}_2/\text{Et}_2\text{O}$  4:1) afforded 1.6 g (85%) of **5a**. Oil;  $[\alpha]_D^{20} = -0.7^\circ$  ( $c = 3.0$ ,  $\text{CH}_2\text{Cl}_2$ ).  $^1\text{H-NMR}$  ( $\text{CDCl}_3$ ): 8.12 ( $s$ ,  $\text{H-C}(2)$ ); 7.9–7.7 ( $m$ , 1 arom. H); 7.6–7.1 ( $m$ , 3 arom. H); 6.22 ( $t'$ ,  $J(1',2') = 6$ ,  $\text{H-C}(1')$ ); 5.77 ( $t'$ ,  $J(2',3') = J(3',4') = 7$ ,  $\text{H-C}(3')$ ); 4.2–3.8 ( $m$ ,  $\text{H-C}(4')$ , 2  $\text{H-C}(5')$ ); 2.66 ( $dd$ , 2  $\text{H-C}(2')$ ); 1.1–0.9 ( $m$ , 4  $\text{Me}_2\text{CH}$ ). Anal. calc. for  $\text{C}_{24}\text{H}_{40}\text{N}_2\text{O}_4\text{Si}_2$  (476.70): C 60.50, H 8.40, N 5.88; found: C 60.39, H 8.42, N 5.89.

**1-(2'-Deoxy- $\beta$ -D-ribofuranosyl)-1H-benzimidazole (5b).** To a THF soln. of 2.0 g (6.3 mmol) of  $\text{Bu}_4\text{NF}$ , 1.5 g (3.1 mmol) of **5a** were added. After stirring for 15 min, the solvent was evaporated and  $\text{H}_2\text{O}$  was added. The resulting soln. was extracted with  $\text{Et}_2\text{O}$ , and the  $\text{H}_2\text{O}$  phase was collected. Evaporation followed by silica-gel chromatography ( $\text{CH}_2\text{Cl}_2/\text{MeOH}$  9:1) gave 0.56 g (80%) of **5b**. M.p.  $152^\circ$  ([5];  $154^\circ$ ).  $[\alpha]_D^{20} = -33.1^\circ$  ( $c = 3.0$ , MeOH; [5]:  $[\alpha]_D = -30.5^\circ$ ).  $^1\text{H-NMR}$  ( $(\text{D}_6)\text{DMSO}$ ): 8.45 ( $s$ ,  $\text{H-C}(2)$ ); 7.7–7.6 ( $m$ , 2 arom. H); 7.3–7.1 ( $m$ , 2 arom. H); 6.40, 6.31 ( $dd$ ,  $J(1',2') = 6.1$ ,  $J(1',2') = 6.3$ ,  $\text{H-C}(1')$ ); 5.32 ( $d$ ,  $J = 4.2$ , OH); 4.94 ( $t$ ,  $J = 5.2$ , OH); 4.4–4.3 ( $m$ ,  $\text{H-C}(3')$ ); 3.9–3.8 ( $m$ ,  $\text{H-C}(4')$ ); 3.6–3.5 ( $m$ , 2  $\text{H-C}(5')$ ); 2.6–2.3 ( $m$ , 2  $\text{H-C}(2')$ ).

**1-(2'-Deoxy-5'-O-dimethoxytrityl- $\beta$ -D-ribofuranosyl)-1H-benzimidazole 3'-[(*p*-Chlorophenyl) (2-Cyanoethyl) Phosphate] (6).** A soln. of 0.5 g (1.9 mmol) of **5b** in pyridine (5 ml) was evaporated to dryness to ensure the removal of traces of  $\text{H}_2\text{O}$ . This operation was repeated twice, and then pyridine (3 ml) as well as 0.8 g (2.3 mmol) of dimethoxytrityl chloride were added. Stirring was immediately started, and after 2 h, MeOH (5 ml) was added. After 30 min, the solvents were evaporated, and the residue was partitioned between  $\text{H}_2\text{O}$  and  $\text{Et}_2\text{O}$ . The org. layer was dried over  $\text{MgSO}_4$  and concentrated to a sirup. The presumed 5'-O-dimethoxytrityl-deoxynucleoside was co-evaporated twice with pyridine (5 ml). The rotatory evaporator was flushed with Ar. Then, dioxane (10 ml), 1.3

g (1.2 ml, 5.0 mmol) of (*p*-chlorophenyl) (2-cyanoethyl) phosphochloridate as well as 1.5 ml (6.0 mmol) of *N*-methylimidazole were added. The soln. was stirred for 1 h and then hydrolyzed with H<sub>2</sub>O/pyridine 1:1 (5 ml) for 15 min. After evaporation of dioxane, a 4% aq. NaHCO<sub>3</sub> soln. was added, and the product was extracted with CH<sub>2</sub>Cl<sub>2</sub>. The org. layer was dried (MgSO<sub>4</sub>), concentrated, and chromatographed over silica gel (CH<sub>2</sub>Cl<sub>2</sub>/MeOH 9:1) to afford 0.8 g (61%) of **6** as a foam.  $[\alpha]_D^{20} = -21.3^\circ$  (*c* = 1.0, CH<sub>2</sub>Cl<sub>2</sub>). <sup>1</sup>H-NMR (CDCl<sub>3</sub>): 8.45 (*s*, H-C(2)); 7.7–7.1 (*m*, 2 arom. H); 6.37 (*t'*,  $J(1',2') = J(1',2'') = 6.2$ , H-C(1')); 4.4–4.2 (*m*, H-C(3'), OCH<sub>2</sub>CH<sub>2</sub>CN); 3.9–3.8 (*m*, H-C(4')); 3.80 (*s*, 2 CH<sub>3</sub>O); 3.6–3.5 (*m*, 2 H-C(5')); 2.7–2.3 (*m*, 2 H-C(2'), OCH<sub>2</sub>CH<sub>2</sub>CN). Anal. calc. for C<sub>42</sub>H<sub>39</sub>ClN<sub>3</sub>O<sub>7</sub>P (764.20): C 66.06, H 5.11, N 5.50; found: C 66.16, H 5.08, N 5.51.

## REFERENCES

- [1] A. M. Hogg, J. G. Kelland, J. C. Vederas, Ch. Tamm, *Helv. Chim. Acta* **1986**, *69*, 908.
- [2] Ch. Papageorgiou, Ch. Tamm, *Tetrahedron Lett.* **1986**, *27*, 555.
- [3] I. Tamm, K. Folkers, C. H. Shunk, *J. Bacteriol.* **1956**, *72*, 54.
- [4] I. Tamm, M. N. Nemes, S. Osterhoud, *J. Exp. Med.* **1960**, 339.
- [5] C. P. Whittle, R. K. Robins, *J. Am. Chem. Soc.* **1965**, *87*, 4940.
- [6] J.-L. Barascut, B. L. Kan, J.-L. Imbach, *J. Heterocycl. Chem.* **1977**, *14*, 1305.
- [7] I. W. Southon, W. Pfeleiderer, *Chem. Ber.* **1978**, *111*, 996.
- [8] Y. Mizumo, M. Ikehara, F. Ishikawa, H. Ikehata, *Chem. Pharm. Bull.* **1962**, *10*, 761.
- [9] M. J. Robins, J. S. Wilson, F. Hansske, *J. Am. Chem. Soc.* **1983**, *105*, 4059.
- [10] S. De Bernardini, F. Waldmeier, Ch. Tamm, *Helv. Chim. Acta* **1981**, *64*, 2142.