

Advanced 

Synthesis & Catalysis

Accepted Article

Title: Synthesis of Benzofulvenes through Palladium-Catalyzed Sequential Three-Component Reactions

Authors: Bo Zhou, Zhuo Wu, Weixin Qi, Xueliang Sun, and Yanghui Zhang

This manuscript has been accepted after peer review and appears as an Accepted Article online prior to editing, proofing, and formal publication of the final Version of Record (VoR). This work is currently citable by using the Digital Object Identifier (DOI) given below. The VoR will be published online in Early View as soon as possible and may be different to this Accepted Article as a result of editing. Readers should obtain the VoR from the journal website shown below when it is published to ensure accuracy of information. The authors are responsible for the content of this Accepted Article.

To be cited as: *Adv. Synth. Catal.* 10.1002/adsc.201801141

Link to VoR: <http://dx.doi.org/10.1002/adsc.201801141>

DOI: 10.1002/adsc.201((will be filled in by the editorial staff))

The Synthesis of Benzofulvenes through Palladium-Catalyzed Sequential Three-Component Reactions

Bo Zhou,^a Zhuo Wu,^a Weixin Qi,^a Xueliang Sun,^a Yanghui Zhang^{a, *}

^a School of Chemical Science and Engineering, Shanghai Key Laboratory of Chemical Assessment and Sustainability, Tongji University, 1239 Siping Road, Shanghai 200092, P. R. China
E-mail: zhangyanghui@tongji.edu.cn; Web: http://zhangyhgroup.tongji.edu.cn

Received: ((will be filled in by the editorial staff))

Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/adsc.201#####>. ((Please delete if not appropriate))


Abstract: An approach for the synthesis of benzofulvenes has been developed through palladium-catalyzed sequential three-component reactions. The reactions likely involve *C,C*-palladacycles as the key intermediates. The palladacycles are generated through cascade reactions of aryl halides and alkynes, and then reacted with CH_2Br_2 to form benzofulvenes the final products.

Keywords: *C,C*-palladacycles; C–H activation; Alkylation; Palladium; Benzofulvenes.

Benzofulvenes are key privileged scaffolds present in natural products and bioactive molecules and have found versatile applications in materials science, medicinal chemistry and organometallics.^[1] In particular, benzofulvenes are important monomers in polymer chemistry, and polybenzofulvenes often show intriguing properties.^[1e, 1f] A great number of methods have been developed for the synthesis of benzofulvenes.^[2] The traditional methods often rely on the use of structurally complex starting materials. Notably, the synthesis of benzofulvenes through C–H activation has also been reported,^[3] However, the reactions are limited to those using aryl ketones as starting materials. Therefore, continuous efforts should still be devoted to seeking new reactions for the synthesis of benzofulvenes.


As an important class of organometallic compounds, metallacycles are very common intermediates in transition-metal catalysis,^[4] In particular in transition-metal-catalyzed C–H functionalization reactions.^[5] In this context, *C,C*-palladacycles are particularly attractive.^[6] *C,C*-palladacycles consist of a C–Pd–C bonding motif and contain two C–Pd bonds. Due to their unique structures, *C,C*-palladacyclic intermediates may exhibit novel reactivity. More importantly, the presence of two C–Pd bonds offers opportunities to develop new transformations. *C,C*-Palladacycles are usually obtained by Pd-catalyzed intramolecular C–H activation of aryl halides as starting materials.^[7]

However, for this method, since *C,C*-palladacycles are synthesized directly from the Pd-mediated cyclization of the substrates, the substrates have complex structures and have to be presynthesized. Therefore, it is highly desirable to develop new methods for the synthesis of *C,C*-palladacycles by employing simple substrates.



Scheme 1. Synthesis of Benzofulvenes *via* Transition-Metal-Catalyzed C–H Activation.

It has been reported that *C,C*-palladacycles could be obtained from aryl halides and alkynes, which represents a straightforward method for the construction of *C,C*-palladacycles from relatively simple precursors.^[8] However, the resulting palladacycles usually underwent intramolecular cyclization. Although the intermolecular arylation of *C,C*-palladacycles formed via reactions of aryl iodides and alkynes has been developed, the aryl iodides were also the starting materials forming the *C,C*-palladacycles.^[9] Actually, this reaction indicates that it should be quite a challenge to functionalize such *C,C*-palladacycles obtained from aryl halides and alkynes with other external reagents. Since multiple reactants, intermediates, and steps are involved in the reactions, each step should proceed in a well-defined sequence to make the reactions occur as desired. Herein, we report the alkylation reaction of *C,C*-palladacyclic intermediates obtained from aryl halides and alkynes. The transformation represents a novel method for the synthesis of benzofulvenes.



^[a]Yields of the the isolated products.

^[b]Observed by GC-MS.

Scheme 2. Alkylation of *C,C*-palladacyclic intermediate with alkyl iodides.

As it has been demonstrated that *C,C*-palladacycles have high reactivity towards alkyl halides,^[7n] we first investigated alkylation reaction of *C,C*-palladacyclic intermediates obtained from aryl halides and alkynes. Therefore, iodobenzene **1a** and diphenylacetylene **2a** were selected for constructing *C,C*-palladacycle, and *n*-butyl iodide was used as the alkylating reagent. After extensive screening, alkylated product **4aaa** was obtained in 52% yield under the conditions as shown in **Scheme 2**. An isomer of **4aaa** and two dialkylated products were also observed by GC-MS. Although the structures of these minor isomers could not be identified due to insufficient quantities, the isomer of **4aaa** may result from the alkylation of the alkene moiety (for the mechanism of the alkylation reaction see the Supporting Information).

Table 1. Optimization of Reaction Conditions for the Benzofulvene-Forming Reaction.




Entry	Base (mmol)	Solvent (mL)	Yield (%) ^[a]
1	KOAc (2.4)	DMA (2)	11
2	KOAc / PhCO ₂ K (2.4 / 0.8)	DMA (2)	21
3	KOAc (3.2)	DMA (2)	15
4	KOAc / PhCO ₂ K (2.4 / 0.8)	DMA / <i>i</i> -PrOH (2 / 0.5)	30
5	KOAc / PhCO ₂ K (2.4 / 0.8)	DMA / PEG (2 / 0.5)	50
6	KOAc / PhCO ₂ K (2.4 / 0.8)	DMA / PEG / <i>i</i> -PrOH (2 / 0.5 / 0.5)	61
7	KOAc / PhCO ₂ K (2.4 / 0.8)	DMA / PEG / <i>i</i> -PrOH (1 / 2 / 0.5)	72

8	KOAc / PhCO ₂ K (7 / 0.8)	PEG / DMA / <i>i</i> -PrOH (2 / 1 / 0.5)	90 (85 ^[b])
---	--------------------------------------	--	-------------------------

^[a]The yields were determined by ¹H NMR analysis of the crude reaction mixture using CHCl₂CHCl₂ as the internal standard. ^[b]Isolated yield. PEG = poly(ethylene glycol) 400. DMA = *N,N*-dimethylacetamide.


The greatest advantage of *C,C*-palladacycles is that they contain two C-Pd bonds. Although the above alkylation reaction demonstrates the feasibility of intermolecular functionalization of *C,C*-palladacycles obtained from aryl halides and alkynes, only one C-Pd bond was functionalized. Our aim is to develop synthetically useful reactions by functionalizing both C-Pd bonds of *C,C*-palladacycles. We found that *C,C*-palladacycles can react with CH₂Br₂, affording fluorenes as the products.^[10] Therefore, we turned to investigate the reaction of the *C,C*-palladacycles formed in situ from aryl halides and alkynes with CH₂Br₂. Unexpectedly, benzofulvene **6aa** was obtained in the reaction of **1a**, **2a**, and **5** in the presence of Pd(OAc)₂ and KOAc (**Table 1**, entry 1). Spurred by this exciting result, we continued to optimize the reaction conditions. The addition of PhCO₂K promoted the yield to 21% (entry 2). The same amount of KOAc was less effective (entry 3). The yield was improved to 30% when 0.5 mL isopropanol was added. The isopropanol should act as a reductant, because the catalytic cycle is expected to start with Pd(0). Gratefully, a yield of 50% was obtained by replacing isopropanol with PEG (entry 5).^[11] The yield increased when the reaction was carried out in a mixture of DMA, PEG, and isopropanol (entries 6 and 7). Finally, benzofulvene was formed in 90% yield when the amount of KOAc was enhanced to 7 mmol. While the role of KOAc is unknown, KOAc is a common base in the C-H activation reaction of aryl halides and can promote C-H activation.^[12]

To gain insight into the mechanism of the benzofulvene-forming reaction, we first synthesized 2,3-diphenyl-1*H*-indene. When 2,3-diphenyl-1*H*-indene was allowed to react with CH₂Br₂, **6aa** was obtained in a yield of 70%. Furthermore, a small amount of 2,3-diphenyl-1*H*-indene was observed under some conditions that were tested during the course of reaction optimization. These outcome indicate that 2,3-diphenyl-1*H*-indene should be formed as the intermediate in the reaction. The benzofulvene was formed in good yields when the reaction was performed in the presence of TEMPO, which is against a SET process.




Scheme 3. Mechanistic studies.

On the basis of these experimental results and previous reports,^[9a, 10, 13] a mechanism is proposed as shown in **Scheme 4**. The catalytic cycle starts with the oxidative addition of iodobenzenes to Pd(0). The resulting arylPd(II) species **A** undergo migratory insertion to form vinylPd(II) species **B**. The subsequent intramolecular C–H activation generates *C,C*-palladacycle **C**. **C** may undergo oxidative addition of CH₂Br₂ to form Pd(IV) complex **D**. The reductive elimination and intramolecular oxidative addition yield six-membered palladacycle **G**. Alternatively, **G** could also be formed *via* a carbene complex **F**. The reductive elimination of **G** generates intermediate 2, 3-diphenyl-1*H*-indene **H** and releases Pd(II), which is reduced to Pd(0) to close the catalytic cycle. Under the basic conditions, **H** can react with CH₂Br₂ to form alkylated products **J**. The subsequent elimination reaction yields final benzofulvene products **6**.




Scheme 4. Proposed mechanism.

Having developed an efficient synthetic protocol for benzofulvenes, we then investigated the substrate scope of the method. We first examined the compatibility of various substituted iodobenzenes. As shown in **Table 2**, a variety of functional groups at the *para* positions, including methyl, *tert*-butyl, methoxy, phenyl, and acetamido, were compatible (**6ba–6fa**), and substrates bearing a *meta*-substituent were also suitable (**6ga** and **6ha**). The reactivity of disubstituted iodobenzenes was also investigated. A range of iodobenzene derivatives were transformed into the corresponding benzofulvenes in moderate yields (**6ia–6la**). It should be mentioned that a trace amount of isomers were observed by GC-MS in most of the reactions. However, the structures of the isomers could not be identified due to insufficient amount, and they may result from the *cis-trans* isomerization of the vinyl palladium complexes (for the possible structures of the isomers and the formation processes see the supporting information).^[9a]

Table 2. Iodobenzene scope of the benzofulvene-forming reaction.^[a]

^[a]Yields of the isolated products.

Next, we examined the performance of substituted diphenylacetylenes. A variety of symmetrical diphenylacetylenes bearing two methyl, *tert*-butyl, methoxy, or fluoro groups at the *para* positions reacted with iodobenzene and CH₂Br₂ to afford various benzofulvene derivatives in moderate or good yields (**Table 3**, **6ab–6ae**). The *meta*-substituted diphenylacetylenes also underwent the cascade reaction (**6af–6ah**), and even the substrate bearing two *ortho*-fluoro groups was compatible (**6ai**). The unsymmetrical alkynes was also examined. 2-Propyl phenylacetylenes was subjected to the standard conditions. The indene **6aj** was obtained instead of the expected benzofulvene. The reason might be that

Table 3. Diarylacetylene scope of the benzofulvene-forming reaction.^[a]

^[a]Yields of the isolated products.

the benzylic C-H bond is less acidic and cannot be deprotonated under the conditions. It should be noted that the reaction exhibit high regioselectivity and **6aj** was the sole product.^[14]

In conclusion, we have developed a new approach for the synthesis of benzofulvenes through palladium-catalyzed sequential three-component reactions. The reactions involve putative *C,C*-palladacycles as the key intermediates. Unlike the common methods for accessing such species from structurally complex starting materials, in this case the palladacycles were accessed from simple aryl halides and alkynes. The resulting *C,C*-palladacycles then react with CH₂Br₂ to form benzofulvenes as the final products. Further studies towards understanding the detailed mechanism and exploring other reactions of *C,C*-palladacycles generated in this way are underway in our lab.

Experimental Section

A 35 mL Schlenk-type tube (with a Teflon high pressure valve and side arm) equipped with a magnetic stir bar was charged with diphenylacetylenes **2** (0.24 mmol), Pd(OAc)₂ (4.5 mg, 0.02 mmol), PhCO₂K (128.2 mg, 0.8 mmol) and KOAc (687.0 mg, 7.0 mmol). Then the mixture was first stirred at room temperature for 1 minute followed by the addition of DMA (1 mL), *i*-PrOH (0.5 mL), aryl iodides **1** (0.2 mmol), CH₂Br₂ **5** (98.2 μL, 1.4 mmol) and PEG (Mn 400, 2 mL). The reaction was frozen with liquid nitrogen, and the tube was then evacuated and backfilled with nitrogen (10 times). The reaction was stirred at 75 °C for 12 h. Upon completion, the reaction mixture was cooled to room temperature, diluted with EtOAc (15 mL) and washed with brine (2X15 mL). The organic phase was dried over Na₂SO₄, filtered and concentrated *in vacuo*. The residue was purified by preparative silica gel TLC to afford the corresponding products.

Acknowledgements

The work was supported by the National Natural Science Foundation of China (No.216721626 and No.2137217) and Shanghai Science and Technology Commission (14DZ2261100). We thank Prof. Keary M. Engle (The Scripps Research Institute) for helpful discussions.

References

- [1] a) B. A. Trofimov, N. A. Nedolya in *Comprehensive Heterocyclic Chemistry*, (Eds.: G. Jones, C. A. Ramsden), Elsevier: Oxford, U.K., **2008**; b) T. Nakano, K. Takewaki, T. Yade, Y. Okamoto, *J. Am. Chem. Soc.* **2001**, *123*, 9182; c) A. S. Felts, B. S. Siegel, S. M. Young, C. W. Moth, T. P. Lybrand, A. J. Dannenberg, L. J. Marnett, K. Subbaramaiah, *J. Med. Chem.* **2008**, *51*, 4911; d) C. Görl, H. G. Alt, *J. Organomet. Chem.* **2007**, *692*, 5727; e) Y. Kosaka, S. Kawauchi, R. Goseki, T. Ishizone, *Macromolecules* **2015**, *48*, 4421; f) Y. Kosaka, K. Kitazawa, S. Inomata, T. Ishizone, *ACS Macro Lett.* **2013**, *2*, 164.
- [2] Selected examples for the synthesis of benzofulvenes and the references cited therein: a) P. Cordier, C. Aubert, M. Malacria, E. Lacôte, V. Gandon, *Angew. Chem. Int. Ed.* **2009**, *48*, 8757; b) J.-M. Lu, Z.-B. Zhu, M. Shi, *Chem. Eur. J.* **2009**, *15*, 963; c) X. Fang, P. Yu, G. Prina Cerai, B. Morandi, *Chem. Eur. J.* **2016**, *22*, 15629; d) S. Ye, J. Wu, *Org. Lett.* **2011**, *13*, 5980; e) R. K. Mohamed, S. Mondal, K. Jorner, T. F. Delgado, V. V. Lobodin, H. Ottosson, I. V. Alabugin, *J. Am. Chem. Soc.* **2015**, *137*, 15441; f) S. Shin, J.-Y. Son, C. Choi, S. Kim, P. H. Lee, *J. Org. Chem.* **2016**, *81*, 11706.
- [3] a) F. W. Patureau, T. Besset, N. Kuhl, F. Glorius, *J. Am. Chem. Soc.* **2011**, *133*, 2154; b) K. Tsuchikama, M. Kasagawa, K. Endo, T. Shibata, *Synlett*, **2010**, *2010*, 97; c) R. K. Chinnagolla, M. Jeganmohan, *Eur. J. Org. Chem.* **2012**, *2012*, 417.
- [4] a) J. Dupont, C. S. Consorti, J. Spencer, *Chem. Rev.* **2005**, *105*, 2527; b) T. R. Cook, P. J. Stang, *Chem. Rev.* **2015**, *115*, 7001; c) I. P. Beletskaya, A. V. Cheprakov, *J. Organomet. Chem.* **2004**, *689*, 4055.
- [5] a) Y. Zhang, G. Shi, J.-Q. Yu, in *Comprehensive Organic Synthesis*, 2nd ed., Vol. 3 (Eds.: P. Knochel, G. A. Molander), Elsevier, Oxford, **2014**, pp. 1101-1209; b) *C-H Activation*, (Eds.: J.-Q. Yu, Z. Shi), Springer, Berlin, **2010**; c) T. W. Lyons, M. S. Sanford, *Chem. Rev.* **2010**, *110*, 1147; d) L. Ackermann, R. Vicente, R. Kapdi Anant, *Angew. Chem. Int. Ed.* **2009**, *48*, 9792; e) J. Miao, H. Ge, *Eur. J. Org. Chem.* **2015**, *2015*, 7859.
- [6] a) C. Juan, P. Pilar, C. Ernesto, *Coordin. Chem. Rev.* **1999**, *193*, 207; b) A. Steffen, R. M. Ward, W. D. Jones, T. B. Marder, *Coordin. Chem. Rev.* **2010**, *254*, 1950; c) H. Yoon, A. Lossouarn, F. Landau, M. Lautens, *Org. Lett.* **2016**, *18*, 6324; d) M. Pérez-Gómez, L. Navarro, I. Saura-Llamas, D. Bautista, M. Lautens, J.-A. García-López, *Organometallics*, **2017**, *36*, 4465; e) A. Lu, X. Ji, B. Zhou, Z. Wu, Y. Zhang, *Angew. Chem. Int. Ed.* **2018**, *57*, 3233.
- [7] a) T. Piou, A. Bunescu, Q. Wang, L. Neuville, J. Zhu, *Angew. Chem. Int. Ed.* **2013**, *52*, 12385; b) J.-X. Yan, H. Li, X.-W. Liu, J.-L. Shi, X. Wang, Z.-J. Shi, *Angew. Chem. Int. Ed.* **2014**, *53*, 4945; c) R. Deng, Y. Huang, X. Ma, G. Li, R. Zhu, B. Wang, Y.-B. Kang, Z. Gu, *J. Am.*

- Chem. Soc.* **2014**, *136*, 4472; d) D.-W. Gao, Q. Yin, Q. Gu, S.-L. You, *J. Am. Chem. Soc.* **2014**, *136*, 4841; e) M. Wang, X. Zhang, Y.-X. Zhuang, Y.-H. Xu, T.-P. Loh, *J. Am. Chem. Soc.* **2015**, *137*, 1341; f) L. Yang, R. Melot, M. Neuburger, O. Baudoin, *Chem. Sci.* **2017**, *8*, 1344; g) G. Dyker, *Angew. Chem. Int. Ed.* **1992**, *31*, 1023; h) G. Dyker, *Angew. Chem. Int. Ed.* **1994**, *33*, 103; i) D. Masselot, J. P. H. Charmant, T. Gallagher, *J. Am. Chem. Soc.* **2006**, *128*, 694; j) J. Ye, Z. Shi, T. Sperger, Y. Yasukawa, C. Kingston, F. Schoenebeck, M. Lautens, *Nat. Chem.* **2016**, *9*, 361; k) M. Pérez-Gómez, J.-A. García-López, *Angew. Chem. Int. Ed.* **2016**, *55*, 14389; l) T.-J. Hu, G. Zhang, Y.-H. Chen, C.-G. Feng, G.-Q. Lin, *J. Am. Chem. Soc.* **2016**, *138*, 2897; m) A. Gutiérrez-Bonet, F. Juliá-Hernández, B. de Luis, R. Martín, *J. Am. Chem. Soc.* **2016**, *138*, 6384; n) D. Chen, G. Shi, H. Jiang, Y. Zhang, Y. Zhang, *Org. Lett.* **2016**, *18*, 2130.
- [8] a) G. Wu, A. L. Rheingold, S. J. Geib, R. F. Heck, *Organometallics*, **1987**, *6*, 1941; b) T. Sakakibara, Y. Tanaka, T.-I. Yamasaki, *Chem. Lett.* **1986**, 797; c) R. C. Larock, M. J. Doty, Q. Tian, J. M. Zenner, *J. Org. Chem.* **1997**, *62*, 7536; d) Q. Tian, R. C. Larock, *Org. Lett.* **2000**, *2*, 3329; e) M. A. Campo, R. C. Larock, *J. Am. Chem. Soc.* **2002**, *124*, 14326; f) J. Zhao, M. Campo, C. Larock Richard, *Angew. Chem. Int. Ed.* **2005**, *44*, 1873; g) N. Chernyak, D. Tilly, Z. Li, V. Gevorgyan, *Chem. Commun.* **2010**, *46*, 150; h) C. Wang, S. Rakshit, F. Glorius, *J. Am. Chem. Soc.* **2010**, *132*, 14006.
- [9] a) G. Dyker, A. Kellner, *Tetrahedron Lett.*, **1994**, *35*, 7633; b) A three-component cross-coupling of aryl halides, alkynes, and arynes might also involve C,C-palladacycles as the intermediate. However, another mechanism was proposed. Z. Liu, R. C. Larock, *Angew. Chem. Int. Ed.* **2007**, *46*, 2535.
- [10] a) G. Shi, D. Chen, H. Jiang, Y. Zhang, Y. Zhang, *Org. Lett.* **2016**, *18*, 2958; b) C. Shao, Z. Wu, X. Ji, B. Zhou, Y. Zhang, *Chem. Commun.* **2017**, *53*, 10429; c) Z. Wu, D. Ma, B. Zhou, X. Ji, X. Ma, X. Wang, Y. Zhang, *Angew. Chem. Int. Ed.* **2017**, *56*, 12288.
- [11] For the use of PEG in organometallic reactions, see: L. Ackermann, R. Vicente, *Org. Lett.* **2009**, *11*, 4922.
- [12] For the use of KOAc in C-H activation, see: a) C. Amatore, A. Jutand, *Acc. Chem. Res.* **2000**, *33*, 314; b) M. Lafrance, C. N. Rowley, T. K. Woo, K. Fagnou, *J. Am. Chem. Soc.* **2006**, *128*, 8754.
- [13] J. Campora, P. Palma, D. del Rio, J. A. Lopez, P. Valerga, *Chem. Commun.* **2004**, 1490.
- [14] a) W. Tao, L. J. Silverberg, A. L. Rheingold, R. F. Heck, *Organometallics*, **1989**, *8*, 2550; b) R. C. Larock, M. J. Doty, S. Cacchi, *J. Org. Chem.* **1993**, *58*, 4579; c) D. V. Kadnikov, R. C. Larock, *Org. Lett.* **2000**, *2*, 3643.

COMMUNICATION

The Synthesis of Benzofulvenes through Palladium-Catalyzed Sequential Three-Component Reactions

Adv. Synth. Catal. **Year**, *Volume*, Page – Page

Bo Zhou, Zhuo Wu, Weixin Qi, Xueliang Sun, Yanghui Zhang *

