Primary Amines as Nucleation Centres for Gold Clustering, and the Structural Chemistry of Polygold Ammonium Cations[†]

Klaus Angermaier and Hubert Schmidbaur*

Anorganisch-chemisches Institut der Technischen Universität München, Lichtenbergstrasse 4, D-85747 Garching, Germany

Treatment of *tert*-butylamine and benzylamine with $[Au(PR'_3)]BF_4$ led to the monoauration products $[(R'_3P)AuNH_2R]^+BF_4^-$ (R' = Me, R = Bu' 1a; R' = Me, R = CH_2Ph 1b; R'_3 = Ph_2Me, R = Bu' 1c). The salts $[\{Au(PMe_3)\}_3NR]^+BF_4^-$ (R' = Me, R = Bu' 2a or CH_2Ph 2b) derived from the same amines were obtained using the corresponding oxonium salt $[\{Au(PMe_3)\}_3O]^+BF_4^-$. A diauration product $[\{Au(PMe_3)\}_3NR]^+BF_4^-$ 3 was generated upon slow hydrolytic degradation of 2b in solution. Treatment of NH(SiMe_3)_2 with $[\{Au(PMe_3)\}_3O]^+BF_4^-$ afforded both the tris- and tetra-aurated ammonium salts, $[\{Au(PMe_3)\}_3NSiMe_3]^+BF_4^-$ 2c and $[\{Au(PMe_3)\}_4N]^+BF_4^-$ 4, depending on the reaction conditions. The cation of 2c is one of the very rare examples of silylammonium species. Compound 4 was also obtained from the oxonium salt and ammonia. All compounds have been characterized through analytical and spectroscopic data. The crystal structures of 1a, 1c and 3 have been determined by X-ray diffraction methods: 1c is a salt with monomeric gold ammonium cations, but 1a and 3 show supramolecular aggregation of the cations into dimers with short intercationic Au····Au contacts.

The affinity of gold for nitrogen is very low indeed, and most compounds with gold-nitrogen bonds are of limited stability.¹ It is therefore most surprising that this affinity is increased if more than one gold atom is present at a given nitrogen centre: polyauration appears to be much more facile than the introductory step of the metallation at nitrogen.² Work in the last two decades has shown that compounds with one, two, three, four, and even five gold atoms at nitrogen can be prepared.³⁻¹⁵ The stability of these clustering products is found to increase remarkably with the size of the nitrogen-centred metal These results have been accumulated from aggregates. occasional observations during various preparative studies in gold chemistry. It was therefore felt that this field of goldnitrogen chemistry should now be given a more systematic consideration by developing straightforward synthetic pathways and elucidating more of the molecular structures of the individual species.

Recent investigations have shown that many of the peculiar stoichiometries and properties of gold compounds have their origin in an unusual structural chemistry.¹⁶⁻¹⁸ Most twoco-ordinate compounds of the type L-Au-X have been found to form supramolecular aggregates with Au · · · Au distances of *ca*. 3.0 Å associated with bond energies of *ca*. 8 kcal mol⁻¹ (*ca*. 33.47 kJ mol⁻¹).¹⁹⁻²² This secondary bonding is representing weak forces between the closed-shell metal atoms (Au^I 5d¹⁰) originating from significant changes in the energy characteristics of the valence electrons of gold by relativistic effects.²³⁻²⁵ In the Periodic Table these effects reach a local maximum for gold in the heavy-element region.^{26.27}

The strength of the Au \cdots Au interactions is influenced also by the electronic effects of the substituents L and X at gold, and some of the experimental findings (above),²⁸⁻³³ as well as theoretical calculations,³⁴ have indicated that soft ligands of low group electronegativity should shorten the Au \cdots Au distance and increase the bond energy. More recent data have shown, however, that even within the series with X = halide the results are not consistent with this simple picture.³⁵ On the other hand it is becoming more and more obvious that steric effects are playing a decisive role, since the weak forces associated with the Au···Au contacts are easily overruled by steric repulsion and other components such as packing forces. (In solution, it is the solvation by solvent molecules which is overruling the aggregation through Au···Au contacts.) In cluster aggregates these Au···Au contacts are synergistic and lead to the stabilization mentioned as the key observation.³⁶⁻⁴⁰

The present study was therefore undertaken also in order to contribute to the open questions regarding the intra- and inter-molecular (supramolecular) aggregation of nitrogencentred gold cluster cations [$\{Au(PR'_3)\}_nNR_{4-n}$]⁺ for different numbers of *n* and different substituents R and R'.

Results and Discussion

Preparative Results.—tert-Butylamine and benzylamine were chosen as substrates because it should be possible to follow the stepwise auration at nitrogen most easily. The steric effect of the Bu¹ group was expected to hinder quick and complete auration, whereas benzyl was expected to give products with good crystallization properties for all steps. For small and simple alkyl and aryl groups it has already been demonstrated in previous experiments that complete auration to give tri(gold)-ammonium cations is virtually instantaneous.²

Treatment of NH_2Bu^t with a solution of equimolar quantities of (trimethylphosphine)gold(1) tetrafluoroborate {prepared *in situ* from [Au(PMe₃)Cl] and AgBF₄ in tetrahydrofuran} gives high yields of the monoaurated ammonium salt **1a** (Scheme 1).

$$[Au(PR'_3)]^+BF_4^- \xrightarrow{NH_2R} [(R'_3P)AuNH_2R]^+BF_4^-$$

$$1a R' = Me, R = Bu'$$

$$1b R' = Me, R = CH_2Ph$$

$$1c R'_3 = Ph_2Me, R = Bu'$$

Scheme 1

[†] Supplementary data available: details of the crystal structure determinations are available from Fachinformationszentrum Karlsruhe GmbH, D-76344 Eggenstein-Leopoldshafen, Germany, by quoting the names of the authors, the journal citation, and the CSD number 58808.

Even if an excess of the aurating agent is used only this primary product is obtained. This result has been confirmed in the reaction of the same amine with (methyldiphenylphosphine)gold(1) tetrafluoroborate, which consistently gave only the analogous product 1c. In a similar way, $NH_2(CH_2Ph)$ was converted into the mononuclear complex salt 1b. It therefore appears that $[Au(PR'_3)]^+BF_4^-$ reagents quite generally lead to monoauration products only. No more highly aurated by-products have been observed in the reaction mixtures by NMR spectroscopy.

By contrast, with μ_3 -oxo-tris[(trimethylphosphine)gold(1)] tetrafluoroborate the two amines, NH₂Bu^t and NH₂(CH₂Ph), are triply aurated immediately even at low temperature and with stoichiometric or sub-stoichiometric quantities of the reagent. The products **2a**, **2b** are generated in high yields and with very satisfactory purity. This behaviour is in agreement with the findings of NH₂Me, NH₂Et, NH₂Ph and other simple mono- or di-functional primary amines (like ethylenediamine or diaminobenzenes).

$$[{Au(PMe_3)}_{3}O]^{+}BF_{4}^{-} \xrightarrow{NH_2R} [{Au(PMe_3)}_{3}NR]^{+}BF_{4}^{-}$$
2a R = Bu¹
2b R = CH₂Ph
Scheme 2

Although the pure crystalline compounds are stable at ambient temperature in an inert atmosphere (below), slow degradation is observed in solution, most notably if humidity is allowed to enter into the storage vessel. The benzyl compound **2b** has been found to undergo hydrolysis thereby (in moist dichloromethane) which leads to the crystalline primary degradation product with only two gold atoms left at the nitrogen atom $[{Au(PMe_3)}_2NH(CH_2Ph)]^+BF_4^- 3$. Compound **2a** also shows this solvolysis, as monitored by NMR spectroscopy, but no pure crystalline material could be isolated.

Hexamethyldisilazane (Me₃Si)₂NH is known to react with $[{Au(PR'_3)}_3O]^+$ salts to give salts of the type $[{Au-(PR'_3)}_4N]^+BF_4^-$. As shown, e.g. for the case with R = Ph, no partially aurated cations are detected in this reaction: both trimethylsilyl groups are replaced by (phosphine)gold units, and the imino hydrogen atom is also lost in the quaternization. In the reaction with $[{Au(PMe_3)}_3O]^+BF_4^-$ we were now able to isolate a triply aurated cation still bearing one trimethylsilyl group at the nitrogen centre $[{Au(PMe_3)}_3NSiMe_3]^+BF_4^-2c$. The reaction was found to proceed further if humidity is allowed to enter the reaction mixture, or if moist tetrahydrofuran is used as a solvent. Under these conditions, the quadruply aurated salt $[{Au(PMe_3)}_4N]^+BF_4^- 4$ is the sole product. This compound is also available from the oxonium salt and ammonia, as suggested by previous work. Compound 2c can be converted into 4 by reaction with another mole of the oxonium reagent, preferably in moist solvent.

No attempts have yet been made to aurate further the cation of 4 aiming at the generation of the five-co-ordinate salt of the type $[{Au(PMe_3)}_5N]^{2+}2BF_4^{-}$. It is expected that future work will confirm the existence of this Me₃P analogue of the known Ph₃P complex.¹⁵

General Properties and Spectroscopic Data.—All new complexes have been obtained as colourless (or pale yellow) crystalline materials, except for **1b** which has been obtained as an oil, which are soluble in polar solvents, preferably di- and tri-chloromethane or tetrahydrofuran. These solutions are not very stable, and decomposition is observed after a few hours or days at room temperature. However, samples of the dry, pure solid materials are stable in dry air at ambient temperature. Their composition has been confirmed by elemental analysis. The cations of the compounds have been registered as parent ions in the mass spectra of these solutions (field desorption, fast atom bombardment, or chemical ionization techniques).

The ¹H, ¹³C, ²⁹Si and ³¹P NMR spectra of the halogenocarbon solutions show the expected patterns. The number of resonances and their relative intensities are consistent with the proposed composition. In a few selected cases the tetrafluoroborate anions have been registered separately by ¹⁹F and ¹¹B NMR spectroscopy, as well as by their IR absorptions and negative-ion mass spectra. Details are given in the Experimental section. While most chemical shift values and (where applicable) coupling constants do not need special comments, it is interesting to mention a more unusual observation in the ³¹P NMR spectrum of compound 2c: the ${}^{31}P-{}^{1}H$ resonance of this cation is a triplet signal with a clear $^{31}P_{-}^{14}N$ coupling (J = 22.8 Hz). We suggest that the three-fold symmetry of the cation leads to a very small field gradient at the nuclei thus eliminating the quadrupolar broadening of the resonance.

Compound 2c is noteworthy in other respects as well. Its cation is one of the few examples of silylated ammonium species. Silylamines, disilazanes and trisilylamines are all exhibiting very low basicity, and hence quaternization is generally not a common process. If it occurs, it is often followed by desilylation leaving behind again only triply bonded nitrogen compounds. It is only in systems stabilized by chelation that silvlated, quadruply bonded nitrogen is observed, preferably in fused rings or cages.⁴¹ In the present case (2c) the system appears to be stabilized by the intimate aggregation of the three gold atoms in a tightly bound metal triangle. Units [(AuL)₃NR]⁺ have been structurally characterized for several substituents R and phosphine ligands L. Their geometrical details are proof for the above assumption of NAu_3 aggregation through bonding $Au \cdots Au$ contacts.² Small Au-N-Au angles and short Au · · · Au distances are diagnostic for the bonding situation in these clusters.

Crystal Structures of Selected Compounds (1a, 1c and 3).— Single crystals were obtained for compounds 1a, 1c and 3. All are ionic materials, with well separated cations and anions in the lattices. The individual cations of 1a and 1c have the expected configuration, with the primary amines as donor ligands to gold(1). The metal atoms are linearly two-co-ordinate with a tertiary phosphine as the second ligand (Figs. 1 and 2). Groups N-Au-P are not very common, and it was only very recently that complexes of this type could be confirmed by structural data.

In agreement with results reported for related compounds, the cations of compound 1c are not associated in the crystal, because steric effects of the bulky Bu' and MePh₂P groups prevent the intimate approach required for metal-metal contacts. In the lattice of compound 1a, however, the cations are associated into pairs to give a short intermolecular Au \cdots Au distance of 3.047(1) Å. The smaller Me₃P ligand (as compared

Fig. 1 Dimeric structure of compound 1a. Selected bond lengths (Å) and angles (°): $Au(1) \cdots Au(2) 3.047(1)$, Au(1)-P(1) 2.236(4), Au(1)-N(1) 2.13(1), Au(2)-P(2) 2.235(3), Au(2)-N(2) 2.11(1); N(1)-Au(1)-P(1)175.7(4), N(2)-Au(2)-P(2) 174.8(3), N(1)-Au(1)-Au(2) 86.6(3), P(1)-Au(1)-Au(2) 96.8(1), N(2)-Au(2)-Au(1) 86.6(3), P(2)-Au(2)-Au(1)97.8(1)





Fig. 2 Monomeric structure of compound 1c. Selected bond lengths (Å) and angles (°): Au-N 2.105(8), Au-P 2.235(3); N-Au-P 176.4(2)

to $MePh_2P$) allows a closer aggregation, which is typical for many LAuX complexes with sterically non-demanding ligands L. Compound 1a is probably the first case, however, where an amine and a phosphine form the ligand sphere of the metals.

The cations of compound 3 feature a benzylamino group bearing two Me₃PAu substituents at the quaternary nitrogen centre. As expected for element-bridged dinuclear gold(1) compounds, the angle Au-N-Au 99.9(3)° is much smaller than the standard tetrahedral angle of 109°, clearly indicating Au ... Au bonding. This situation resembles that in cations $[(AuL)_2OR]^+$, $[(AuL)_2SR]^+$ and $[(AuL)_2SeR]^+$ which are (valence) isoelectronic.⁴²⁻⁴⁵ The individual cationic units of compound 3 are associated into centrosymmetrical dimers, as shown in Fig. 3. The two intramolecular Au...Au bonds $[Au(1) \cdots Au(2) \quad 3.171(1)]$ and the two intermolecular contacts $[Au(1) \cdots Au(2') 3.143(1)]$ form the edges of the rectangular skeleton of gold atoms in the dication. This type of aggregation to give a rectangle of gold atoms is again analogous to the configuration of the related oxonium and sulfonium cations (above). It is only for very small ligands that an even more intimate clustering is observed, where the gold atoms join up to form a tetrahedron.⁴⁶

There is no evidence for hydrogen bonding originating from the NH groups in the aggregate of compound 3, and from the NH₂ groups in compounds 1a and 1c. The bond distances and angles of the ligands L and the amino groups $RNH_{(2)}$ of all three compounds show no anomalies. Selected data are given in the figure captions.

Experimental

All experiments were carried out under pure, dry nitrogen. Solvents were purified, dried, and stored over molecular sieves in a nitrogen atmosphere. The NMR spectra were recorded in CDCl₃ as solvent, with SiMe₄ as internal standard for ¹H, ²⁹Si and ¹³C-{¹H}, H₃PO₄ (85%) as external standard for ³¹P-{¹H}, on JEOL GX 270 and GX 400 spectrometers. Mass spectra were obtained on a Varian MAT 90 spectrometer.

(tert-Butylamine)(trimethylphosphine)gold(1) Tetrafluoroborate 1a.—To a solution of [Au(PMe₃)]BF₄ {freshly prepared from [Au(PMe₃)Cl] (0.17 g, 0.55 mmol) and AgBF₄ (0.11 g, 0.56 mmol)} in tetrahydrofuran (20 cm³) was added NH₂Bu^t (0.1 cm³) at 0 °C. After stirring the solution for 2 h the crude product was precipitated with pentane. Compound 1a was purified from a dichloromethane solution by carefully precipitating with diethyl ether. Colourless crystals, yield 0.17 g (77%), decomp. 152 °C (Found: C, 20.60; H, 5.60; N, 3.90. C₇H₂₀AuBF₄NP requires C, 20.90; H, 5.00; N, 3.50%). NMR: ¹H, δ 1.3 (CCH₃, s), 1.6 [PCH₃, d, J(PH) = 11.0] and 3.7 (NH₂, s); ¹³C-{¹H}, δ 14.9 [PCH₃, d, J(PC) = 41.4 Hz], 31.4 (CCH₃, s) and 53.8



Fig. 3 Dimeric structure of compound 3. Selected bond lengths (Å) and angles (°): $Au(1) \cdots Au(2) 3.171(1)$, $Au(1) \cdots Au(2') 3.143(1)$, Au(1)-N 2.071(8), Au(2)-N 2.073(8), Au(1)-P(1) 2.246(3), Au(2)-P(2) 2.248(3); Au(1)-Au(2)-Au(1') 91.3(1), Au(2)-Au(1)-Au(2') 88.7(1), Au(1)-N-Au(2) 99.9(3)

(CCH₃, s); ³¹P-{¹H}, δ -9.7 (s). FAB mass spectrum: m/z = 346.4 (100%).

(Benzylamine)(trimethylphosphine)gold(1) Tetrafluoroborate **1b**.—Procedure as described for compound **1a**, a solution of [Au(PMe₃)]BF₄ {freshly prepared from [Au(PMe₃)Cl] (0.15 g, 0.49 mmol) and AgBF₄ (0.10 g, 0.54 mmol)} in tetrahydrofuran (20 cm³) being treated with NH₂(CH₂Ph) (0.1 cm³). Colourless oil, yield 0.13 g (57%) (Found: C, 25.70; H, 3.35; N, 3.10. C₁₀H₁₈AuBF₄NP requires C, 25.70; H, 3.90; N, 3.00%). NMR: ¹H, δ 1.5 [PCH₃, d, J(PH) = 11.6], 3.7 (NH₂, s), 4.2 (CH₂, s) and 7.3–7.5 (aryl, m); ¹³C-{¹H}, δ 14.6 [PCH₃, d, J(PC) = 41.4 Hz], 48.9 (CH₂, s), 128.3 (C⁵, s), 128.8 (C^{4.6}, s), 129.1 (C^{3.7}, s) and 138.4 (C², s); ³¹P-{¹H}, δ –10.5 (s). FAB mass spectrum: m/z = 380.3 (100%).

(tert-Butylamine)(methyldiphenylphosphine)gold(1) Tetrafluoroborate 1c.—Procedure as described for compound 1a, a solution of [Au(PPh₂Me)]BF₄ {freshly prepared from [Au(PPh₂Me)Cl] (0.31 g, 0.72 mmol) and AgBF₄ (0.14 g, 0.72 mmol)} in tetrahydrofuran (30 cm³) being treated with NH₂Bu^t (0.1 cm³). Colourless crystals, yield 0.37 g (92%), decomp. 175 °C (Found: C, 36.60; H, 4.45; N, 2.05. C₁₇H₂₄AuBF₄NP requires C, 36.65; H, 4.35; N, 2.50%). NMR: ¹H, δ 1.3 (CCH₃, s), 2.1 [PCH₃, d, J(PH) = 10.6], 4.3 (NH₂, s) and 7.4–7.6 (aryl, m); ¹³C-{¹H}, δ 14.0 [PCH₃, d, J(PC) = 43.4 Hz], 31.9 (CCH₃, s), 54.1 (CCH₃, s) and 129.5–132.9 (aryl, m); ³¹P-{¹H}, δ 16.0 (s). FAB mass spectrum: m/z = 470.2 (35%).

μ-Benzylamido-bis[(trimethylphosphine)gold(1)] Tetrafluoroborate 3.—Slow removal of the solvent from a solution of compound 2b in humid dichloromethane within 5 d afforded 3 as a colourless crystalline product. ³¹P-{¹H} NMR: δ – 10.6 (s). FAB mass spectrum: m/z = 652.2 (22%).

 $μ_3$ -tert-Butylimido-tris[(trimethylphosphine)gold(I)] Tetrafluoroborate **2a**.—To a solution of [{Au(PMe_3)}₃O]⁺BF₄⁻ (0.12 g, 0.13 mmol) in dichloromethane (20 cm³) was added NH₂Bu⁺ (0.05 cm³). The solution was initially kept at 0 °C for 1 h, then allowed to warm to room temperature and stirred for 3 h. After reducing the volume to 10 cm³ in a vacuum, diethyl ether was added to precipitate the solids. The pure, crystalline product was obtained at -25 °C from a dichloromethane solution by carefully adding pentane. Colourless crystals, yield 0.06 g (47%), decomp. 135 °C (Found: C, 15.90; H, 3.45; N, 1.75. C₁₃H₃₆Au₃BF₄NP₃ requires C, 16.00; H, 3.70; N, 1.45%). NMR: ¹H, δ 1.3 (CCH₃, s) and 1.6 [PCH₃, d, J(PH) = 10.0]; ¹³C-{¹H}, δ 15.5 [PCH₃, d, J(PC) = 37.7 Hz], 43.4 (CCH₃, s) and 63.7 (CCH₃, s); $^{31}P\text{-}\{^{1}H\}, \delta$ -11.2 (s). FAB mass spectrum: m/z = 890.6 (61.2%).

 $\mu_{3}\text{-}\textit{Benzylimido-tris}[(\textit{trimethylphosphine})gold(I)] \textit{Tetrafluoro-}$ borate 2b.--Procedure as described for compound 2a, [{Au- $(PMe_3)_3O]^+BF_4^-$ (0.24 g, 0.26 mmol) being treated with NH₂(CH₂Ph) (0.07 cm³). Colourless crystals, yield 0.26 g (79%), decomp. 146 °C (Found: C, 18.90; H, 3.45; N, 1.45. $C_{16}H_{34}Au_3BF_4NP_3$ requires C, 19.00; H, 3.40; N, 1.40%). NMR: ¹H, δ 1.5 [PCH₃, d, J(PH) = 10.4], 4.8 [CH₂, q, J(PH) = 4.9¹³C-{¹H}, δ 126.7 (C⁵, s) ³¹P-{¹H}, δ

 μ_3 -Trimethylsilylimido-tris[(trimethylphosphine)gold(I)] Tetrafluoroborate 2c.—Procedure as described for compound 2a, $[{Au(PMe_3)}_3O]^+BF_4^-$ (0.30 g, 0.32 mmol) being treated with NH(SiMe_3)₂ (0.05 cm³). Colourless crystals. NMR: ¹H, δ 1.5 [PCH₃, d, J(PH) = 10.7] and 0.1 (SiCH₃, s); ¹³C-{¹H}, δ 8.1 (SiCH₃, s) and 15.7 [PCH₃, d, J(PC) = 39.1]; ²⁹Si, δ 6.4 [SiCH₃, dct, J(SiH) = 7.4]; ³¹P-{¹H}, δ -12.1 [PAuN, t, J(PC) = 22.8 [PAuN, t, J(PC)] J(PN) = 22.8 Hz]. FAB mass spectrum: m/z = 906.3 (100) and 618.2 (73%).

gold(I)] Tetrafluoro- $\{0,1\}_{3}O]^{+}BF_{4}^{-}(0.22 \text{ g},$ was added NH_3 (0.10 -55 °C for 3 h, then

Table I C	rystallographic data for compounds 1a	, 1c and 3		
		1a	1c	3
	Formula M	C ₇ H ₂₀ AuBF ₄ NP 432.99	C ₁₇ H ₂₄ AuBF ₄ NP 557.12	C ₁₃ H ₂₆ Au ₂ BF ₄ NP ₂ 739.03
	Crystal system	Monoclinic	Orthorhombic	Monoclinic
	Space group	<i>C</i> 2/ <i>c</i> (no. 15)	<i>Pccn</i> (no. 56)	$P2_1/n$ (no. 14)
	a/Å	18.242(2)	12.246(2)	9.783(1)
	b/Å	11.058(1)	16.577(1)	13.512(1)
	c/Å	28.902(4)	20.215(2)	15.759(1)
	β/°	106.05(2)	90	92.38(1)
	$U/Å^3$	5602.9	4103.6	2081.4
	$D_{\rm c}/{\rm g~cm^{-3}}$	2.04	1.80	2.36
	Z	16	8	4
	$\mu(Mo-K\alpha)/cm^{-1}$	106.3	72.8	142.6
	Crystal dimensions/mm	$0.18 \times 0.25 \times 0.35$	$0.30 \times 0.45 \times 0.45$	$0.07 \times 0.08 \times 0.40$
	<i>T</i> ∕°C	-62	- 68	25
	Absorption correction	Empirical	Empirical	None
	T_{\min}, T_{\max} (%)	55.96, 99.80	66.89, 99.96	
	Scan mode	ω	$\theta - \theta$	$\theta - \theta$
	Scan range θ/°	3–26	2–27	2-27
	hkl ranges	$0-22, 0-13, \pm 35$	0-15, 0-21, 0-25	± 12, 0–17, 0–20
	Measured data	5952	4858	4680
	Unique data	4743	3864	3951
	Observed data $[F_o \ge 4\sigma(F_o)]$	3535	3065	3154
	No. of parameters	207	200	204
	R	0.051	0.053	0.038
	R'	0.047	0.051	0.036
	Weighting scheme, w	$\left[\sigma^2(F_o) +\right]$	$[\sigma^2(F_0) +$	$[\sigma^2(F_o) +$
		$0.000\ 100\ F_0^2]^{-1}$	$0.000\ 245\ F_0^2$] ⁻¹	$0.000\ 379\ F_0^2]^{-1}$
	Residual electron density/e Å ⁻³	1.42 2.07	1.84 1.21	1.78 0.80

Table 1 Crys

 Table 2 Positional parameters for the non-hydrogen atoms of compound 1a

Atom	x	у	z	Atom	x	у	z
Au(1)	-0.07395(3)	-0.15103(6)	-0.59916(2)	$C(142)^{b}$	-0.094(1)	-0.188(2)	-0.5003(7)
Au(2)	0.008 88(3)	-0.355 00(6)	-0.63474(2)	$C(152)^{b}$	-0.167(1)	-0.123(2)	-0.5141(7)
P(1)	-0.0863(2)	-0.0287(4)	-0.6624(1)	C(162) ^b	-0.033(1)	-0.092(2)	-0.4786(7)
P(2)	0.127 4(2)	-0.2830(4)	-0.6069(1)	$C(172)^{b}$	-0.100(1)	-0.278(2)	-0.4611(7)
N(1)	-0.0704(7)	-0.263(1)	-0.5388(4)	B(1)	0.815(2)	0.195(3)	0.181(1)
N(2)	-0.0987(6)	-0.439(1)	-0.6599(4)	F (1)	0.738 7(7)	0.218(1)	0.184 8(4)
C(21)	0.138 7(9)	-0.169(2)	-0.5632(6)	F(2)	0.859 9(8)	0.169(1)	0.220 4(5)
C(22)	0.164 4(9)	-0.218(2)	-0.6623(5)	F(3)	0.830 1(8)	0.318(1)	0.167 8(5)
C(23)	0.192 7(9)	-0.399(2)	-0.5805(7)	F(4)	0.811 9(9)	0.118(1)	0.146 1(6)
C(24)	-0.1104(7)	-0.522(1)	-0.7014(5)	B (21) ^c	0.144(1)	0.325(2)	0.035 9(7)
C(25)	-0.048 0(9)	-0.612(1)	-0.6936(5)	F(11) °	0.171(1)	0.227(2)	0.016 8(7)
C(26)	-0.1874(9)	-0.585(1)	-0.7071(6)	F(21) ^c	0.150(1)	0.303(2)	0.083 4(7)
C(27)	-0.1148(8)	-0.446(1)	-0.7453(5)	F(31) ^c	0.185(1)	0.427(2)	0.031 9(7)
C(11)	-0.096(1)	-0.106(1)	-0.7183(5)	F(41) °	0.069(1)	0.341(2)	0.011 0(7)
C(12)	-0.168(1)	0.066(2)	-0.6730(5)	$\mathbf{B}(22)^d$	0.110 6(6)	0.359(1)	0.046 7(4)
C(13)	-0.009(1)	0.068(1)	-0.6598(6)	$F(12)^{d}$	0.084 6(6)	0.451(1)	0.069 7(4)
C(141) ^a	-0.120(1)	-0.240(2)	-0.5117(7)	$F(22)^d$	0.089 4(6)	0.250(1)	0.061 7(4)
C(151) ^a	-0.116(1)	-0.328(2)	-0.471 5(7)	F(32) ^d	0.188 5(6)	0.364(1)	0.057 5(4)
C(161) ^a	-0.200(1)	-0.243(2)	-0.546 5(7)	$F(42)^{d}$	0.079 8(6)	0.370(1)	-0.0020(4)
C(171) ^a	-0.103(1)	-0.112(2)	-0.4908(7)				

^a Split atom (s.o.f. = 0.48). ^bSplit atom (s.o.f. = 0.52). ^c Split atom (s.o.f. = 0.25). ^d Split atom (s.o.f. = 0.75).

Table 3 Positional parameters for the non-hydrogen atoms of compound 1c

Atom	x	у	Ξ
Au	0.090 31(3)	0.120 73(2)	0.350 66(2)
Р	0.179 3(2)	0.103 7(2)	0.446 2(1)
N	0.014 9(6)	0.133 3(5)	0.257 6(4)
C(11)	0.323 9(8)	0.123 4(6)	0.434 5(5)
C(12)	0.399 3(9)	0.102 4(7)	0.482 3(5)
C(13)	0.509 0(9)	0.118 0(7)	0.473 0(5)
C(14)	0.544 5(9)	0.153 8(6)	0.414 0(6)
C(15)	0.469 5(9)	0.173 7(6)	0.365 9(5)
C(16)	0.360 0(9)	0.159 9(6)	0.377 4(5)
C(21)	0.171 3(8)	0.000 5(7)	0.474 2(6)
C(22)	0.165(1)	-0.0200(9)	0.540 4(7)
C(23)	0.164(1)	-0.099(1)	0.559(1)
C(24)	0.165(1)	-0.160(1)	0.516(1)
C(25)	0.169(1)	-0.139 6(8)	0.448(1)
C(26)	0.173(1)	-0.0601(7)	0.427 6(7)
C(3)	0.135 8(9)	0.167 5(7)	0.512 6(6)
C(41)	-0.0022(9)	0.055 3(7)	0.220 1(5)
C(42)	-0.052(1)	0.079 2(9)	0.153 5(6)
C(43)	-0.077(1)	0.004 3(9)	0.258 4(7)
C(44)	0.108(1)	0.016 3(9)	0.210 7(7)
B (1)	0.25	0.25	0.184 7(8)
B (2)	0.25	0.75	0.237(1)
F(1)	0.204(1)	0.190(1)	0.146 9(8)
F(2)	0.173(1)	0.277(1)	0.229 3(8)
F(3)	0.168(1)	0.780(1)	0.276(1)
F(4)	0.207(1)	0.691(1)	0.196(1)

allowed to warm to 0 °C and stirred for 3 h. After reducing the volume to 10 cm³ in a vacuum, diethyl ether was added to precipitate the solids. Pure product was obtained from an acetonitrile solution by carefully adding pentane.

(b) Compound 4 was also obtained by reacting 2c with $[{Au(PMe_3)}_3O]^+BF_4^-$ for 20 h at room temperature in humid tetrahydrofuran, but in lower yields compared with route (a). Colourless. Yield 0.11 g (51%) [route (a)], decomp. 122 °C (Found: C, 12.30; H, 3.15; N, 1.00. C₁₂H₃₆Au₄BF₄NP₄ requires C, 12.05; H, 3.05; N, 1.15%). NMR: ¹H, δ 1.6 [PCH₃, d, J(PH) = 10.8]; ¹³C-{¹H}, δ 15.8 [PCH₃, d, J(PC) = 37.2 Hz]; ³¹P-{¹H}, δ -15.1 (s). FAB mass spectrum: m/z = 1106.0 (84%).

Crystal Structure Determinations.—All samples were mounted in glass capillaries. Graphite-monochromated Mo-K α radiation was used. The structures were solved by direct methods (SHELXTL PLUS).⁴⁷ That of compound **1a** was complicated by disorder of Bu' and BF₄ groups. These distributions were accounted for by using split models of rigid groups with site occupation factors (s.o.f.s) of 0.25 and 0.75 for one BF₄ group and 0.48 and 0.52 for one Bu' group. Rigid-group refinement was also used for the BF₄ in **1c**. An empirical absorption correction was not carried out after data collection for compound **3** because of some decomposition of the crystal during the measurements of the intensity data. Hydrogen atoms of non-disordered groups were included in idealized, fixed positions. The final unit-cell parameters and specific data collection parameters are summarized in Table 1; atomic coordinates are given in Tables 2–4.

Additional material available from the Cambridge Crystallographic Data Centre comprises H-atom coordinates, thermal parameters and remaining bond lengths and angles.

Acknowledgements

This work was supported by the Deutsche Forschungsgemeinschaft, the Fonds der Chemischen Industrie and through the donation of chemicals by Degussa AG and Heraeus GmbH.

Table 4Positional parameters for the non-hydrogen atoms ofcompound 3

Atom	x	У	Z
Au(1)	0.074 40(4)	0.841 98(3)	0.507 11(2)
Au(2)	0.199 74(4)	1.052 88(3)	0.552 18(2)
P(1)	0.081 1(3)	0.728 5(2)	0.403 2(2)
P(2)	0.331 6(3)	1.170 0(2)	0.495 8(2)
N(1)	0.084 8(9)	0.942 5(6)	0.606 4(5)
C(11)	-0.007(1)	0.762 0(9)	0.304 1(7)
C(12)	0.011(1)	0.609 4(8)	0.433 2(8)
C(13)	0.255(1)	0.698(1)	0.371 6(8)
C(21)	0.255(1)	1.229(1)	0.402 4(8)
C(22)	0.373(1)	1.268 3(9)	0.572 3(9)
C(23)	0.498(1)	1.128 3(9)	0.465 4(8)
C(1)	0.141(1)	0.898 8(8)	0.688 1(7)
C(2)	0.165(1)	0.977 9(8)	0.756 5(7)
C(3)	0.290(1)	1.029 2(9)	0.761 5(9)
C(4)	0.312(2)	1.102(1)	0.821(1)
C(5)	0.215(2)	1.124(1)	0.878(1)
C(6)	0.090(2)	1.070(1)	0.876 5(8)
C(7)	0.064(1)	0.995(1)	0.813 1(8)
В	0.683(2)	0.037(1)	0.718(1)
F(1)	0.613(1)	-0.050 8(6)	0.727 8(6)
F(2)	0.790(1)	0.024(1)	0.673 0(8)
F(3)	0.712(2)	0.074(1)	0.792 3(9)
F(4)	0.591(1)	0.096(1)	0.676 0(9)

We thank Professor F. R. Kreissl for recording the mass spectra and Mr. J. Riede for collecting the X-ray data sets.

References

- R. J. Puddephatt, in *Comprehensive Coordination Chemistry*, eds. G. Wilkinson, R. D. Gillard and J. A. McCleverty, Pergamon, Oxford, 1987, ch. 55.9 and refs. therein.
- 2 A. Grohmann, J. Riede and H. Schmidbaur, J. Chem. Soc., Dalton Trans., 1991, 783.
- 3 J. P. Vollano, D. H. Pickler and J. A. Statler, *Inorg. Chim. Acta*, 1989, 155, 31.
- 4 E. M. Kinsch and D. W. Stephan, Inorg. Chim. Acta, 1984, 91, 263.
- 5 R. Uson, A. Laguna and M. D. Villacampa, *Inorg. Chim. Acta*, 1984, 81, 25.
- 6 E. G. Perevalova, K. I. Grandberg, E. I. Smyslova, L. G. Kuz'mina, V. I. Korsunskii and D. N. Kravtsov, *Organomet. Chem. USSR*, 1989, **2**, 523.
- 7 A. Grohmann and H. Schmidbaur, Inorg. Chem., 1992, 31, 3378.
- 8 H. Schmidbaur, A. Kolb and P. Bissinger, *Inorg. Chem.*, 1992, 31, 4370.
- 9 V. Ramamoorthy and P. R. Sharp, Inorg. Chem., 1990, 29, 3336.
- 10 Yi. Yang, V. Ramamoorthy and P. R. Sharp, *Inorg. Chem.*, 1993, 32, 1946.
- 11 L. G. Kuz'mina, Zh. Neorg. Khim., 1993, 38, 994.
- 12 E. Zeller, H. Beruda, A. Kolb, P. Bissinger, J. Riede and H. Schmidbaur, *Nature (London)*, 1991, **352**, 141.
- 13 Yu. L. Slovokhotov and Yu. T. Struchkov, J. Organomet. Chem., 1984, 277, 143.
- 14 E. G. Perevalova, E. I. Smyslova, V. P. Dyadchenko, K. I. Grandberg and A. N. Nesmeyanov, *Izv. Akad. Nauk SSSR, Ser. Khim.*, 1980, 1455.
- 15 A. Grohmann, J. Riede and H. Schmidbaur, *Nature (London)*, 1990, 345, 140.
- 16 H. Schmidbaur, Gold Bull., 1990, 23, 11.
- 17 H. Schmidbaur, Interdiscip. Sci. Rev., 1992, 17, 213.
- 18 H. Schmidbaur, Pure Appl. Chem., 1993, 65, 691.
- 19 K. Dziwok, J. Lachmann, D. L. Wilkinson, G. Müller and H. Schmidbaur, *Chem. Ber.*, 1990, **123**, 423.
- 20 H. Schmidbaur, K. Dziwok, A. Grohmann and G. Müller, Chem. Ber., 1989, 122, 893.
- 21 H. Schmidbaur, W. Graf and G. Müller, Angew. Chem., Int. Ed. Engl., 1988, 27, 417.
- 22 R. Narayanaswamy, M. A. Young, E. Parkhurst, M. Ouellette, M. E. Kerr, D. M. Ho, R. C. Elder, A. E. Bruce and M. R. M. Bruce, *Inorg. Chem.*, 1993, 32, 2506.
- 23 J. Li and P. Pyykkö, Chem. Phys. Lett., 1992, 197, 586.

- 24 J. Li and P. Pyykkö, Inorg. Chem., 1993, 32, 2630.
- 25 P. Pyykkö and Y.-F. Zhao, Angew. Chem., Int. Ed. Engl., 1991, 30, 604.
- 26 P. Pyykkö, Chem. Rev., 1988, 88, 563.
- 27 P. Pyykkö and J. P. Desclaux, Acc. Chem. Res., 1979, 12, 276.
- 28 H. Schmidbaur, G. Weidenhiller, O. Steigelmann and G. Müller, Z. Naturforsch., Teil B, 1990, 47, 747.
- 29 H. Schmidbaur, G. Weidenhiller, O. Steigelmann and G. Müller, Chem. Ber., 1990, 123, 285.
- 30 S. Ahrland, B. Aurivillius, K. Dreisch, B. Noren and A. Oskarsson, Acta Chem. Scand., 1992, 46, 262.
- 31 H. Schmidbaur, K. Angermaier and E. Zeller, J. Organomet. Chem., 1994, 472, 371.
- 32 D. B. Dyson, R. V. Parish, C. A. McAuliffe, D. G. Pritchard, R. Fields and B. Beagley, J. Chem. Soc., Dalton Trans., 1989, 907.
- 33 S. Ahrland, K. Dreisch, B. Noren and A. Oskarsson, Acta Chem. Scand., Ser. A, 1987, 41, 499.
- 34 P. Pyykkö, J. Li and N. Runeberg, Chem. Phys. Lett., 1994, 218, 133.
- 35 K. Angermaier and H. Schmidbaur, unpublished work.36 F. Scherbaum, A. Grohmann, B. Huber, C. Krüger and
- H. Schmidbaur, Angew. Chem., Int. Ed. Engl., 1988, 27, 1544.
 F. Scherbaum, A. Grohmann, G. Müller and H. Schmidbaur, Angew. Chem., Int. Ed. Engl., 1989, 28, 463.

- J. CHEM. SOC. DALTON TRANS. 1995
- 38 F. Scherbaum, B. Huber, G. Müller and H. Schmidbaur, Angew. Chem., Int. Ed. Engl., 1988, 27, 1542.
- 39 E. Zeller and H. Schmidbaur, J. Chem. Soc., Chem. Commun., 1993, 69.
- 40 A. Blumenthal, H. Beruda and H. Schmidbaur, J. Chem. Soc., Chem. Commun., 1993, 1005.
- 41 M. Veith, F. Goffing, S. Becker and V. Huck, J. Organomet. Chem., 1991, **406**, 105.
- 42 Y. Yang, V. Ramamoorthy and P. R. Sharp, *Inorg. Chem.*, 1993, 32, 1946.
- 43 P. G. Jones, G. M. Sheldrick and E. Hädicke, *Acta Crystallogr.*, Sect. B, 1980, 36, 2777.
- 44 K. Angermaier and H. Schmidbaur, Chem. Ber., 1994, 127, 2387.
- 45 S. Wang and J. P. Fackler, Inorg. Chem., 1990, 29, 4404.
- 46 K. Angermaier and H. Schmidbaur, Inorg. Chem., 1994, 33, 2069.
- 47 G. M. Sheldrick, SHELXTL PLUS, release 4.0 for Siemens R3 Crystallographic Research Systems, Siemens Analytical X-Ray Instruments, Madison, WI, 1989.

Received 8th July 1994; Paper 4/04161E