CHEMISTRY LETTERS, pp. 2049-2052, 1986.

Asymmetric Microbial Oxidation of Formaldehyde Dithioacetals¹⁾

Yasushi OKAMOTO, Hiromichi OHTA,^{*} and Gen-ichi TSUCHIHASHI Department of Chemistry, Faculty of Science and Technology, Keio University, Hiyoshi 3-14-1, Kohoku-ku, Yokohama 223

Incubation of formaldehyde dithioacetals with growing cells of <u>Corynebacterium equi</u> IFO 3730 afforded a variety of oxidation products depending on the structure of the substrates. The products with sulfoxide moiety were optically pure and had <u>R</u> configuration.

A dithioacetal group serves not only as a protecting group of carbonyl compounds, but also as a "masked acyl anion", because two mercapto groups stabilize the central methylenic anion.²⁾ Thus, it will be potentially interesting to introduce an asymmetric center to a dithioacetal group because the resulting compound will serve as a chiral acyl anion equivalent. It has been already demonstrated by Ogura <u>et al</u>. that the chirality on sulfur atom of dithioacetals can be transferred to a carbon atom.³⁾

Microbial transformations have been demonstrated to be often effective for the introduction of chirality to prochiral compounds. Hitherto, only a few examples of enzymatic oxidation of dithioacetals have been demonstrated, and the optical purities of the products were not enough for the synthetic application.⁴⁾

Studies on microbial transformation in our laboratory have shown that <u>Coryne-bacterium</u> <u>equi</u> IFO 3730 has an enzyme system which catalyzes the oxidation of various sulfides to optically active sulfoxides and sulfones.⁵⁾ Now, we wish to report the results of application of this system to oxidation of thioacetals.

In a 500-ml Sakaguchi flask, were added 90 ml of the sterilized inorganic medium,⁶⁾ 2 ml of hexadecane, and 0.1 to 0.4 ml of a dithioacetal <u>1</u>. The mixture was inoculated with 10 ml of a suspension of grown cells of <u>C</u>. <u>equi</u>, and incubated for 3 to 7 days at 30 ^OC on a reciprocatory shaker. The broth was extracted with a 100 ml portion of ethyl acetate three times, and the combined extracts were concentrated under reduced pressure. Separation of the remaining hexadecane by column chromatography on silica gel gave a mixture of the substrate and the oxidized products amounting to almost quantitative recovery of sulfur-containing compounds. The oxidation products were isolated and purified by preparative TLC on silica gel and identified by comparing their ¹H-NMR, IR, and mass spectra with those of authentic specimens.⁷⁾ The major products are summarized in Table 1.

A variety of products were obtained depending on the structure of substrates. The substrates which have alkyl chain with large numbers of carbons tend to be oxidized smoothly (compd b, d). The similar tendency has been observed in the oxidation of terminal olefins,⁸⁾ secondary alcohols,⁹⁾ and alkyl aryl sulfides⁵⁾ by <u>C</u>. equi. Diphenyl dithioacetal was most resistant to oxidation and afforded

the mono sulfone in a low yield (compd a). In oxidation of unsymmetric alkyl aryl dithioacetals, the sulfur atom linked to the alkyl chain was oxidized in all cases (compd d, e, f).

Oxidation of dibutyl dithioacetal and methyl phenyl dithioacetal resulted in the formation of products which have sulfoxide moiety (2c, 2f).¹⁰⁾ Both sulfoxides were confirmed to be optically pure by ¹H-NMR in the presence of chiral shift reagent (Fig. 1).

The absolute configuration of sulfoxides is determined by Pirkle's method¹¹⁾ and CD spectra.¹²⁾ Sulfoxides are known to form 1:1 solvates with (S)-(+)-2,2,2-trifluoro-1-(9-anthryl)-ethanol <u>3</u> (TFAE) (Fig. 2). The shielding effect of the

Table 1. Oxidation of Formaldehyde Dithioacetals

$H_2C_{SR}^{'SR}$	<u>C. equi</u>	^{H2C'} S(0) _m R ¹
1		2

Compd	Substrate	Concn /% ^{a)}	Cult./day	Product	Yield/%	e.e./%	Δε of CD(nm)
a	H ₂ C ^{/SPh} sph	0.1	7	H ₂ C ^{/SO} 2 ^{Ph} SPh	34 ^{b)}	-	-
b	H ₂ C ^{/SC} 8 ^H 17 SC8 ^H 17	0.2	3	H ₂ C ^{/SO₂C₈H₁₇ SO₂C₈H₁₇}	100	-	_
с	^{H2C^{/SC4^H9} SC4^H9}	0.1	7	H ₂ C ^{′SO₂C₄H₉} SOC ₄ H ₉	70	>95	-2.2(223) ^{d)} -1.9(229) ^{e)}
đ	H2C ^{/SPh} SC8H17	0.2	3	^H 2 ^{C′ SPh} SO ₂ C ₈ H ₁₇	76	-	-
e	H ₂ C ^{/SPh} SC ₄ H ₉	0.1	7	H ₂ C ^{/SPh} SO ₂ C ₄ H ₉	44 ^c)	-	-
f	H ₂ C ^{/SPh} SCH ₃	0.1	7	H ₂ C ^{/SPh} SOCH ₃	47 ^{b)}	>95	-1.1(218) ^{d)} -1.4(231) ^{e)}

- a) ml substrate / 100 ml medium
- b) The remaining part is recovered starting material.
- c) A small amount of ${\rm PhSCH}_2{\rm SOC}_4{\rm H}_9$ (12%) was identified in addition to starting material.
- d) Measured in EtOH.
- e) Measured in $C-C_6H_{12}$.

2050

anthryl group allows us to expect that the methylene proton resonances occur at higher field for the (Sc,Ss)-solvates (Fig. 2-A, 2-C) than the (Sc,Rs)-solvates (Fig. 2-В, 2-D). ¹H-NMR The signal due to the methylene group of racemic 2c split into three peaks (doublet and singlet), corresponding to two enantiomers by adding 2 mol equivalents of a chiral solvent TFAE 3. Under the same conditions <u>2c</u> obtained from microbial oxidation showed а methylene signal of doublet in the lower field due to one enantiomer.¹³⁾ These results show that the methylene



Fig. 1. $Eu(hfc)_3$ -shifted ¹H-NMR Spectra of <u>2</u>. (A) <u>2c</u>; (B) <u>2f</u> (1) microbial oxidation product; (2) racemic

ОН 3



Fig. 2. Configuration of solvates of 2 with TFAE 3.

group of the product $\underline{2c}$ is \underline{trans} to the anthryl group (Fig. 2-B), which led to the conclusion that $\underline{2c}$ has \underline{R} configuration. In a similar manner, while the signal due to methylene of racemic $\underline{2f}$ split into two sets of quartet, the one of the product $\underline{2f}$ appeared as only one set of quartet in lower field.¹³⁾ Hence, the product $\underline{2f}$ is determined to have \underline{R} configuration.

Mislow <u>et al</u>. have shown that a relation exists between the absolute configuration of alkyl methyl and alkyl aryl sulfoxides and the ORD Cotton effect. The sulfoxides which show a negative Cotton effect, centered at the absorption band of S-O group, have been demonstrated to have <u>R</u> configuration. We studied the Cotton effect with CD spectra. Compared with ORD spectra, CD spectra are independent of other absorptions, and give the simple and important informations for the study of stereochemistry. The results were summarized in Table 1. In both cases of the products <u>2c</u> and <u>2f</u>, the CD spectra showed negative values, which indicates that both sulfoxides have <u>R</u> configuration. This result agreed with the one obtained by Pirkle's method.

References

- 1) Studies on Enzymatic Oxidation of Sulfides, Part VI. For Part V, H. Ohta, Y. Kato, and G. Tsuchihashi, J. Org. Chem., submitted.
- 2) B. T. Gröbel and D. Seebach, Synthesis, 1977, 357.
- K. Ogura, M. Fujita, T. Inaba, K. Takahashi, and H. Iida, Tetrahedron Lett., 24, 503 (1983).
- 4) M. Poje, O. Nota, and K. Balenović, Tetrahedron, <u>36</u>, 1895 (1980); K. Ogura, M. Fujita, and H. Iida, Tetrahedron Lett., <u>21</u>, 2233 (1980); B. J. Auret, D. R. Boyd, F. Breen, and R. M. E. Greene, J. Chem. Soc., Perkin Trans. 1, <u>1981</u>, 930; B. J. Auret, D. R. Boyd, E. S. Cassidy, F. Turley, A. F. Drake, and S. F. Mason, J. Chem. Soc., Chem. Commun., <u>1983</u>, 282; B. J. Auret, D. R. Boyd, E. S. Cassidy, R. Hamilton, and F. Turley, J. Chem. Soc., Perkin Trans. 1, <u>1985</u>, 1547.
- 5) H. Ohta, Y. Okamoto, and G. Tsuchihashi, Chem. Lett., <u>1984</u>, 205; Agric. Biol. Chem., <u>49</u>, 671 (1985); ibid., <u>49</u>, 2229 (1985).
- 6) The medium consists of $(NH_4)_2HPO_4$ 10 g, KH_2PO_4 2 g, $MgSO_4 \cdot 7H_2O$ 0.3 g, $FeSO_4 \cdot 7H_2O$ 10 mg, $ZnSO_4 \cdot 7H_2O$ 8 mg, $MnSO_4 \cdot 4H_2O$ 8 mg, yeast extract 0.2 g, and H_2O to make 1000 ml (pH 7.2).
- 7) Authentic samples were prepared as follows: (C₈H₁₇SO₂)₂CH₂, C₄H₉SO₂CH₂SOC₄H₉, and PhSCH₂SOCH₃, by oxidation of dithioacetals with H₂O₂ in acetic acid; PhSCH₂SO₂Ph, PhSCH₂SO₂C₈H₁₇, and PhSCH₂SO₂C₄H₉, by the coupling of phenylthio-methylmagnesium chloride with alkanesulfonyl chloride; K. Ogura, M. Fujita, K. Takahashi, and H. Iida, Chem. Lett., <u>1982</u>, 1697.
- 8) H. Ohta and H. Tetsukawa, Agric. Biol. Chem., 43, 2099 (1979).
- 9) H. Ohta, H. Yamada, and G. Tsuchihashi, Agric. Biol. Chem., 49, 2459 (1985).
- 10) $C_4H_9SO_2CH_2SOC_4H_9 \ \underline{2c}$: IR (NaCl) 2915, 2848, 1450, 1315, 1295, 1025 cm⁻¹; NMR (CDCl₃) δ 1.00 (6H, t, J=7.5 Hz, 2 x CH₃), 1.64 (4H, m, 2 x CH₂CH₃), 3.00 (2H, t, J=7.2 Hz, SOCH₂CH₂), 3.26 (2H, t, J=7.2 Hz, SO₂CH₂CH₂), 4.17 (2H, s, CH₂); MS m/z (rel. intensity) 241 (37, M+1), 223 (10), 184 (40), 127 (44), 103 (63), 63 (100), 57 (45). PhSCH₂SOCH₃ <u>2f</u>: IR (NaCl) 3050, 2960, 2900, 1575, 1475, 1435, 1040, 740, 690 cm⁻¹; NMR (CDCl₃) δ 2.57 (3H, s, CH₃), 3.78 (1H, d, J=12 Hz, CH₂), 4.02 (1H, d, J=12 Hz, CH₂), 7.33 (5H, m, Ph); MS m/z (rel. intensity) 186 (1, M⁺), 123 (100), 109 (5), 77 (13), 45 (34).
- 11) W. H. Pirkle, S. D. Beare, and R. L. Muntz, Tetrahedron Lett., <u>1974</u>, 2295; W. H. Pirkle and D. L. Sikkenga, J. Org. Chem., <u>40</u>, 3430 (1975).
- 12) K. Mislow, M. M. Green, P. Laur, J. T. Melillo, T. Simmons, and A. L. Ternay, Jr., J. Am. Chem. Soc., <u>87</u>, 1958 (1965); M. Axelrod, P. Bickart, M. L. Goldstein, M. M. Green, A. Kjaer, and K. Mislow, Tetrahedron Lett., <u>1968</u>, 3249.
- 13) Chemical shift of CH₂ flanked to two sulfur atoms: <u>2c</u>; <u>3.9509</u>, <u>3.9511</u>,
 3.9401. <u>2f</u>; <u>4.0415</u>, <u>4.0073</u>, 3.9975, 3.9633, <u>3.8314</u>, <u>3.7985</u>, 3.8021, 3.7679.
 All peaks were observed for racemic compounds, while underlined peaks were strongly observed for enzymatic oxidation products.

(Received September 13, 1986)

2052