

A Journal of the Gesellschaft Deutscher Chemiker A Deutscher Chemiker GDCh International Edition www.angewandte.org

Accepted Article

Title: Efficient Palladium-Catalyzed Aerobic Arylative Carbocyclization of Enallenynes

Authors: Jan-E. Bäckvall, Jie Liu, Alexander Ricke, and Bin Yang

This manuscript has been accepted after peer review and appears as an Accepted Article online prior to editing, proofing, and formal publication of the final Version of Record (VoR). This work is currently citable by using the Digital Object Identifier (DOI) given below. The VoR will be published online in Early View as soon as possible and may be different to this Accepted Article as a result of editing. Readers should obtain the VoR from the journal website shown below when it is published to ensure accuracy of information. The authors are responsible for the content of this Accepted Article.

To be cited as: Angew. Chem. Int. Ed. 10.1002/anie.201810501 Angew. Chem. 10.1002/ange.201810501

Link to VoR: http://dx.doi.org/10.1002/anie.201810501 http://dx.doi.org/10.1002/ange.201810501

WILEY-VCH

WILEY-VCH

Efficient Palladium-Catalyzed Aerobic Arylative Carbocyclization of Enallenynes

Jie Liu, Alexander Ricke, Bin Yang,* and Jan-E. Bäckvall*

Dedicated to Prof. Elias J. Corey and Prof. Xiyan Lu on the occasion of their 90th birthday

Abstract: Herein, we communicate a selective and efficient protocol for oxidative arylating carbocyclization of enallenynes using O_2 as the oxidant. The key to success for this aerobic transformation is the application of a specific electron transfer mediator (ETM) – a bifunctional catalyst consisting of a metal-macrocycle and quinone moieties. This catalyst significantly facilitates the reoxidation of Pd(0) to Pd(II) under atmospheric pressure of O_2 . Diverse functionalized enallenynes react with aryl boronic acids to afford the corresponding cyclic tetraenes in moderate to good yields.

The use of molecular oxygen (O₂) in enzyme-catalyzed oxidation reactions is highly widespread in nature.^[1] In the perspective of synthetic organic chemistry, molecular oxygen is an inexpensive. abundant and highly atom-efficient oxidant that generates no toxic byproducts, thus fulfilling the requirements of "green chemistry".^[2] Over the past decades, palladium-catalyzed aerobic oxidations have provided the basis for streamlined conversion of various feedstocks into valuable products.^[3] Illustrative examples include Wacker oxidations,^[4] alcohol oxidations,^[5] alkene functionalizations^[6] and oxidative C-H activations.^[7] In spite of the significant synthetic progress offered by these reactions, the relative low catalyst efficiency due to palladium deactivation still remains challenging in most cases.^[8] This "Oxidation Problem" can be explained by the slow electron transfer directly between Pd(0) and O₂, compared to the rapid precipitation of palladium black from active palladium species (Scheme 1a).

To circumvent the problem of getting Pd black in direct reoxidation of Pd(0) by O_2 , considerable attention has been focused on the development of ancillary air-stable ligands such as amines, pyridines, sulfoxides, and carbene derivatives that can inhibit precipitation of Pd black during the catalytic cycle.^[9] On the other hand, inspired by nature, a coupled catalyst system with electron transfer mediators (ETMs) can facilitate the transport of the electrons from the reduced palladium catalyst to O_2 , thereby increasing the efficiency of the reoxidation of Pd(0) to Pd(II).^[10] After the original work by Bäckvall in the late 1980s,^[11] several groups have explored this concept by developing mild and efficient Pd(II)-catalyzed aerobic oxidative reactions.^[12] The key to success for these transformations is the use of macrocyclic metal complexes and quinones as ETMs under aerobic conditions. Additionally, improved coupled catalyst sys-

 [a] Dr. J. Liu, A. Ricke, Dr. B. Yang and Prof. Dr. J.-E. Bäckvall Department of Organic Chemistry, Arrhenius Laboratory, Stockholm University, SE-10691 Stockholm, Sweden
 E-mail: <u>binyang@organ.su.se</u> jeb@organ.su.se
 Homepage: <u>http://www.organ.su.se/jeb/</u>

Supporting information for this article is given via a link at the end of the document.((Please delete this text if not appropriate))

tems in which a metal-macrocycle and quinones are merged into one molecule have been reported by our group.^[13] These bifunctional catalysts led to an increased efficiency of the electron transfer compared to that observed for the system with the quinone and metal-macrocycle as separate molecules.^[13] Based on these state-of-the-art works, we present work on the use of such a bifunctional catalyst (Co(salophen)-HQ; HQ = hydroquinone, Scheme 1b) as an efficient electron transfer mediator in Pdcatalyzed aerobic oxidative carbocyclizations of enallenynes.



Scheme 1. a) The "Oxidation Problem" in palladium-catalyzed aerobic reactions and the solutions. b) The cobalt-based bifunctional catalyst (Co(salophen)-HQ) as an electron transfer mediator in this study.

Allenes constitute an important class of synthons in organic synthesis, which can be applied to construct a variety of valuable molecules.^[14] Our group has a long-standing interest in Pd-catalyzed C-C bond-forming reactions from allenesubstituted unsaturated compounds, such as enallenes, allenynes, and bisallenes.^[15] In the present work, we report for the first time an example of Pd-catalyzed oxidative carbocyclization of rationally designed enallenynes. These interesting molecules contain three different C-C m-bond functionalities (allene, olefin and alkyne groups), all of which can be potential reaction sites.^[16] However, the presence of these three π systems provides a challenge concerning control of regioselectivity in the reaction. As shown in Scheme 2, there are three possible pathways for the transformation of vinylpalladium species Int-A, which is generated from an initial allenic C-H cleavage^[15] of enallenyne 1 by Pd(II): (1) According to our previous work,^[17] the close-by olefin insertion of Int-A could lead to a highly strained cyclobutene complex Int-B, followed by arylation to give a fourmembered carbocycle 3' (Path I). (2) Intermediate Int-A can also be directly trapped by a nucleophile such as ArB(OH)₂ to give an acyclic arylated product 3" (Path II).[18] (3) A ligand exchange of the olefin for alkyne in Int-A would give Int-C, which on subsequent alkyne insertion can generate six-

membered ring species *Int-D*. Trapping of *Int-D* by ArB(OH)₂ would lead to cyclic tetraene **3** (Path III).



Scheme 2. Possible pathways for palladium-catalyzed oxidative functionalization (arylation) of enallenynes.

With these possibilities in mind, our goal was to develop a general and efficient catalyst system for oxidative functionalization of enallenynes in a selective manner. More specifically, in light of recent growing interest in green and sustainable chemistry, we have undertaken preliminary investigations to use molecular oxygen as terminal oxidant.

At the outset of our investigations, the palladium-catalyzed aerobic oxidation of a readily accessible enallenyne 1a and phenylboronic acid 2a in H2O/acetone was chosen as the benchmark reaction (Table 1).^[19] In the absence of ETMs, we did not observe any formation of the arylated product under aerobic conditions and the starting material 1a could be recovered in 93% yield (entry 1). When catalytic amounts of BQ (p-benzoquinone) was added, a highly unsaturated sixmembered carbocycle 3a was selectively formed albeit in low yield (15%, entry 2).^[20] It is noteworthy that this unexpected cyclization takes place between the allene moiety and distal triple bond of enallenyne 1a, while the olefin group remains intact.^[21] This interesting result motivated us to further explore various established metal-based electron transfer mediators to improve the reaction efficiency. The application of VO(acac)₂, Fe(Pc), and Co(Pc) (Pc = phthalocyanine) as ETMs did not lead to any significant improvement of the transformation of 1a to 3a. (21%-24% yields, entries 3-5). To our delight, the use of cobaltbased salen and salophen complexes led to good yields of 3a (71 and 74% yields, entries 6 and 7). As to the reduced state of BQ, HQ (hydroquinone) also furnished the cyclization product 3a in 63% yield (entry 8). Interestingly, a bifunctional cobalt catalyst (Co(salophen)-HQ), combining a metal-macrocycle and quinones moieties, was found to be the best performing catalyst and afforded the desired product 3a in 79% yield (entry 9). Notably, even under lower catalyst loading (2 mol% Pd(OAc)₂ and 5 mol% Co(salophen)-HQ), no significant decrease in yield was observed (76% yield, entry 10 vs. 79% yield, entry 9).

Table 1. Evaluation of different electron transfer mediators (ETMs) for oxidative carbocyclization of enallenyne ${\bf 1a.}^{[a]}$







[a] Unless otherwise noted, the following reaction condition were employed with 1a (0.1 mmol, 1.0 equiv.), 2a (0.15 mmol, 1.5 equiv.), Pd(OAc)₂ (5 mol%), ETM₁ (10 mol%), ETM₂ (20 mol%), H₂O (1.0 equiv.) in 0.1 M acetone, O₂ (1 atm) at room temperature (23 °C) for 48 h. Yield and conversion are determined by ¹H NMR using anisole as internal standard. [b] 10 mol% Co(salophen)-HQ was added. [c] 5 mol% Co(salophen)-HQ with 2 mol% Pd(OAc)₂ was added.



Figure 1. Reaction progress with different ETMs.

Next, the catalytic reaction progress with different electron transfer mediators (ETMs) was examined using 1 mol% of Pd(OAc)₂, 2.5 mol% Co(salophen)-HQ or 2.5 mol% Co(salophen) and 5 mol% quinone, at 30 °C under 1 atm of air. As shown in Figure 1, the use of Co(salophen)-HQ resulted in a faster reaction rate than that with Co(salophen) and quinone as separate ETMs. This result indicates that the intramolecular electron transfer between the hydroquinone unit and the oxidized metal-macrocycle of this bifunctional catalyst leads to a more efficient overall reaction under aerobic conditions.^[22]

After optimizing the reaction conditions, we continued to explore the scope of this transformation with various arylboronic acids (Scheme 3). First, when applying PhB(OH)₂ as the arylating partner, 77% isolated yield of cyclization product 3a was achieved. Importantly, this aerobic catalytic system appeared to be equally effective in comparison to the application of stoichiometric BQ as oxidant (76% yield in Scheme 3), therefore highlighting the advancement of this efficient and sustainable method. In addition to PhB(OH)2, arylboronic acids bearing an electron neutral group (3-Me), donating group (3-OMe) and withdrawing group (4-Acyl) all reacted well and produced the corresponding cyclic tetraenes (3b-3d) in good yields (61-77%, respectively). Various halogen-containing substrates (3-F, 3-Cl, and 4-Br) were well compatible with this methodology and gave the desired products (3e-3g) in 54-74% yields. 2-Naphthylboronic acid also reacted smoothly and a moderate yield (53% of 3h) could be achieved. Moreover, when 4-vinylphenylboronic acid was used as the substrate, the olefin bond remained intact and 52% yield of the desired product 3i was obtained. However, 2-thienylboronic acid, 4-pyridinylboronic acid and trans-2phenylvinylboronic acid did not give the corresponding tetraenes under the optimal reaction conditions.



Scheme 3. Substrate scope of different arylboronic acids 2. Reaction conditions: 1a (1.0 equiv.), $ArB(OH)_2$ 2 (1.5 equiv.), $Pd(OAc)_2$ (5 mol%), Co(salo-

phen)-HQ (10 mol%), H₂O (1.0 equiv.) in 0.1 M acetone, O₂ (1 atm) at room temperature (23 $^{\circ}$ C) for 48 h. Isolated yield. [a] Conditions employing 1.1 equiv. BQ instead of 10 mol% Co(salophen)-HQ under Ar. [b] Reaction time of 72 h.

We next investigated the reactivity of structurally diverse enallenynes 1 in this aerobic arylating carbocyclization (Scheme 4). Various substitutions on the allene moiety of 1, such as phenyl (3j), cyclopentyl (3k) and cyclohexyl (3l) groups were well tolerated, furnishing products in 61%-81% yields. Moreover, functional groups on the alkyne moiety, bearing -OAc (3m), -OBn (3n), and -CN (3o) were compatible with the aerobic conditions and gave the corresponding carbocycles in moderate to good yields (65%-79%). Under the optimized aerobic condition, cyclohexylacetylene-substituted enallenyne substrate 1p afforded 3p in only 36% yield. We attribute this diminished reactivity to the increased steric bulk of the alkyne moiety. In addition to an ester group, enallenynes containing butyl (3q), benzyl (3r), hydroxyl (3s), silyl (3t), sulfonamide (3u), and imide (3v) functionalities underwent this transformation smoothly highlighting the broad substrate scope of this protocol. Latestage oxidative reaction is a powerful approach for the streamlining and diversification of complex natural products and medicinal compounds. Here, an estrone-derived substrate, participated in this reaction efficiently to afford a useful yield (75%) of a functionalized complex molecule 3w.



WILEY-VCH

Furthermore, to demonstrate the necessity of the olefin group $(X = C=CH_2)$ in substrate 1, control experiments were performed with substrates lacking the olefin group (Table 2). At first, allenyne **1m**', without the olefin group in the β -position (X = CH₂) of the allene moiety, was applied under the standard reaction conditions. Notably, the six-membered carbocycle 3m' was not observed (entry 2 vs. entry 1). Allenynes containing heteroatom linkers such as O (1x) and NTs (1y) were evaluated in this aerobic arylative carbocyclization; however, neither of them was found to be effective (entries 3 and 4). In addition to an olefin group, we recently showed that a hydroxyl functionality, can trigger allene attack by weak coordination to the Pd(II) center, enabling efficient oxidative carbocyclizations of enallenols.^[23] We therefore examined the reactivity of an allenynol 1z, with a hydroxyl group at the β -position of the allene moiety. However, the desired oxidative carbocyclization did not take place in this case (entry 5). These experiments show that the initial step of allenic C(sp³)-H cleavage by Pd(II) requires the coordination of a pending olefin bond.^[18] A recent computational study by the Liu group supports that the assisting olefin group plays an indispensable role in the formation of the vinylpalladium intermediate.[24]

 Table 2. Investigation of different linkers in allenynes for Pd-catalyzed aerobic arylating carbocyclization.^[a]



[a] Reaction conditions: allene 1 (1.0 equiv.), PhB(OH)₂ 2a (1.5 equiv.), Pd(OAc)₂ (5 mol%), Co(salophen)-HQ (10 mol%), H₂O (1.0 equiv.) in 0.1 M acetone, O₂ (1 atm) at room temperature (23 $^{\circ}$ C) for 48 h.

On the basis of these experimental findings, the mechanism given in Scheme 5 is proposed for this oxidative arylating carbocyclization. Initially, the coordination of allene and olefin units to the Pd(II) center leads to a chelate palladium complex *Int-II.* This special coordination of the close-by olefin to Pd(II) is essential for triggering the allenic C(sp³)-H cleavage and generating a vinylpalladium intermediate *Int-III.*^[18] Next, instead of direct arylation to give **3**" or olefin insertion to form a cyclobutene complex *Int-B* (vide supra, Scheme 2), the envisioned ligand exchange of olefin^[25] by the distant alkyne moiety takes place to give *Int-IV*. Subsequent carbocyclization

of *Int-IV* by alkyne insertion gives a cyclic vinylpalladium *Int-V*. Transmetallation of *Int-V* with arylboronic acid **2**,^[26] followed by reductive elimination provides the target product **3** and Pd(0) species *Int-VI*, respectively. Finally, with the assistance of the cobalt hybrid catalyst as electron transfer mediator, aerobic oxidation of Pd(0) regenerates Pd(II) to close the catalytic cycle.^[27]



Scheme 5. Proposed mechanism.

In summary, we have developed an efficient Pd(II)-catalyzed oxidative carbocyclization of enallenynes using molecular oxygen as the oxidant. By applying a bifunctional catalyst (Co(salophen)-HQ) as an efficient electron transfer mediator, a wide range of enallenynes and arylboronic acids were transformed into the corresponding six-membered carbocycles in moderate to good yields. In view of the reaction efficiency and good selectivity, this protocol is expected to complement the current approach for oxidative functionalization in homogeneous catalysis and organic synthesis.

Acknowledgments

We are grateful for the financial support from European Research Council (ERC AdG 247014), Swedish Research Council (2016-03897, 2018-00830), and Carl Tryggers Foundation. We thank Dr. Zoltán Bacsik for the characterization of catalyst.

Keywords: palladium • aerobic oxidation • carbocyclization • enallenyne • electron transfer mediator •

- a) M.-C. Tang, Y. Zou, K. Watanabe, C. T. Walsh, Y. Tang, *Chem. Rev.* 2017, *117*, 5226–5333; b) X. Huang, J. T. Groves, *Chem. Rev.* 2018, *118*, 2491–2553.
- [2] J.-E. Bäckvall, Modern Oxidation Methods, Second Edition, Wiley-VCH Verlag & Co. KGaA, Weinheim, Germany, 2011.
- [3] S. S. Stahl, Science 2005, 309, 1824.
- [4] a) J. Reinhard, Angew. Chem. Int. Ed. 2009, 48, 9034–9037; Angew. Chem. 2009, 121, 9196–9199; b) J. A. Keith, P. M. Henry, Angew.

Chem. Int. Ed. 2009, 48, 9038–9049; Angew. Chem. 2009, 121, 9200–9212.

- [5] a) R. A. Sheldon, I. W. C. E. Arends, G.-J. ten Brink, A. Dijksman, Acc. Chem. Res. 2002, 35, 774–781; b) M. S. Sigman, D. R. Jensen, Acc. Chem. Res. 2006, 39, 221–229; c) E. J. Horn, B. R. Rosen, P. S. Baran, ACS Cent. Sci. 2016, 2, 302–308.
- [6] For recent reviews, see: a) R. I. McDonald, G. Liu, S. S. Stahl, *Chem. Rev.* 2011, *111*, 2981–3019; b) M. S. Sigman, E. W. Werner, *Acc. Chem. Res.* 2012, *45*, 874–884; c) W. Wu, H. Jiang, *Acc. Chem. Res.* 2012, *45*, 1736–1748; d) X.-F. Wu, X. Fang, L. Wu, R. Jackstell, H. Neumann, M. Beller, *Acc. Chem. Res.* 2014, *47*, 1041–1053; e) J. J. Dong, W. R. Browne, B. L. Feringa, *Angew. Chem. Int. Ed.* 2015, *54*, 734–744; *Angew. Chem.* 2015, *127*, 744–755; f) S. E. Mann, L. Benhamou, T. D. Sheppard, *Synthesis* 2015, *47*, 3079–3117; g) G. Yin, X. Mu, G. Liu, *Acc. Chem. Res.* 2016, *49*, 2413–2423; h) Z. Dong, Z. Ren, S. J. Thompson, Y. Xu, G. Dong, *Chem. Rev.* 2017, *117*, 9333–9403; i) D. Zhang, J. Liu, A. Córdova, W.-W. Liao, *ACS Catal.* 2017, *7*, 7051–7063; j) H. Sommer, F. Juliá-Hernández, R. Martin, I. Marek, *ACS Cent. Sci.* 2018, *4*, 153–165.
- [7] For recent reviews, see: a) C. Liu, J. Yuan, M. Gao, S. Tang, W. Li, R. Shi, A. Lei, *Chem. Rev.* 2015, *115*, 12138–12204; b) J. F. Hartwig, M. A. Larsen, *ACS Cent. Sci.* 2016, *2*, 281–292; c) A. V. Iosub, S. S. Stahl, *ACS Catal.* 2016, *6*, 8201–8213; d) H. M. L. Davies, D. Morton, *ACS Cent. Sci.* 2017, *3*, 936–943; e) J. He, M. Wasa, K. S. L. Chan, Q. Shao, J.-Q. Yu, *Chem. Rev.* 2017, *117*, 8754–8786; f) Y.-F. Liang, N. Jiao, *Acc. Chem. Res.* 2017, *50*, 1640–1653; g) Y. Yang, J. Lan, J. You, *Chem. Rev.* 2017, *117*, 8787–8863; h) H. Sterckx, B. Morel, B. U. W. Maes, *Angew. Chem. Int. Ed.* 2018, *57*, 10.1002/anie.201804946.
- [8] a) A. N. Campbell, S. S. Stahl, Acc. Chem. Res. 2012, 45, 851–863; b)
 L. Jin, A. Lei, Sci. China Chem. 2012, 55, 2027–2035; c) Z. Shi, C.
 Zhang, C. Tang, N. Jiao, Chem. Soc. Rev. 2012, 41, 3381–3430.
- [9] D. Wang, A. B. Weinstein, P. B. White, S. S. Stahl, Chem. Rev. 2018, 118, 2636–2679.
- a) J. Piera, J.-E. Bäckvall, Angew. Chem. Int. Ed. 2008, 47, 3506–3523; Angew. Chem. 2008, 120, 3558–3576; b) A. Vasseur, J. Muzart, J. Le
 Bras, Eur. J. Org. Chem. 2015, 2015, 4053–4069; c) A. E. Wendlandt,
 S. S. Stahl, Angew. Chem. Int. Ed. 2015, 54, 14638–14658; Angew. Chem. 2015, 127, 14848–14868.
- a) J.-E. Bäckvall, A. K. Awasthi, Z. D. Renko, J. Am. Chem. Soc. 1987, 109, 4750–4752; b) J.-E. Bäckvall, R. B. Hopkins, H. Grennberg, M. Mader, A. K. Awasthi, J. Am. Chem. Soc. 1990, 112, 5160–5166.
- [12] For recent examples, see: a) J. Piera, K. Naerhi, J.-E. Bäckvall, Angew. Chem. Int. Ed. 2006, 45, 6914–6917; Angew. Chem. 2006, 118, 7068– 7071; b) J. Piera, A. Persson, X. Caldentey, J.-E. Bäckvall, J. Am. Chem. Soc. 2007, 129, 14120–14121; c) B. Morandi, Z. K. Wickens, R. H. Grubbs, Angew. Chem. Int. Ed. 2013, 52, 2944–2948; Angew. Chem. 2013, 125, 3016–3020; d) C. M. R. Volla, J.-E. Bäckvall, Angew. Chem. Int. Ed. 2013, 52, 14209–14213; Angew. Chem. 2013, 125, 14459– 14463; e) C. C. Pattillo, I. I. Strambeanu, P. Calleja, N. A. Vermeulen, T. Mizuno, M. C. White, J. Am. Chem. Soc. 2016, 138, 1265–1272; f) L. Ta, A. Axelsson, H. Sunden, Green Chem. 2016, 18, 686–690.
- a) H. Grennberg, S. Faizon, J.-E. Bäckvall, Angew. Chem., Int. Ed. Engl.
 1993, 32, 263–264; Angew. Chem. 1993, 105, 269–271; b) B. W. Purse,

L.-H. Tran, J. Piera, B. Åkermark, J.-E. Bäckvall, *Chem. Eur. J.* **2008**, *14*, 7500–7503; c) E. V. Johnston, E. A. Karlsson, S. A. Lindberg, B. Åkermark, J.-E. Bäckvall, *Chem. Eur. J.* **2009**, *15*, 6799–6801; d) E. V. Johnston, E. A. Karlsson, L.-H. Tran, B. Aakermark, J.-E. Bäckvall, *Eur. J. Org. Chem.* **2009**, 3973–3976.

- [14] a) N. Krause, A. S. K. Hashmi, *Modern Allene Chemistry*, Wiley-VCH Verlag & Co. KGaA, Weinheim, Germany, 2008; b) C. Aubert, L. Fensterbank, P. Garcia, M. Malacria, A. Simonneau, *Chem. Rev.* 2011, *111*, 1954–1993; c) F. Inagaki, S. Kitagaki, C. Mukai, *Synlett* 2011, *2011*, 594–614; d) N. Krause, C. Winter, *Chem. Rev.* 2011, *111*, 1994–2009; e) F. López, J. L. Mascareñas, *Chem. Eur. J.* 2011, *17*, 418–428; f) P. Rivera-Fuentes, F. Diederich, *Angew. Chem. Int. Ed.* 2012, *51*, 2818–2828; *Angew. Chem.* 2012, *124*, 2872–2882; g) S. Yu, S. Ma, *Angew. Chem. Int. Ed.* 2012, *51*, 3074–3112; *Angew. Chem.* 2012, *124*, 3128–3167; h) J. Ye, S. Ma, *Acc. Chem. Res.* 2014, *47*, 989–1000.
- [15] B. Yang, Y. Qiu, J.-E. Bäckvall, Acc. Chem. Res. 2018, 51, 1520–1531.
- [16] a) N. Saito, T. Ichimaru, Y. Sato, *Chem. Asian J.* 2012, *7*, 1521–1523;
 b) Y. Ohta, S. Yasuda, Y. Yokogawa, K. Kurokawa, C. Mukai, *Angew. Chem. Int. Ed.* 2014, *54*, 1240–1244; *Angew. Chem.* 2015, *127*, 1256–1260; c) C. Raviola, S. Protti, D. Ravelli, M. Fagnoni, *Chem. Soc. Rev.* 2016, *45*, 4364–4390; d) D. Cassú, T. Parella, M. Solà, A. Pla-Quintana, A. Roglans, *Chem. Eur. J.* 2017, *23*, 14889–14899.
- [17] Y. Qiu, B. Yang, C. Zhu, J.-E. Bäckvall, Angew. Chem. Int. Ed. 2016, 55, 6520–6524; Angew. Chem. 2016, 128, 6630–6634.
- [18] C. Zhu, B. Yang, T. Jiang, J.-E. Bäckvall, Angew. Chem. Int. Ed. 2015, 54, 9066–9069; Angew. Chem. 2015, 127, 9194–9197.
- [19] There is an equilibrium between PhB(OH)₂ and its anhydride form (PhBO)₃. We speculate that the addition of water can promote the equilibrium toward the formation of PhB(OH)₂. Please see the Supporting Information for details.
- [20] The side product **3a**' and **3a**'' in Scheme 2 were not obtained. For a discussion on the selectivity, please see the Supporting Information.
- [21] The isomer of arene from aromatization of tetraene 3a was not observed.
- [22] A proposed catalytic cycle for reoxidation of Pd(0) by Co(salophen)-HQ is given in the Supporting Information. For a related mechanistic study, see: C. W. Anson, S. Ghosh, S. Hammes-Schiffer, S. S. Stahl, *J. Am. Chem. Soc.* 2016, *138*, 4186–4193.
- [23] D. Posevins, Y. Qiu, J.-E. Bäckvall, J. Am. Chem. Soc. 2018, 140, 3210–3214.
- [24] L. Han, T. Liu, Org. Biomol. Chem. 2017, 15, 5055–5061.
- [25] a) C. Zhu, B. Yang, Y. Qiu, J.-E. Bäckvall, *Angew. Chem. Int. Ed.* 2016, 55, 14405–14408; *Angew. Chem.* 2016, 128, 14617–14620; b) Y. Qiu, B. Yang, C. Zhu, J.-E. Bäckvall, *Chem. Sci.* 2017, 8, 616–620.
- [26] In the absence of ArB(OH)₂, no product was obtained. The *Int-V* could not be detected because a fast trapping of *Int-V* by ArB(OH)₂ occurs and generates the corresponding cyclic tetraene.
- [27] In addition to facilitate electron transfer from the reduced palladium catalyst to O_2 , we speculate the quinone moiety of the oxidized Co(salophen)-HQ can act as a ligand that coordinates to the Pd(II) intermediate during the catalysis. Quinone coordination to Pd(II) could withdraw electron density from the Pd(II) center, therefore promoting the subsequent reductive elimination step.

lanusc

WILEY-VCH

COMMUNICATION

Entry for the Table of Contents (Please choose one layout)

COMMUNICATION



Co catalyst as co-catalyst: We report the first example of a Pd(II)-catalyzed oxidative carbocyclization of the rational designed enallenynes using molecular oxygen. By applying a bifunctional catalyst (Co(salophen)-HQ) as an efficient electron transfer mediator, enallenynes are effectively arylated to the corresponding six-membered carbocycles in good yields.

Jie Liu, Alexander Ricke, Bin Yang,* and Jan-E. Bäckvall*

Page No. – Page No.

Efficient Palladium-Catalyzed Aerobic Arylative Carbocyclization of Enallenynes