

Journal of Carbohydrate Chemistry

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/lcar20>

Synthesis of (Methyl 3-O-Benzyl-4,6-O-benzylidene-2-deoxy- α -D-altropyranosid-2-yl)thiophene Derivatives as Precursors of New Iso-C-nucleoside Analogues

Lidcay Herrera ^a, Holger Feist ^a, José Quincoces ^b, Manfred Michalik ^c & Klaus Peseke ^a

^a Fachbereich Chemie, Universität Rostock, D-18051, Rostock, Germany

^b Universidade Bandeirante de São Paulo, Rua Maria Candida, 1813, Vila Guilherme, São Paulo, CEP: 02071-013, Brazil

^c Institut für Organische Katalyseforschung, Buchbinderstraße 5-6, D-18055, Rostock, Germany

Published online: 28 Oct 2011.

To cite this article: Lidcay Herrera, Holger Feist, José Quincoces, Manfred Michalik & Klaus Peseke (2003) Synthesis of (Methyl 3-O-Benzyl-4,6-O-benzylidene-2-deoxy- α -D-altropyranosid-2-yl)thiophene Derivatives as Precursors of New Iso-C-nucleoside Analogues, Journal of Carbohydrate Chemistry, 22:3-4, 171-178, DOI: [10.1081/CAR-120021697](https://doi.org/10.1081/CAR-120021697)

To link to this article: <http://dx.doi.org/10.1081/CAR-120021697>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

Synthesis of (Methyl 3-*O*-Benzyl-4,6-*O*-benzylidene-2-deoxy- α -D-altropyranosid-2-yl)thiophene Derivatives as Precursors of New Iso-*C*-nucleoside Analogues

Lidcay Herrera,¹ Holger Feist,¹ José Quincoces,²
Manfred Michalik,³ and Klaus Peseke^{1,*}

¹Fachbereich Chemie, Universität Rostock, Rostock, Germany

²Universidade Bandeirante de São Paulo, Vila Guilherme, São Paulo, Brazil

³Institut für Organische Katalyseforschung, Rostock, Germany

ABSTRACT

Treatment of 2-(methyl 3-*O*-benzyl-4,6-*O*-benzylidene-2-deoxy- α -D-altropyranosid-2-yl)ethanal (**3**) with malononitrile in the presence of aluminium oxide provided 2-cyano-4-(methyl 3-*O*-benzyl-4,6-*O*-benzylidene-2-deoxy- α -D-altropyranosid-2-yl)crotononitrile (**4**). Starting from **4**, cyclization with sulphur and triethylamine yielded 2-amino-5-(methyl 3-*O*-benzyl-4,6-*O*-benzylidene-2-deoxy- α -D-altropyranosid-2-yl)thiophene-3-carbonitrile (**5**). Further cyclization could be achieved with triethyl orthoformate/ammonia to furnish 4-amino-6-(methyl 3-*O*-benzyl-4,6-*O*-benzylidene-2-deoxy- α -D-altropyranosid-2-yl)thieno[2.3-*d*]pyrimidine (**8**).

Key Words: Nucleoside analogues; Branched-chain sugars; Malononitrile; Thiophene; Thieno[2.3-*d*]pyrimidine.

*Correspondence: Klaus Peseke, Fachbereich Chemie, Universität Rostock D-18051 Rostock, Germany; Fax: 49-381-498-6412; E-mail: klaus.peseke@chemie.uni-rostock.de.



INTRODUCTION

The synthesis of *C*-nucleoside analogues in the last years has been carried out in a search for compounds having anticancer, antibacterial and antiviral activities.^[1–4] Furthermore, iso-*C*-nucleosides in which the nucleobase is linked by a carbon-carbon bond to the sugar moiety at a carbon other than *C*-1 are a special type of *C*-nucleosides. These substances may have an increased chemical and enzymatic stability under physiological conditions.^[5]


Several syntheses of iso-*C*-nucleosides have been reported.^[4–6] Recently we described a method for the preparation of *C*-nucleosides on the basis of (α -D-glycopyranosyl) thiophene and thienopyrimidine derivatives^[7] due to the wide spectrum of interesting biological properties of thiophene derivatives such as serine protease inhibitions,^[8] dihydrofolate reductase inhibitions,^[9] analgesic,^[10,11] and anti-inflammatory^[11] properties, among others. In the present paper we describe the synthesis of the corresponding iso-*C*-nucleosides, the (methyl 3-*O*-benzyl-4,6-*O*-benzylidene-2-deoxy- α -D-altropyranosid-2-yl)thiophene and (methyl 3-*O*-benzyl-4,6-*O*-benzylidene-2-deoxy- α -D-altropyranosid-2-yl)thieno[2.3-*d*]pyrimidine.

RESULTS AND DISCUSSION

Methyl 2,3-anhydro-4,6-benzylidene- α -D-allopyranoside **1** was reacted in two steps to furnish the known alkene derivative **2**.^[12,13] Oxidation of **2** with osmium tetroxide and sodium periodate^[14] afforded in two hours 2-(methyl 3-*O*-benzyl-4,6-*O*-benzylidene-2-deoxy- α -D-altropyranosid-2-yl)ethanal **3** in 66% yield. The signals in the ¹H and ¹³C NMR spectra found at δ = 9.66 and 199.0, respectively, are typical of the aldehyde group in structure **3**.

Starting from **3**, the Knoevenagel reaction with excess malononitrile could be performed to yield the crotononitrile derivative **4** (Scheme 1). The reaction was catalyzed by aluminium oxide,^[15,16] permitting complete transformation of the starting material in only one hour. After isolation of the product by column chromatography, the spectroscopic data of the product clearly verified the formation of compound **4**. In the IR spectrum both cyano groups appeared at 2242 cm⁻¹ and the mass spectrum gave the corresponding molecular peak.

The degree of acidity of the methylene group at the neighbouring position to the dicyanomethylene moiety allowed the preparation of 2-amino-5-(methyl 3-*O*-benzyl-4,6-*O*-benzylidene-2-deoxy- α -D-altropyranosid-2-yl)thiophene-3-carbonitrile (**5**) by treatment of **4** with elementary sulphur and triethylamine.^[17] In the original procedure for the synthesis of such thiophenes, ethanol was used as solvent.^[18] However, we carried out the reaction in *N,N*-dimethylformamide and isolated the thiophene-3-carbonitrile **5** as light yellow solid in 82% yield. As expected, in the IR spectrum of **5** only one CN band was found at a lower wave number 2204 cm⁻¹ due to the conjugation of the cyano group with the thiophene ring and the amino group. In the ¹³C NMR spectrum of **5** only one CN signal at δ = 115.2 was present and in the ¹H NMR spectrum the signals of H-4 of the thiophene ring and amino group appeared at δ = 6.51 and 4.71, respectively.




Scheme 1.

For further cyclization the enamionitrile unit located in the thiophene ring of **5** could be used.^[19] Treatment of thiophene **5** with triethyl orthoformate under reflux afforded a crude syrupy formimidic acid ester **6** which furnished, by reaction with a saturated ethanolic ammonia solution at room temperature, the 2-[aminomethylenamino]thiophene-3-carbonitrile **7** in 41% yield. On the other hand, **6** reacted with a saturated ethanolic ammonia solution under reflux gave the corresponding thieno[2.3-*d*]pyrimidine-4-amine **8** in 64% yield (Scheme 2).

In the IR and ^{13}C NMR spectra of compound **7** the expected CN peaks were displayed and in the ^1H NMR spectrum a broad singlet at $\delta = 7.83$ could be assigned to the aminomethylenamino group of the thiophene ring. The absence of CN peaks in the IR and in the ^{13}C NMR spectra of the (α -D-altropyranosid-2-yl)thieno[2.3-*d*]pyrimidine **8** proved the successful ring closure. Furthermore, in the ^1H NMR spectrum a defined singlet appeared at $\delta = 8.42$ belonging to H-2 of the thieno[2.3-*d*]pyrimidine ring.





Scheme 2.

EXPERIMENTAL

General procedures. Melting points were determined with a Boëtius melting point apparatus and are corrected. Optical rotations were measured with a Polar L μ P (IBZ Meßtechnik) polarimeter. IR spectra were recorded with a Nicolet 205 FT-IR spectrometer. ^1H and ^{13}C NMR spectra were recorded with a Bruker AC 250 (250.1 MHz and 62.9 MHz, respectively) and a Bruker ARX 300 (300.1 MHz and 75.5 MHz, respectively). The calibration of spectra was carried out by means of solvent peaks (CDCl_3 : δ ^1H 7.25; δ ^{13}C 77.0). The ^{13}C NMR signals were assigned by DEPT and/or two-dimensional ^1H , ^{13}C correlation spectra. Mass spectra were obtained with an AMD 402/3 spectrometer (AMD Intectra GmbH). Elemental analyses were performed with a Leco CHNS-932. Column chromatography was carried out on silica gel 60 (230–400 μm , Merck). Thin-layer chromatography (TLC) was performed on silica gel 60 GF $_{254}$ foils (Merck) with detection by UV-light and by charring with sulphuric acid. Solvents and liquid reagents were purified and dried according to recommended procedures.



2-(Methyl 3-*O*-benzyl-4,6-*O*-benzylidene-2-deoxy- α -D-altropyranosid-2-yl)ethanal (3). To methyl 3-*O*-benzyl-4,6-*O*-benzylidene-2-deoxy-2-*C*-(prop-2-enyl)- α -D-altropyranoside (2) (1.98 g, 5 mmol) dissolved in dioxane/water 10:1 (50 mL), was added an OsO₄-solution (0.4 mg in dioxane, 0.4 mL), and the resulting reaction mixture was stirred at room temperature for 45 min. Then NaIO₄ (2.14 g), as a saturated solution in water was added, the mixture stirred for 2 h, filtered, and the solvent completely removed in vacuo. Compound 3 was obtained as a white solid after column chromatography (toluene/ethyl acetate 3:1): yield 1.31 g (66%); mp 95–97°C; $[\alpha]_D^{21} +45.7$ (*c* 1.0, CHCl₃); *R*_f 0.35 (toluene/ethyl acetate 3:1). IR (KBr): 1717 cm⁻¹ (CO). ¹H NMR (250.1 MHz, CDCl₃), δ 9.66 (t, 1H, ³*J*_{1,2} = 1.2 Hz, H-1); 7.47–7.15 (m, 10H, ArH); 5.54 (s, 1H, CH-Ph); 4.82, 4.78 (q (AB), 2H, ²*J*_{A,B} = 13.1 Hz, CH₂-Ph); 4.44 (m, 1H, H-5'); 4.41 (d, 1H, ³*J*_{1',2'} = 0.5 Hz, H-1'); 4.31 (dd, 1H, ²*J*_{6'a,6'b} = 10.3 Hz, ³*J*_{5',6'a} = 5.4 Hz, H-6'a); 3.75–3.64 (m, 3H, H-3', H-4', H-6'b); 3.41 (s, 3H, OMe); 2.90 (m, 1H, H-2'); 2.66 (ddd, 1H, ²*J*_{2a,2b} = 18.2, Hz, ³*J*_{2a,2'} = 8.0 Hz, ³*J*_{1,2a} = 1.2 Hz, H-2a); 2.47 (ddd, 1H, ²*J*_{2a,2b} = 18.2, Hz, ³*J*_{2b,2'} = 6.3 Hz, ³*J*_{1,2b} = 1.2 Hz, H-2b). ¹³C NMR (62.9 MHz, CDCl₃), δ 199.0 (C-1); 138.7, 137.5 (2 × *i*-Ph); 129.0 (*p*-Ph), 128.2, 128.1, 127.5, 126.2 (*o*-,*m*-Ph); 127.2 (*p*-Ph); 102.2 (CH-Ph); 101.4 (C-1'); 77.2 (C-4'); 74.4 (C-3'); 72.1 (CH₂-Ph); 69.4 (C-6'); 58.2 (C-5'); 55.7 (OMe); 44.7 (C-2); 37.5 (C-2'). MS (EI), *m/z*: 398 [M]⁺.

Anal. Calcd for C₂₃H₂₆O₆ (398.45): C, 69.33; H, 6.58. Found: C, 69.18; H, 6.70.

2-Cyano-4-(methyl 3-*O*-benzyl-4,6-*O*-benzylidene-2-deoxy- α -D-altropyranosid-2-yl)crotononitrile (4). To a solution of 3 (0.398 g, 1 mmol) in dried toluene (30 mL), were added malononitrile (0.165 g, 2.5 mmol) and Al₂O₃ (0.1 g), and the mixture was stirred under reflux for 1 h. After filtration the solvent was evaporated, the residue was purified by column chromatography (toluene/ethyl acetate 3:1) and recrystallized from ethanol to give 4 as white crystals: yield 0.388 g (87%); mp 148–150°C; $[\alpha]_D^{21} -56.7$ (*c* 1.0, CHCl₃); *R*_f 0.45 (toluene/ethyl acetate 3:1). IR (KBr): 2242 cm⁻¹ (CN). ¹H NMR (250.1 MHz, CDCl₃), δ 7.52–7.33 (m, 10H, ArH); 6.86 (t, 1H, ³*J*_{3,4} = 7.8 Hz, H-3); 5.57 (s, 1H, CH-Ph); 4.89, 4.67 (2d, 2H, ²*J*_{A,B} = 12.5 Hz, CH₂-Ph); 4.46–4.29 (m, 3H, H-1', H-5', H-6'a); 3.78–3.69 (m, 2H, H-4', H-6'b); 3.58 (dd, 1H, ³*J*_{3',4'} = 3.0 Hz, ³*J*_{2',3'} = 2.0 Hz, H-3'); 3.41 (s, 3H, OMe); 2.63 ('t', 2H, ³*J*_{3,4} = ³*J*_{2',4} = 7.8 Hz, H-4); 2.31 (dt, 1H, ³*J*_{2',4} = 7.8 Hz, ³*J*_{2',3'} = 2.0 Hz, H-2'). ¹³C NMR (62.9 MHz, CDCl₃), δ 165.7 (C-3); 138.0, 137.4 (2 × *i*-Ph); 129.1 (*p*-Ph), 128.4, 128.3, 128.2, 126.2 (*o*-,*m*-Ph); 128.0 (*p*-Ph); 111.3, 110.1 (2 × CN); 102.3 (CH-Ph); 100.8 (C-1'); 91.7 (C-2); 76.9 (C-4'); 73.1 (CH₂-Ph); 72.6 (C-3'); 69.3 (C-6'); 58.3 (C-5'); 55.7 (OMe); 43.6 (C-2'); 33.6 (C-4). MS (CI), *m/z*: 447 [M+1]⁺.

Anal. Calcd for C₂₆H₂₆N₂O₅ (446.50): C, 69.94; H, 5.87; N, 6.27. Found: C, 69.81; H, 5.64; N, 6.39.

2-Amino-5-(methyl 3-*O*-benzyl-4,6-*O*-benzylidene-2-deoxy- α -D-altropyranosid-2-yl)thiophene-3-carbonitrile (5). Compound 4 (0.446 g, 1 mmol) was dissolved in dry DMF (5 mL). To this solution sulphur (0.048 g, 1.5 mmol) and triethylamine (0.21 mL) were added, and the entire mixture was stirred at room temperature for 2 h. Water (30 mL) was added, the mixture was extracted with CH₂Cl₂ (3 × 50 mL), and the organic phase was washed with water (2 × 50 mL), dried (Na₂SO₄) and concentrated under reduced pressure. The residue was purified by column chromato-



graphy (toluene/ethyl acetate 3:1) to give **5** as a light yellow solid: yield 0.392 g (82%); mp 158–160°C; $[\alpha]_D^{21} -65.3$ (*c* 1.0, CHCl₃); *R*_f 0.40 (toluene/ethyl acetate 3:1). IR (KBr): 2204 cm⁻¹ (CN). ¹H NMR (250.1 MHz, CDCl₃), δ 7.50–7.20 (m, 10H, ArH); 6.51 (d, 1H, ⁴*J*_{2',4} = 0.9 Hz, H-4); 5.52 (s, 1H, CH-Ph); 4.89 (d, 1H, ²*J*_{A,B} = 12.5 Hz, 1 × CH₂-Ph); 4.82 (br s, 1H, ³*J*_{1',2'} = 0.6 Hz, H-1'); 4.74 (d, 1H, ²*J*_{A,B} = 12.5 Hz, 1 × CH₂-Ph); 4.71 (br, 2H, NH₂); 4.48 (m, 1H, H-5'); 4.36 (dd, 1H, ²*J*_{6'a,6'b} = 10.1 Hz, ³*J*_{5',6'a} = 5.5 Hz, H-6'a); 3.92 (dd, 1H, ³*J*_{4',5'} = 9.7 Hz, ³*J*_{3',4'} = 3.0 Hz, H-4'); 3.84 (br “t”, 1H, ³*J*_{3',4'} = 3.0 Hz, ³*J*_{2',3'} = 2.5 Hz, H-3'); 3.77 (“t”, 1H, ²*J*_{6'a,6'b} = ³*J*_{5',6'b} = 10.3 Hz, H-6'b); 3.43 (s, 3H, OMe); 3.39 (m, 1H, H-2'). ¹³C NMR (62.9 MHz, CDCl₃), δ 161.3 (C-2); 138.4, 137.5 (2 × *i*-Ph); 129.0 (*p*-Ph); 128.3, 128.2, 127.8, 126.2, (*o*-, *m*-Ph); 127.6 (*p*-Ph); 126.0 (C-5); 123.5 (C-4); 115.2 (CN); 102.3 (CH-Ph); 100.8 (C-1'); 87.3 (C-3); 76.4 (C-4'); 75.9 (C-3'); 72.8 (CH₂-Ph); 69.4 (C-6'); 58.5 (C-5'); 55.7 (OMe); 45.5 (C-2'). MS (CI), *m/z*: 479 [M+1]⁺.

Anal. Calcd for C₂₆H₂₆N₂O₅S (478.56): C, 65.26; H, 5.48; N, 5.85; S, 6.70. Found: C, 65.56; H, 5.69; N, 5.76; S, 6.56.

5-(Methyl 3-*O*-benzyl-4,6-*O*-benzylidene-2-deoxy- α -D-altropyranosid-2-yl)-2-[aminomethylenamino]thiophene-3-carbonitrile (7). Compound **5** (0.478 g, 1 mmol) was dissolved in triethyl orthoformate (10 mL) and the reaction mixture was refluxed for 1 h. The solvent was then evaporated under reduced pressure and the residue dissolved in a saturated solution (10 mL) of NH₃ in ethanol. The mixture was then stirred for 1 h, and the solvent was evaporated. The residue was purified by column chromatography (toluene/ethyl acetate 1:1) to give **7** as a white solid: yield 0.207 g (41%); mp 113–115°C; $[\alpha]_D^{21} -75.8$ (*c* 0.23, CHCl₃); *R*_f 0.41 (toluene/ethyl acetate 1:1). IR (KBr): 2215 cm⁻¹ (CN). ¹H NMR (250.1 MHz, CDCl₃), δ 7.83 (br, 1H, CH = N); 7.50–7.15 (m, 10H, ArH); 6.71 (d, 1H, ⁴*J*_{2',4} = 0.9 Hz, H-4); 5.53 (s, 1H, CH-Ph); 5.46 (br, NH₂); 4.89 (d, 1H, ²*J*_{A,B} = 12.8 Hz, 1 × CH₂-Ph); 4.86 (s, 1H, H-1'); 4.76 (d, 1H, ²*J*_{A,B} = 12.8 Hz, 1 × CH₂-Ph); 4.50 (m, 1H, H-5'); 4.37 (dd, 1H, ²*J*_{6'a,6'b} = 10.2 Hz, ³*J*_{5',6'a} = 5.5 Hz, H-6'a); 3.93 (dd, 1H, ³*J*_{4',5'} = 9.5 Hz, ³*J*_{3',4'} = 3.0 Hz, H-4'); 3.88 (br “t”, 1H, ³*J*_{3',4'} = 3.0 Hz, ³*J*_{2',3'} = 2.5 Hz, H-3'); 3.79 (“t”, 1H, ²*J*_{6'a,6'b} = ³*J*_{5',6'b} = 10.2 Hz, H-6'b); 3.48 (m, 1H, H-2'); 3.44 (s, 3H, OMe). ¹³C NMR (62.9 MHz, CDCl₃), δ 164.8 (C-2); 153.8 (CH=N); 138.4, 137.5 (2 × *i*-Ph); 131.1 (C-5); 129.0 (*p*-Ph); 128.3, 128.2, 127.7, 126.2, (*o*-, *m*-Ph); 127.6 (*p*-Ph); 124.4 (C-4); 116.1 (CN); 102.3 (CH-Ph); 100.8 (C-1'); 96.9 (C-3); 76.3 (C-4'); 76.0 (C-3'); 72.8 (CH₂-Ph); 69.4 (C-6'); 58.5 (C-5'); 55.7 (OMe); 46.1 (C-2'). MS (CI), *m/z*: 506 [M+1]⁺.

Anal. Calcd for C₂₇H₂₇N₃O₅S (505.58): C, 64.14; H, 5.38; N, 8.31; S, 6.34. Found: C, 64.28; H, 5.45; N, 8.38; S, 6.30.

4-Amino-6-(methyl 3-*O*-benzyl-4,6-*O*-benzylidene-2-deoxy- α -D-altropyranosid-2-yl)thieno[2.3-*d*]pyrimidine (8). Compound **5** (0.478 g, 1 mmol) was dissolved in triethyl orthoformate (10 mL) and the reaction mixture was refluxed for 1 h. The solvent was evaporated under reduced pressure, and the residue was dissolved in a saturated solution (10 mL) of NH₃ in ethanol. The mixture was stirred for 1 h and then heated until the product **7** disappeared (TLC control). The solvent was removed and the residue purified by column chromatography (ethyl acetate) to give **8** as a white solid: yield 0.323 g (64%); mp 100–102°C; $[\alpha]_D^{21} -85.3$ (*c* 1.0, CHCl₃); *R*_f 0.33 (ethyl acetate). IR (KBr): 3230, 3196 cm⁻¹ (NH₂). ¹H NMR (250.1 MHz, CDCl₃), δ 8.42



(s, 1H, H-2); 7.48–7.27 (m, 10H, ArH); 6.97 (s, 1H, H-5); 5.67 (br d, 2H, NH₂); 5.49 (s, 1H, CH-Ph); 5.01 (s, 1H, H-1'); 4.93, 4.79 (q (AB), 2H, $^2J_{A,B}$ = 12.8 Hz, CH₂-Ph); 4.55 (m, 1H, H-5'); 4.38 (dd, 1H, $^2J_{6'a,6'b}$ = 10.3 Hz, $^3J_{5',6'a}$ = 5.5 Hz, H-6'a), 4.02 (br "t", 1H, H-3'); 3.95 (dd, 1H, $^3J_{4',5'}$ = 9.5 Hz, $^3J_{3',4'}$ = 3.1 Hz, H-4'); 3.78 (t, 1H, $^2J_{6'a,6'b}$ = $^3J_{5',6'b}$ = 10.3 Hz, H-6b'); 3.69 (br d, 1H, $^3J_{2',3'}$ = 2.5 Hz, H-2'); 3.47 (s, 3H, OMe). ¹³C NMR (62.9 MHz, CDCl₃), δ 166.9, 157.2 (C-4, C-7a); 153.7 (C-2); 139.8, 116.9 (C-4a, C-6); 138.2, 137.4 (2 $\times$ *i*-Ph); 129.0 (*p*-Ph); 128.3, 128.2, 127.9, 126.1 (*o*-, *m*-Ph); 127.6 (*p*-Ph); 115.7 (C-5); 102.2 (CH-Ph); 100.7 (C-1'); 76.3 (C-4'); 76.0 (C-3'); 72.9 (CH₂-Ph); 69.4 (C-6'); 58.6 (C-5'); 55.8 (OMe); 46.4 (C-2'). MS (CI), m/z : 506 [M+1]⁺.

Anal. Calcd for C₂₇H₂₇N₃O₅S (505.58): C, 64.14; H, 5.38; N, 8.31; S, 6.34. Found: C, 63.92; H, 5.54; N, 8.15; S, 6.63.

ACKNOWLEDGMENTS

We are grateful to the Deutsche Forschungsgemeinschaft and the Fonds der Chemischen Industrie for financial support. J. Q. is grateful to the Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP, Brasil) and the Deutscher Akademischer Austauschdienst for financial support. L. H. would like to thank the Deutscher Akademischer Austauschdienst for a scholarship.

REFERENCES

1. Simons, C. *C-nucleosides*. In *Nucleoside Mimetics: Their Chemistry and Biological Properties*; Simons, C., Ed.; Gordon and Breach Science Publishers: Australia, 2001; 117–135.
2. Buchanan, J.G.; Wightman, R.H. The *C*-nucleoside antibiotics. *Prog. Chem. Org. Nat. Prod.* **1984**, *44*, 243–299.
3. Humber, D.C.; Mulholland, K.R.; Stoodley, R.J. *C*-nucleosides. Part 1. Preparation of tiazofurin and *N*-substituted tiazofurins from benzyl(2',3',5'-tri-*O*-benzoyl- β -D-ribofuranosyl)penicillinate. *J. Chem. Soc., Perkin Trans. 1* **1990**, 283–292.
4. Yu, H.W.; Zhang, H.Y.; Yang, Z.J.; Min, J.M.; Ma, L.T.; Zhang, L.H. Studies on the syntheses and biological activities of isonucleosides. *Pure Appl. Chem.* **1998**, *70*, 435–438.
5. Zhang, M.; Zhang, H.; Yang, Z.; Ma, L.; Min, J.; Zhang, L. Synthesis of 3-*C*-(Methyl β -D-xylofuranosid-3-yl)-5-phenyl-1,2,4-oxadiazole. *Carbohydr. Res.* **1999**, *318*, 157–161.
6. Casiraghi, G.; Cornia, M.; Rassu, G.; Sante, C.D.; Spanu, P. Synthesis and transformation of pyrrole *C*-glycoconjugates. *Tetrahedron* **1992**, *48*, 5619–5628.
7. García, I.; Feist, H.; Cao, R.; Michalik, M.; Peseke, K. Synthesis of (2,3,4,6-Tetra-*O*-acetyl- α -D-glycopyranosyl)thiophene derivatives as new *C*-nucleoside analogues. *J. Carbohydr. Chem.* **2001**, *20*, 681–687.
8. Pinto, I.L.; Jarvest, R.L.; Serafinowska, H.T. The synthesis of 5-alkoxy and 5-amino substituted thiophenes. *Tetrahedron Lett.* **2000**, *41*, 1597–1600.
9. Rosowsky, A.; Mota, C.E.; Wright, J.E.; Freisheim, J.H.; Heusner, J.J.;



- McCormack, J.J.; Queener, S.F. 2,4-diaminothieno[2,3-d]pyrimidine analogues of trimetrexate and piritrexim as potential inhibitors of *Pneumocystis carinii* and *Toxoplasma gondii* dihydrofolate reductase. *J. Med. Chem.* **1993**, *36*, 3103–3112.
10. Huybrechts, L.; Buffel, D.; Freyne, E.; Hoornaert, G. The synthesis of 5-carboxamido-4-hydroxy-3-(β -D-ribofuranosyl)-thiophene derivatives, analogues of pyrazofurin. *Tetrahedron* **1984**, *40*, 2479–2485.
11. Dave, C.G.; Shah, P.R.; Shah, A.B.; Dave, K.C.; Patel, V.J. Synthesis and biological activity of pyrido[3',2':4,5]thieno[3,2-d]pyrimidines. *J. Indian Chem. Soc.* **1989**, *66*, 48–50.
12. Hughes, A.B.; Stick, R.V.; Tilbrook, D.M. The synthesis of some carbohydrate-derived precursors to tylonolide, the aglycon of the antibiotic tylosin. *Aust. J. Chem.* **1990**, *43*, 1681–1695.
13. Wood, A.J.; Holt, D.J.; Domínguez, M.; Jenkins, P.R. The stereoselective preparation of an enantiomerically pure cyclopentane using intramolecular aldol cyclopentaannulation of a glucose derivative. *J. Org. Chem.* **1998**, *63*, 8522–8529.
14. Tronchet, J.M.J.; Zsély, M.; Geoffroy, M. Spin-labelled glycolipid analogues: D-glucose series. *Carbohydr. Res.* **1995**, *275*, 245–258.
15. Peseke, K.; Feist, H.; Hanefeld, W.; Kopf, J.; Schulz, H. Branched-chain sugars as push-pull butadienes. *J. Carbohydr. Chem.* **1995**, *14*, 317–325.
16. Texier-Boullet, F.; Focaud, A. Knoevenagel condensation catalyzed by aluminium oxide. *Tetrahedron Lett.* **1982**, *23*, 4927–4928.
17. Gewald, K.; Kleinert, M.; Thiele, B.; Hetschel, M. Zur basenkatalysierten Reaktion von methylenaktiven Nitrilen mit Schwefel. *J. Prakt. Chem.* **1972**, *314*, 303–314.
18. Gómez, M.; Quincoces, J.; Peseke, K.; Michalik, M.; Reinke, H.J. Synthesis of anellated pyranose derivatives on the basis of levoglucosenone. *J. Carbohydr. Chem.* **1999**, *18*, 851–865.
19. Taylor, E.C.; Berger, J. Heterocyclic syntheses from *o*-aminonitriles. XXIX. A new synthesis of 5-substituted pyrimidines. *J. Org. Chem.* **1967**, *32*, 2376–2378.

Received September 27, 2002

Accepted February 5, 2003

