

Allylic Rearrangements During the Rhodium-catalysed Reactions of 2-Allyloxybenzylamines and 2-(*N*-Allyl-*N*-benzylamino)benzylamine

Eva M. Campi, W. Roy Jackson,* Quentin J. McCubbin and Andrew E. Trnacek

Department of Chemistry, Monash University, Clayton, Vic., Australia 3168

Reactions of 2-allyloxybenzylamines with H₂/CO in the presence of rhodium catalysts give 1,3-benzoxazines *via* a reaction which involves allylic cleavage and carbonylation as demonstrated by crossover experiments.

Rhodium catalysed reactions of some 2-allyloxybenzylamines **1** gave very high isolated yields of substituted 1,3-benzoxazines **2**. Reaction of 2-allyloxybenzylamine **1a** gave 2-isopropyl-3,4-dihydro-1,3-benzoxazine **2a** in 90% isolated yield and the ¹H and ¹³C NMR spectra of the total reaction mixture showed no evidence for the presence of other compounds (Table 1). An authentic sample of **2a** was prepared by condensation of 2-hydroxybenzylamine with 2-methylpropanal in the presence of molecular sieves.¹

Introduction of a 5-bromo substituent into the aryl ring **1b** also gave a quantitative yield of oxazine **2b**. However, a 3-methoxy substituent gave only polymeric material. Methyl substitution at either end of the allyl moiety (**1d**, **1e**) gave the same product (**2d**, **2e**). This result suggested to us that the products arise *via* a mechanism which involves cleavage of the allyl moiety followed by regioselective carbonylation at the internal carbon atom (Scheme 1).

Support for this mechanism came from a reaction of 2-allyloxy-*N,N*-dimethylbenzylamine **3** with H₂/CO/[Rh] in the presence of 2-hydroxybenzylamine **4** which gave the oxazine **2a** together with 2-hydroxy-*N,N*-dimethylbenzylamine, and the oxazine **6**. In the absence of 2-hydroxybenzylamine **4**, reaction of **3** gave an equimolar mixture of the






Table 1 Reactions of 2-allyloxybenzylamines **1** with H₂/CO^a

Substituents	R ¹	R ²	R ³	Yield of 2 (%)
a	H	H	H	90
b	H	H	5-Br	90
c	H	H	3-MeO	— ^b
d	Me	H	H	70
e	H	Me	H	90

^a Reactions were carried out with alkene, [Rh (OAc)₂]₂, and PPh₃ (200:1:4) or alkene and HRh(CO)(PPh₃)₃ (200:1) for 20 h at 40–60 °C, typically using 3 mmol of alkene in ethyl acetate or benzene (10 ml) and 1:1 molar mixture of H₂/CO (2.76 MPa). Yields of material are after distillation. ^b Polymeric material only.



Scheme 1

deallylated material **5** and the linear aldehyde resulting from terminal hydroformylation of **3**. This aldehyde presumably condenses with **4** in the above sequence leading to the oxazine **6**.

Allylic migration in metal catalysed reactions has been reported recently during palladium catalysed alkylation reactions² and in rhodium catalysed silylaminocarbonylations.³ Precedents for the regioselective carbonylation of the allyl residue on the central carbon cannot be found but nucleophilic substitution at the central carbon of a π -allylplatinum complex has been reported.⁴

An analogous rearrangement/carbonylation has also been demonstrated in the rhodium catalysed reaction of the nitrogen analogue **7** which gave the quinazoline derivative **8** in 80–90% yield (NMR) together with a small amount of an unidentified compound.

A rhodium complex **9** of **1a** has been prepared following the method of Krafft⁵ but attempts to induce rearrangement to a π -allylrhodium has so far not been successful.

We thank the Australian Research Council for support including postgraduate awards to (Q. J. McC. and A. E. T.), Johnson Matthey Ltd for a loan of rhodium, and Professor I. Ojima, State University of New York, Stony Brook for discussion.

Received, 2nd September 1994; Com. 4/05367B

References

- A. F. McDonagh and H. E. Smith, *J. Org. Chem.*, 1968, **33**, 1.
- Y. I. M. Nilsson, P. G. Andersson and J. E. Backvall, *J. Am. Chem. Soc.*, 1993, **115**, 6609.
- R. D. Donovan, D. Machnik and I. Ojima, *208th ACS National Meeting*, August, 1994, Washington, DC, Abstracts, ORGN 317.
- K. Ohe, H. Matsuda, T. Morimoto, S. Ogoshi, N. Chatani and S. Murai, *J. Am. Chem. Soc.*, 1994, **116**, 4125.
- M. Krafft, L. J. Wilson and K. D. Onan, *Organometallics*, 1988, **7**, 2528.