



Pergamon

Tetrahedron Letters 40 (1999) 8815–8818

TETRAHEDRON
LETTERS

A new method for synthesis of allenes, including an optically active form, from aldehydes and alkenyl aryl sulfoxides with carbon–carbon bond-formation

Tsuyoshi Satoh,* Yuko Kuramochi and Yujiro Inoue

Department of Chemistry, Faculty of Sciences, Science University of Tokyo, Kagurazaka, Shinjuku-ku, Tokyo 162-8601, Japan

Received 30 July 1999; revised 26 August 1999; accepted 27 August 1999

Abstract

Ligand exchange reaction of β -mesyloxy sulfoxides or β -acetoxy sulfoxides, which were derived from alkenyl aryl sulfoxides and aldehydes in two steps, with *n*-BuLi or EtMgBr at low temperature gave allenes in good yields. Optically active allenes were synthesized starting from optically active 2-phenylethenyl *p*-tolyl sulfoxides. © 1999 Elsevier Science Ltd. All rights reserved.


Allenes are quite important compounds in organic chemistry and a large number of studies have been reported on their chemistry and synthesis.¹ Some allenes have axial chirality and exist as an optically active form. Recently, asymmetric synthesis of allenes has received much attention.²

We previously reported³ a method for synthesizing allenes **2** from sulfoxides **1** by the ligand exchange reaction of sulfoxides with *n*-BuLi.⁴ In continuation of our study on the ligand exchange reaction of sulfoxides in organic synthesis, herein we report a new method for a synthesis of allenes **5**, including an optically active form, from alkenyl aryl sulfoxides **3** and aldehydes via mesylate or acetate **4** (Scheme 1).


According to the procedure reported by Marino,^{5a} addition of the lithium carbanion of propenyl phenyl sulfoxide **6** (a mixture of geometrical isomers) to 3-phenylpropionaldehyde gave the adduct **7a** as a diastereomeric mixture in almost quantitative yield.⁵ Mesylation of the adduct afforded the mesylate **7b** in good yield. Treatment of the mesylate **7** with alkylmetals was carried out in THF at -78°C with three alkylmetals as shown in Scheme 2. The elimination of both the mesyloxy and sulfinyl groups was realized to give the desired allene **8** in good yield. In the cases with *n*-BuLi and *t*-BuLi, the reactions were completed within 30 min; however, the reaction with EtMgBr required 1.5 h for completion.

Other examples of the synthesis of allenes from alkenyl aryl sulfoxides and aldehydes are summarized in Table 1. When R¹ and/or R² is an aromatic group, the mesylate (**4**: R=Ms) is quite unstable due to the formation of an allylic cation. In these cases we obtained the acetates (**4**: R=COCH₃) in high yields (entries 1–4). As shown in entry 1, the elimination reaction of the acetate with both *n*-BuLi and

* Corresponding author.



Scheme 1.



Scheme 2.

t-BuLi gave only a complex mixture. Fortunately, the elimination was successful with EtMgBr to give the desired allene in 88% yield. Entry 4 shows that in the case that both R¹ and R² are an alkyl group, much better yields were obtained with the mesylate than with the acetate (compare the results in entry 4 and that in Scheme 2). Entry 6 shows the synthesis of 3-(2-naphthyl)-1,2-propadiene. The addition of the carbanion of phenyl vinyl sulfoxide with β-naphthaldehyde gave a moderate yield of the adduct; however, the elimination of the acetate with EtMgBr gave a quantitative yield.




Next, we investigated the asymmetric synthesis of allenes starting from the optically active 2-phenylethenyl *p*-tolyl sulfoxide⁶ **9** with β-naphthaldehyde and 3-phenylpropionaldehyde (Scheme 3).

Addition of the lithium carbanion of **9** to β-naphthaldehyde at -100°C gave a mixture of two diastereomers, which were separated by silica gel column chromatography (**10a**, 53%; **11a**, 43%). The newly created chiral centers were assigned as *R* and *S*, respectively, as shown in Scheme 3, by comparison of the data with those of Marino's report.^{5a}

Acetylation of the adducts **10a** and **11a** gave the acetates **12a** ([α]_D +121) and **13a** ([α]_D +1.5), respectively, in almost quantitative yields. The elimination of the acetate **12a** in THF with EtMgBr was found to be rather sluggish; however, use of 10 equivalents of EtMgBr at -25°C for 20 min gave the desired allene (*R*)-**14a**⁷ ([α]_D -531) in 92% yield as colorless crystals. The enantiomeric excess was calculated to be 68% by HPLC using a chiral column (Daicel, Chiralpak AD, hexane:*i*-PrOH=9:1). Recrystallization of the product with AcOEt-hexane gave the allene having much higher optical purity (mp 85.5–87°C; [α]_D -668 (c 0.55, CHCl₃), 95% ee by HPLC). Similar treatment of **13a** in THF at -25°C with 4 equivalents of EtMgBr for 10 min gave the optically active allene (*S*)-**15a** ([α]_D +515; 74% ee) as colorless crystals in 73% yield.

Next, the adducts of **9** with 3-phenylpropionaldehyde (**10b** and **11b**) were acetylated to give the acetates **12b** ([α]_D +68) and **13b** ([α]_D +51). Treatment of **12b** with 14 equivalents of EtMgBr at -30°C

Table 1
Synthesis of allenes **5** from alkenyl sulfoxides **3** and aldehydes through the ligand exchange reaction of sulfoxides

Entry	3		R ² CHO	4	Alkylmetal	5
	Ar	R ¹	R ² (Yield / %) ^{a)}	R (Yield / %) ^{b)}		Yield / % ^{c)}
1	Ph	Ph	PhCH ₂ CH ₂ (87)	Ac (96)	<i>n</i> -BuLi <i>t</i> -BuLi EtMgBr	— ^{d)} — ^{d)} 88
2	Ph	Ph	 (99)	Ac (92)	EtMgBr ^{e)}	92
3	Ph	CH ₃	 (77)	Ac (94)	EtMgBr ^{f)}	77
4	Ph	CH ₃	PhCH ₂ CH ₂ (92)	Ac (86)	EtMgBr	42
5	<i>p</i> -Tol	CH ₂ (CH ₂) ₃	PhCH ₂ CH ₂ (82)	Ms (82)	<i>n</i> -BuLi EtMgBr	90 77
6	Ph	H	 (53)	Ac (95)	EtMgBr	99

a) The yield of the reaction of **3** with the aldehyde. b) The yield for the acetylation or mesylation. c) Isolated purified yield after silica gel column chromatography. d) A complex mixture was obtained. e) at -25 °C, 20 min. f) at -100 °C, 2 h.


for 1 h gave the allene (*R*)-**14b** ($[\alpha]_D -63$; 21% ee, by Daicel Chiralcel OD) in 93% yield as an oil. Similar treatment of **13b** with 6 equivalents of EtMgBr at -30 °C for 1 h gave allene (*S*)-**15b** ($[\alpha]_D +101.5$; 61% ee) in 94% yield. The reason why the optical yield of **14b** is so low is obscure at present.

By comparison of the absolute configurations of **12** and **13** with those of **14** and **15**, the elimination was thought to proceed mainly via the *anti*-conformation as shown in Scheme 3.

We are continuing to study the scope and limitations of this method and to improve the optical yield of the elimination step.

Acknowledgements

This work was supported by a Grant-in-Aid for Scientific Research no. 11640545 from the Ministry of Education, Science, Sports, and Culture, which is gratefully acknowledged.



Scheme 3.

References

- (a) Patai, S. *The Chemistry of Ketenes, Allenes, and Related Compounds*; John Wiley & Sons: Chichester, 1980; Parts 1 and 2. (b) Brandsma, L.; Verkruisje, H. D. *Synthesis of Acetylenes, Allenes, and Cumulenes*; Elsevier: Amsterdam, 1981. (c) Schuster, H. F.; Coppola, G. M. *Allenenes in Organic Synthesis*; John Wiley & Sons: New York, 1984.
- Some selected recent papers for asymmetric synthesis of allenes: (a) Marek, I.; Mangeney, P.; Alexakis, A.; Normant, J. F. *Tetrahedron Lett.* **1986**, *27*, 5499. (b) Alexakis, A.; Marek, I.; Mangeney, P.; Normant, J. F. *J. Am. Chem. Soc.* **1990**, *112*, 8042. (c) Gooding, O. W.; Beard, C. C.; Jackson, D. Y.; Wren, D. L.; Cooper, G. F. *J. Org. Chem.* **1991**, *56*, 1083. (d) Shepard, M. S.; Carreira, E. M. *Tetrahedron* **1997**, *48*, 16253. (e) Li, A.-H.; Dai, L.-X.; Aggarwal, V. K. *Chem. Rev.* **1997**, *97*, 2341. (f) Tanaka, K.; Fujii, K. *J. Syn. Org. Chem. Jpn.* **1998**, *56*, 521. (g) Franck-Neumann, M.; Martina, D.; Neff, D. *Tetrahedron: Asymmetry* **1998**, *9*, 697. (h) Noguchi, Y.; Takiyama, H.; Katsuki, T. *Synlett* **1998**, 543.
- Satoh, T.; Itoh, N.; Watanabe, S.; Koike, H.; Matsuno, H.; Matsuda, K.; Yamakawa, K. *Tetrahedron* **1995**, *51*, 9327.
- Satoh, T. *J. Syn. Org. Chem. Jpn.* **1996**, *54*, 481.
- (a) Marino, J. P.; Viso, A.; Lee, J.-D. *J. Org. Chem.* **1997**, *62*, 645. (b) *Organosulfur Chemistry, Synthetic and Stereochemical Aspects*; Page, P., Ed.; Academic Press: San Diego, 1998.
- Makolajczyk, M.; Perlikowska, W.; Omelanczuk, J.; Cristau, H.-J.; Perraud-Darcy, A. *J. Org. Chem.* **1998**, *63*, 9716.
- The absolute configuration was determined by the sign of optical rotation: see Ref. 1c, pp. 31–32.