

# RSC Advances

## Accepted Manuscript

This article can be cited before page numbers have been issued, to do this please use: G. Xia, H. Wang and Y. Yuan, *RSC Adv.*, 2013, DOI: 10.1039/C3RA43608J.



This is an *Accepted Manuscript*, which has been through the RSC Publishing peer review process and has been accepted for publication.

*Accepted Manuscripts* are published online shortly after acceptance, which is prior to technical editing, formatting and proof reading. This free service from RSC Publishing allows authors to make their results available to the community, in citable form, before publication of the edited article. This *Accepted Manuscript* will be replaced by the edited and formatted *Advance Article* as soon as this is available.

To cite this manuscript please use its permanent Digital Object Identifier (DOI®), which is identical for all formats of publication.

More information about *Accepted Manuscripts* can be found in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics contained in the manuscript submitted by the author(s) which may alter content, and that the standard [Terms & Conditions](#) and the [ethical guidelines](#) that apply to the journal are still applicable. In no event shall the RSC be held responsible for any errors or omissions in these *Accepted Manuscript* manuscripts or any consequences arising from the use of any information contained in them.

Cite this: DOI: 10.1039/c0xx00000x

www.rsc.org/xxxxxx

ARTICLE TYPE

# Regioselectivity and Synthetic Mechanism of 1,2-Benzimidazole Squaraines: Combined Experimental and Theoretical Studies

Guomin Xia, Zhiwei Wu, Yanli Yuan, and Hongming Wang\*

Received (in XXX, XXX) Xth XXXXXXXXX 200X, Accepted Xth XXXXXXXXX 200X  
DOI: 10.1039/b000000x

In this paper, a synthetic method to produce 1,2-benzimidazole squaraines with a yield of up to 89% is developed. It is found that both strong organic and inorganic bases have satisfied catalyzed activity for this reaction. Theoretical studies provide detailed explanations for the 1,2 *versus* 1,3 condensation regiochemistry of the squaraines. Experimental and theoretical studies agree well with each other, paving a practical way to efficiently synthesize 1,2-squaraines.

## Introduction

Squaraine (SQ) dyes are the product of condensation of squaric acid and two equivalents of a suitable electron-rich precursor<sup>1</sup> and be notable for their exceptionally high absorption coefficients extending from the green to the near-infrared region.<sup>2,3</sup> This has prompted their exploitation in a number of technologically relevant applications, including photoconductivity,<sup>4</sup> data storage,<sup>5</sup> light-emitting field-effect transistors,<sup>6</sup> solar cells,<sup>7</sup> and fluorescent histological probes.<sup>8</sup> Since the pioneering work of Cohen in 1959,<sup>9</sup> SQs have also been used in the fields of bioorganic and medicinal chemistry as well.<sup>10</sup> For example, Xie *et al.* demonstrated that SQ is an effective pharmacophore for the design of tyrosine phosphatase inhibitors.<sup>11</sup> The synthesis of squaraine dyes was first reported in the 1960s.<sup>12</sup> Then numerous reports describing the synthesis and properties of SQs have appeared in recent decades.<sup>13</sup> According to previous investigations, SQs are the condensation product of two electron-rich derivatives (activated arenes, 2,4-dimethylpyrrole and benzothiazole) and squaric acid or their esters, which generally affords the 1,3-regioisomer as the main product and the 1,2-isomer as a by-product. Modification of the reaction conditions only slightly increases the regioselectivity toward 1,2-SQ; the reaction is regioselective and the major products are still the 1,3-isomers. Although some studies indicate that 1,2-SQ can be synthesized through a two-step Friedel-Crafts acylation reaction between squaric acid chloride and bromo-substituted thiophene, its direct synthesis from squaric acid or squarate have seldom been reported. As we all know, 1,2-SQs are considered to be promising as sensitizers in organic photoconducting devices and electron-acceptor components for construction of high-performance D-A-type conjugated photovoltaic polymers,<sup>[14]</sup> so a facile and high selectivity route to synthesize 1,2-SQs is urgently required. Ramaiah's group achieved the selective synthesis of 1,2- and 1,3-

semisquaraine<sup>15</sup> and identified the 1,3-isomer as the reactive intermediate in squaraine dye formation rather than the neutral 1,2-isomer. Their investigations indicate that the base influences the regioselectivity of 1,2- and 1,3-semisquaraine. Ronchi *et al.* identified the main electronic factors in this reaction and changed its conditions, increasing the yield of 1,2-SQs up to 66%.<sup>16</sup>

It is clear that the synthesis of 1,2-SQs is of significant challenge, and the yield as well as selectivity still remains unsatisfactory. In this paper, we report a synthetic method to produce 1,2-benzimidazole squaraines (3a-b) at a high yield. To explore the reason for the high regioselectivity of 3a-b, the overall reaction mechanisms were studied by density functional theory (DFT) calculations as well. By combining the experimental and computational findings, we are able to propose a method of chemical regulation that changes the regioselectivity of the product. This work sheds light on the factors influencing reaction via the 1,2- versus the 1,3-condensation pathway, which could provide guidelines for the synthesis of desired 1,2-SQs in the future.


## Results and Discussion

Considering the most of synthesis of 1,3-SQs in previous reports, we planned to produce 1,3-SQ (2a) by reacting benzimidazole salt 1a with 3,4-diethyl squarate in the presence of a frequently-used base (triethylamine (Et<sub>3</sub>N)) in EtOH (Scheme 1), however, no products formed after heating under reflux for 12 h, and the 1a didn't react at all. Then, a fairly strong base was been used to catalyze the reaction. As a result, the reaction mixture turned from yellow to deep red instantly when diazabicycloundecene (DBU) was been added, after 3h reflux, a dull-red product was isolate with 68% yield and crystallized by the slow evaporation method. To our surprise, X-ray diffraction analysis of the product showed that the red product was 1,2-SQ 3a (Figure 1) instead of 1,3-SQ 2a. Also, 3a was characterized by various spectroscopic and analytical techniques: <sup>1</sup>H and <sup>13</sup>C NMR spectroscopy, FT-

IR and elemental analysis.

The results show that we should draw to the thought that the basicity of the base has a large effect towards the products; this prompted us to use a stronger base to verify this inference.

Sodium ethoxide (EtONa), as a common organic base, was used to catalyze the condensation reaction between 1a-b and 3,4-diethyl squarate. As the way that was expected, red deposits formed immediately as long as EtONa was been added, after 3h reflux, the reactions were completed to produce 3a and 3b in 89% and 86% yield, respectively. Moreover, in these reactions, 1,3-SQs (2a and 2b) were not been observed. However, when squaric acid was reacted with 1a-b in EtOH solvent, no red product 1,2-SQs formed after heating under reflux for 3h (Scheme 2). The weaker base pyridine was also been used to catalyze the condensation reactions between 1a-b and 3,4-diethyl squarate; the mixture was always beige, and no red deposit formed in these reactions, after heating under reflux for 12h, no 1,2- or 1,3-SQs formed, which is similar to the reaction using Et<sub>3</sub>N as the base.




Scheme 1. The reaction of benzimidazole salt with 3,4-diethyl squarate.

Table 1. Reaction of Benzimidazole Salts (1a-b) and 3,4-Diethyl Squarate catalyzed by various bases.

	R	Base	Time (h)	Product	Yield (%)
1	PhCH <sub>2</sub>	pyridine	12	<b>3a</b>	0
2	CH <sub>3</sub>	pyridine	12	<b>3b</b>	0
3	PhCH <sub>2</sub>	Et <sub>3</sub> N	12	<b>3a</b>	0
4	CH <sub>3</sub>	Et <sub>3</sub> N	12	<b>3b</b>	0
5	PhCH <sub>2</sub>	DBU	3	<b>3a</b>	68
6	CH <sub>3</sub>	DBU	3	<b>3b</b>	61
7	PhCH <sub>2</sub>	EtONa	3	<b>3a</b>	89
8	CH <sub>3</sub>	EtONa	3	<b>3b</b>	86

Based on the results above, we wanted to determine that whether an inorganic base can be used to catalyze these reactions. The yields of the reactions utilizing various inorganic bases are presented in Table 2. It indicated that inorganic bases also can catalyze this condensation reactions and the basicity of the base worked as the same as organic bases: only trace amounts of **3a** and **3b** were observed (inspected by TLC chromatography) when a weak inorganic bases (NaHCO<sub>3</sub>, KHCO<sub>3</sub>, Na<sub>2</sub>CO<sub>3</sub> and K<sub>2</sub>CO<sub>3</sub>) was been used after refluxing 12h; However, with 250%mol strong base Cs<sub>2</sub>CO<sub>3</sub>, NaOH, KOH, the yield of **3a** and **3b** were dramatically improved towards to 76% and 73%, 81% and 75%, 85% and 80%, respectively.



Scheme S2. The reaction between benzimidazole salt and squaric acid catalyzed by EtONa.

Table 2. Reaction of Benzimidazole Salts (1a-b) and 3,4-Diethyl Squarate under Various Inorganic Bases.

	R	Base	Time (h)	Product	Yield (%)
1	PhCH <sub>2</sub>	NaHCO <sub>3</sub>	12	<b>3a</b>	trace
2	CH <sub>3</sub>	NaHCO <sub>3</sub>	12	<b>3b</b>	trace
3	PhCH <sub>2</sub>	KHCO <sub>3</sub>	12	<b>3a</b>	trace
4	CH <sub>3</sub>	KHCO <sub>3</sub>	12	<b>3b</b>	trace
5	PhCH <sub>2</sub>	Na <sub>2</sub> CO <sub>3</sub>	12	<b>3a</b>	trace
6	CH <sub>3</sub>	Na <sub>2</sub> CO <sub>3</sub>	12	<b>3b</b>	trace
7	PhCH <sub>2</sub>	K <sub>2</sub> CO <sub>3</sub>	12	<b>3a</b>	trace
8	CH <sub>3</sub>	K <sub>2</sub> CO <sub>3</sub>	12	<b>3b</b>	trace
9	PhCH <sub>2</sub>	Cs <sub>2</sub> CO <sub>3</sub>	3	<b>3a</b>	76
10	CH <sub>3</sub>	Cs <sub>2</sub> CO <sub>3</sub>	3	<b>3b</b>	73
11	PhCH <sub>2</sub>	NaOH	3	<b>3a</b>	81
12	CH <sub>3</sub>	NaOH	3	<b>3b</b>	75
13	PhCH <sub>2</sub>	KOH	3	<b>3a</b>	85
14	CH <sub>3</sub>	KOH	3	<b>3b</b>	80



Figure 1. ORTEP diagram of **3a**.

Absorption spectra of **3a** and **3b** in selected solvents are presented in Fig. 2A and 3A (see the ESI). The spectra indicate a solvent-dependent change in the absorption maxima and intensity of **3a**. The absorption maximum changes considerably in methanol compared with in other solvents. The absorption coefficient of **3a** in dichloromethane is larger than in any other solvent. Similar results were also observed for **3b**. In Fig. 2B, the absorption peak at 524 nm decreases and that at 399 nm increases gradually as the concentration of TFA increases, which implies a strong pH dependence of the visible absorption spectra of **3a**. **3a** was also investigated by time-dependent density functional theory (TDDFT) calculations. Fig. S5 shows the calculated ground state structures and molecular orbitals of **3a**. From TDDFT calculations, the long-wavelength absorption (524 nm) is mainly generated by the HOMO→LUMO transition. The short-wavelength absorption (433 nm) corresponds to the promotion of an electron from the HOMO-2 to the LUMO-1 (see Figure. S5).




Figure 2. Absorption spectra of the 1,2-SQ 3a in various solvents (A). Absorption spectra change of 3a in acetone upon addition of different ratio of trifluoroacetic acid (TFA) (B).





Figure 3. Absorption spectra of the 1,2-SQ 3b in various solvents (A). Absorption spectra change of 3a in acetone upon addition of different ratio of trifluoroacetic acid (TFA) (B).

To examine the different reaction activity of various bases and their effect on the regioselectivity of this reaction, DFT calculations were performed to explore the detailed reaction mechanism[17] (Scheme S3). In the first step of this reaction, 1b reacted with different bases (EtONa, DBU, Et3N, and pyridine) to produce PRa complexes. Fig. 4A shows the energy profile for this deprotonation step. The H1 atom is attacked by the bases (EtONa, DBU, Et3N, and pyridine) to produce an INa (INa1, INa2, INa3, and INa4) complex. All of these complexes have lower energies than the reactants (Fig. 2A). Because of the strong electron-donating ability of the lone pairs of electrons of the bases, the H1 atom is then transferred from 1b to the base to afford 1a-base (INb) intermediates via transition states (TSa) (Fig. 4A). Also, Fig. 4A shows that TSa1 has lower energy (-18.69 kcal mol<sup>-1</sup>) than the reactant. However, the energy barriers of TSa2, TSa3 and TSa4 are 7.42, 14.84 and 16.37 kcal mol<sup>-1</sup>, respectively, which are higher than those of the reactants. Subsequently, the products, PRa1, PRa2, PRa3 and PRa4 are formed from the INb complexes. Fig. 4A reveals that PRa1 and PRa2 are more stable than the reactants by 30.68 and 0.44 kcal mol<sup>-1</sup>, respectively. However, the energies of PRa3 and PRa4 are 14.10 and 23.47 kcal mol<sup>-1</sup> higher than those of the corresponding reactants. Because of the low energy barriers and exothermic characteristics, the reactions catalyzed by EtONa and DBU proceed easily. For Et3N and pyridine, however, both reactions are strongly endothermic with relatively high energy barriers, it is understandable that these two reactions do not proceed readily. In addition, EtONa has the highest activity during all of the bases used. These conclusions are consistent with the experimental results mentioned above where the reaction catalyzed by EtONa gives a high yield of 1,2-SQ (89%).







Figure 4. Energy profiles for (A) the deprotonation of benzimidazole salt by base, and (B) the formation of PRb.

After deprotonation, the benzimidazole methylene (PRA) was formed (Scheme S1 and Fig. 4A). Because of nucleophilicity of PRA, it is able to undergo a 1,4-conjugated attack at the squarate (FS) double bond (Fig. 4B). The step involves a transition state (TSb) that has a low active barrier (13.3 kcal mol<sup>-1</sup>). An intermediate (INC) is then formed with an energy 2.31 kcal mol<sup>-1</sup> above that of the reactants. The next step is a 1,5-proton shift between the methylene of PRA and enolate (see Fig. 4B and Fig. S6 in the ESI). Finally, thermally-catalyzed elimination of ethanol gives exclusively PRb. In fact, this step (TSc) has a very high energy barrier (44.01 kcal mol<sup>-1</sup>). This barrier is too large to enable the reaction to be conducted experimentally at ambient temperature, so heating is required. Because the energy barriers involve proton transfer, we considered that weather EtOH played an important role in this process. When EtOH was introduced as the proton transfer catalyst to assist in forming TScs (six-member ring transition state), the energy barrier was reduced considerably to 16.21 kcal mol<sup>-1</sup> (Fig. 4B). This experiment also indicates that the yield and reaction rate would be increased considerably in ethanol compared with that in DMF catalyzed by DBU (Scheme 3).



**Scheme 3.** Reaction of Benzimidazole Salts (1a-b) and 3,4-Diethyl Squarate catalyzed by DBU in dry DMF.




Figure 5. Energy profiles for the formation of 3b (pathways: I-A) and the structures of transition states (B: TSd, C: TSe).

Because of its nucleophilicity, PRA can continue to attack the C1 and C2 atoms of PRb through two reaction pathways: I-A and I-B (see Scheme S1). In reaction pathway I-A, the C3 atom of PRA attacks the C1 atom of PRb to produce an intermediate INd. This process involves transition state TSd with an energy barrier of 16.3 kcal mol<sup>-1</sup> (figure 5 and Figure. S8). After INd, transition state TSe with an energy 37.58 kcal mol<sup>-1</sup> above that of the reactants forms (Figure. S8). TSe involves a proton transfer (H8) between the O9 and C7 atoms, and leads to products 3b and EtOH. When EtOH is introduced as a cocatalyst, the energy barrier (TSe) decreases to 18.49 kcal mol<sup>-1</sup>.




Figure 6. Energy profiles for the formation of 2b (pathways: I-B) and the structures of transition states (B: TSg, C: TSi).



As displayed in the figure 6 and Figure. S9, in reaction pathway **I-B**, the C3 atom of PRa attacks the para-C2 atom of PRb to produce an intermediate INF *via* transition state TSf with an energy barrier of 16.12 kcal mol<sup>-1</sup>. After INF, a transition state TSg involving proton transfer (H8) from PRa to PRb with an energy barrier of 45.12 kcal mol<sup>-1</sup> occurs. When EtOH is introduced as a proton transfer catalyst to assist this step, the energy barrier was reduced to 17.32 kcal mol<sup>-1</sup>. In the next step, an OH group on the C2 atom transfers to the C1 atom. This process proceeds *via* the transition state TSh, which is 36.1 kcal mol<sup>-1</sup> higher in energy than the reactants. After TSh, INg is formed with an energy of 3.07 kcal mol<sup>-1</sup> above that of the reactants. The last step for this pathway is proton transfer from the OH group on C1 to O atom of the ethoxyl. This process results in product **2b** *via* transition state TSi. The computed activation barrier for TSi is 19.97 kcal mol<sup>-1</sup>. From the analyses above, it can be concluded that pathway **I-A** is the favorable reaction route, because its highest activation barrier (17.58 kcal mol<sup>-1</sup>) is lower than that of channel **I-B** (36.1 kcal mol<sup>-1</sup>). This explains well the high regioselectivity for 1,2-SQs (**3a-b**) in our experiments.

## Conclusions

In summary, a method to prepare 1,2-benzimidazole squaraines in high yield *via* one step reaction is developed. The yield of 1,2-benzimidazole squaraines is up to 89%. The total reaction mechanism is explored by DFT calculations. By combining experimental and computational findings, we proposed a chemical regulation method to change the regioselectivity of the product. Our experimental and theoretical studies agree well with each other, paving a practical way to synthesize of 1,2-squaraines in high yield. Syntheses of other 1,2-SQ complexes will be reported in the future.

## Experimental Section

### Calculation detailed

Geometrical optimization, including transition-structure searches, was carried out with the standard 6-311G(d,p) basis set by using the three-parameter hybrid functional developed by Becke in the formulation implemented in the Gaussian 09 program<sup>17</sup> (B3LYP), which is slightly different from the original proposed by Becke.<sup>18</sup> The transition states are ascertained by vibrational analysis with only one imaginary frequency mode. In the case of TS, the vibration associated with the imaginary frequency was checked to correspond with a movement in the direction of the reaction coordinate.<sup>19</sup> The values of the relative energies,  $\Delta E$ , have been calculated on the basis of the total energies of the stationary points. The solvent effects have been considered using a relatively simple self-consistent reaction field (SCRF) method, based on the polarizable continuum model (PCM).<sup>20</sup> The solvent used in this calculation is ethanol.

### General Remarks

All starting chemicals and solvents were purchased from commercial suppliers and used as received unless explicitly stated. Absorption spectrometry was performed using Gold S54T spectrophotometer of Lenggung Company. Melting points were

determined in open capillaries. <sup>1</sup>H NMR and <sup>13</sup>C NMR spectra were measured on a Bruker AVANCE 400 spectrometer in DMSO-d<sub>6</sub> using TMS as an internal standard. The crystals for X-ray diffraction analyses were grown in dichloromethane by adding n-hexane to slow diffuse. The X-ray diffraction experiment was carried out using Bruker SMART APEX-II Single-crystal diffractometer.

### Synthesis of 2-methyl benzimidazole (a)

Phenylene diamine (0.3mol, 32.4g) and acetic acid (1mol, 60ml) were refluxed for 2h. Ice and KOH was added to PH=8~10 and the light violet solid filtered out, then active carbon was used and recrystallized from water to giving 2-methyl benzimidazole (a) as light yellow needle crystals (30g, yield 77%), M.p 176~178 °C.<sup>21</sup>

### Synthesis of 1,3-dibenzyl-2-methyl-benzimidazolium chloride (1a)

To a solution of 2-methylbenzimidazole (1 g, 7.57 mmol) in DMF (20 ml) was added benzyl chloride (2.66 ml, 22.7 mmol), potassium carbonate (1.25 g, 9.08 mmol) and a catalytic amount of tetra-n-butylammonium bromide. The mixture was stirred at room temperature for 24 h. The solution was filtered and the solvent removed under reduced pressure. The residue was recrystallized from ethanol to afford 1,3-dibenzyl-2-methylbenzimidazolium chloride (1a) as colorless amorphous crystals (0.8g, yield 27%). M.p 312~314 °C<sup>22</sup>; <sup>1</sup>H NMR (400 MHz, DMSO-d<sub>6</sub>): 7.92 (d, J = 1.67 Hz, 2H), 7.57 (d, J = 0.76 Hz, 2H), 5.81 (s, 4H), 7.36 (s, 10H), 2.99 (s, 3H); <sup>13</sup>C NMR (400 MHz, DMSO-d<sub>6</sub>  $\delta$ ): 152.9 (1C), 134.6 (2C), 131.5 (2C), 129.4 (4C), 128.8 (4C), 127.49 (2C), 126.8 (2C), 113.8 (2C), 48.8 (2C), 11.5(1C).

### Synthesis of 1,2,3-trimethyl benzimidazolium iodide (1b)

In 250ml round bottomed flask, Na (0.04mol 0.9g) was dissolved in 16ml ethanol, after this 2-methyl benzimidazole (0.04mol 5.2g), 100ml benzene and MeI (0.12mol 7.5ml) were added successively, the mixtures were refluxed for 18h, then allowed to cool to room temperature, and the solvent was removed under reduced pressure. The crude product was purified by recrystallization from 95% ethanol to afford 1,2,3-trimethyl benzimidazolium iodide (1b) as colorless needle crystals (9.2g, yield 80%). M.p 258~259 °C<sup>23</sup>; <sup>1</sup>H NMR (400 MHz, DMSO-d<sub>6</sub>  $\delta$ ): 7.97 (dd, J = 5.97, 2.93 Hz, 2H), 7.62 (dd, J = 6.01, 2.83 Hz, 2H), 3.99 (d, J = 14.04 Hz, 6H), 2.85 (s, 3H); <sup>13</sup>C NMR (400 MHz, DMSO-d<sub>6</sub>  $\delta$ ): 152.7 (1C), 131.8 (2C), 126.2 (2C), 113.1(2C), 32.2 (2C), 11.1(1C).

### General Procedure for the Synthesis of 1,2-Squaraines

All of the experiments described above were performed according to the following general procedure. A mixture of 2mmol of squaric acid in 30ml anhydrous ethanol was refluxed at 100 °C for 3h under a nitrogen atmosphere, then the solvent was removed under reduced pressure and an other 30ml steamed anhydrous ethanol was added, the mixture was refluxed for 30min and the solvent was removed again, repeating this for at least three times in order to get diethyl squarate generated completely. The base (10mmol Organic base: EtONa, DBU,

Et<sub>3</sub>N or pyridine; 5mmol Inorganic base: NaHCO<sub>3</sub>、KHCO<sub>3</sub>、Na<sub>2</sub>CO<sub>3</sub>、K<sub>2</sub>CO<sub>3</sub>、Cs<sub>2</sub>CO<sub>3</sub>、NaOH or KOH)、4mmol of the benzimidazole salt and 15ml anhydrous ethanol was added, the mixture was refluxed for 3h, then allowed to cool to room temperature. The crude reaction mixture is inspected by TLC chromatography. The products were determined by comparison with the analytical samples of the various squaraines obtained by the procedures described in details in the following paragraph.

#### 10 Synthesis of 3,4-bis((1,3-dibenzyl-1H-benzo[d]imidazol-2(3H)-ylidene)methyl) cyclobut-3-ene-1,2-dione (3a)

A mixture of squaric acid (2mmol, 228mg) in 30ml anhydrous ethanol was refluxed at 100 °C for 3 h under a nitrogen atmosphere, then the solvent was removed under reduced pressure and an other 30ml steamed anhydrous ethanol was added, the mixture was refluxed for 30min and the solvent was removed again, repeating this for at least three times in order to made diethyl squarate generated as much as possible. In the meantime Na (230mg, 10mmol) was dissolved in 10ml anhydrous ethanol, then the solution of sodium ethoxide、1a (4mmol, 1.39g) and 15ml anhydrous ethanol was added, the mixtures turned to deep red at once and then was refluxed for 3h. the solvent was removed under reduced pressure and 15ml isopropanol was added to separate garnet solid out, the solution was filtered and the residue was washed by isopropanol、ether、petroleum ether alternately to give 3a as a garnet solid (1.25g, yield 89%). M.p 217~218°C; <sup>1</sup>H NMR (400 MHz, DMSO-d<sub>6</sub>): 7.35 (dd, J = 4.88, 2.57 Hz, 4H), 7.17 (t, J = 6.53 Hz, 12H), 7.11-7.07 (m, 4H), 7.01 (d, J = 6.91 Hz, 8H), 5.42 (s, 8H), 4.70 (s, 2H); <sup>13</sup>C NMR (400 MHz, DMSO-d<sub>6</sub>): 189.4 (2C), 170.9 (2C), 151.1 (2C), 136.6 (4C), 133.3 (4C), 129.1 (8C), 128.0 (8C), 127.2 (4C), 123.3 (4C), 110.3 (4C), 65.9 (2C), 47.8 (4C); Anal. Calcd for 3a: C, 81.95; H, 5.41; N, 7.96. Found: C, 81.73; H, 5.87; N, 7.67.

#### 15 Synthesis of 3,4-bis((1,3-dimethyl-1H-benzo[d]imidazol-2(3H)-ylidene)methyl) cyclobut-3-ene-1,2-dione (3b).

A mixture of squaric acid (2mmol, 228mg) in anhydrous ethanol was reacted as the same as the steps described above. In the meantime Na (230mg, 10mmol) was dissolved in 10ml anhydrous ethanol. The solution of sodium ethoxide、1b (4mmol, 1.15g) and 15ml anhydrous ethanol was added, the mixture turned from yellow to blood red immediately and was refluxed for 3h, then allowed to cool to room temperature, the solution was filtered and the residue was washed by ethanol、ether、petroleum ether in turn to give a pure bright red solid product 3b (684mg, yield 86%). M.P>300 °C; <sup>1</sup>H NMR (400 MHz, DMSO-d<sub>6</sub>): 7.39-7.34 (m, 4H), 7.22-7.18 (m, 4H), 4.75-4.72 (m, 2H), 3.67 (s, 12H); Anal. Calcd for 3b: C, 72.36; H, 5.53; N, 14.07. Found: C, 72.25; H, 5.64; N, 14.03.

#### Acknowledgments

The support of the National Natural Science Foundation of China (10947171 and 21103082) is gratefully acknowledged.

#### Notes and references

- <sup>a</sup> Institute for Advanced Study, Nanchang University, <sup>b</sup> Department of Chemistry, Nanchang University, Nanchang, China, 330031. Fax: 0086-791-3969552 Tel: 0086-791-3969552 E-mail: Hongmingwang@ncu.edu.cn
- † Electronic Supplementary Information (ESI) available: Cartesian coordinates and calculated energies for all structures used in the quantum chemical calculations. See DOI: 10.1039/b000000x/
1. (a) J. D. Park, S. Cohen, J. R. Lacher, J. Am. Chem. Soc. 2002, 84, 2919-2922; (b) J. J. McEwen, K. Wallace, J. Chem. Commun. 2009, 6339-6351.
2. (a) F. Silvestri, M. D. Irwin, L. Beverina, A. Facchetti, G. A. Pagani, T. J. Marks, J. Am. Chem. Soc. 2008, 130, 17640-17641; (b) G. D. Wei, R. R. Lunt, K. Sun, S. Y. Wang, M. E. Thompson, S. R. Forrest, Nano Lett. 2010, 10, 3555-3559; (c) U. Mayerhöfer, F. Wüthner, Chem. Sci. 2012, 3, 1215-1220.
3. (a) A. Ajayaghosh, P. Chithra, R. Varghese, Angew. Chem. Int. Ed. 2007, 46, 230-233; (b) U. Mayerhöfer, F. Wüthner, Angew. Chem. Int. Ed. 2012, 51, 5615-5619; (c) S. Sreejith, P. Carol, P. Chithra, A. Ajayaghosh, J. Mater. Chem. 2008, 18, 264-274.
4. (a) X. Zhang, J. Jie, W. Zhang, C. Zhang, L. Luo, Z. He, X. Zhang, W. Zhang, C. Lee, S. Lee, Adv. Mater. 2008, 20, 2427-2432; (b) K. Y. Law, Chem. Rev. 1993, 93, 449-486; (c) H. Choi, J.-J. Kim, K. Song, J. Ko, M. K. Nazeeruddin, M. Grätzel, J. Mater. Chem. 2010, 20, 3280-3286.
5. (a) A. Dualeh, J. H. Delcamp, M. K. Nazeeruddin, M. Grätzel, App. Phys. Lett. 2012, 100, 173512; (b) M. Emmelius, G. Pawlowski, H. Vollmann, Angew. Chem. Int. Ed. 1989, 28, 1445-1471; (c) R. Petermann, M. Tian, S. Tatsuura, M. Furuki, Dyes and Pigments. 2003, 57, 43-54.
6. E. C. P. Smits, S. Setayesh, T. D. Anthopoulos, M. Buechel, W. Nijssen, R. Coehoorn, P. W. M. Blom, B. de Boer, D. M. de Leeuw, Adv. Mater. 2007, 19, 734-738.
7. (a) A. Piechowski, G. Bird, D. Morel, E. Stogryn, J. Phys. Chem. 1984, 88, 934-950; (b) K. Liang, K.-Y. Law, D. G. Whitten, J. Phys. Chem. 1995, 99, 16704-16708; (c) J.-H. Yum, P. Walter, S. Huber, D. Rentsch, T. Geiger, F. Nuesch, F. D. Angelis, M. Grätzel, M. K. Nazeeruddin, J. Am. Chem. Soc. 2007, 129, 10320-10321; (d) Y. Shi, R. B. M. Hill, J.-H. Yum, A. Dualeh, S. Barlow, M. Grätzel, S. R. Marder, M. K. Nazeeruddin, Angew. Chem. Int. Ed. 2011, 50, 6619-6621; (e) U. Mayerhöfer, K. Deing, K. Gruff, H. Braunschweig, K. Meerholz, F. Wüthner, Angew. Chem. Int. Ed. 2009, 48, 8776-8779; (f) G. Wei, X. Xiao, S. Wang, J. D. Zimmerman, K. Sun, V. V. Diev, M. E. Thompson, S. R. Forrest, Nano Lett. 2011, 11, 4261-4264; (g) X. Xiao, G. Wei, S. Wang, J. D. Zimmerman, C. K. Renshaw, M. E. Thompson, S. R. Forrest, Adv. Mater. 2012, 24, 1956-1960.
8. (a) E. Arunkumar, C. C. Forbes, B. C. Noll, B. D. Smith, J. Am. Chem. Soc. 2005, 127, 3288-3289; (b) U. Mayerhöfer, B. Fimmel, F. Wüthner, Angew. Chem. Int. Ed. 2012, 51, 164-167; (c) J. M. Baumes, J. J. Gassensmith, J. Giblin, J.-J. Lee, A. G. White, W. J. Culligan, W. M. Leevy, M. Kuno, B. D. Smith, Nature Chem. 2010, 2, 1025-1030.
9. (a) S. Cohen, J. R. Lacher, J. D. Park, J. Am. Chem. Soc. 1959, 81, 3480-3481; (b) J. D. Park, S. Cohen, J. R. Lacher, J. Am. Chem. Soc. 1962, 84, 2919-2922; (c) S. Cohen, S. G. Cohen, J. Am. Chem. Soc. 1966, 88, 1533-1536.
10. (a) M. B. Onaran, A. B. Comeau, C. T. Seto, J. Org. Chem. 2005, 70, 10792-10803; (b) Y. Xu, N. Yamamoto, D. I. Ruiz, D. S. Kubitz, K. D. Janda, Bioorg. Med. Chem. Lett. 2005, 15, 19, 4304-4307; (c) A. Tevyashova, F. Sztaricskai, G. Batta, P. Herczegh, A. Jeney, Bioorg. Med. Chem. Lett. 2004, 14, 18, 4783-4789; (d) J. R. Porter, S. C. Archibald, K. Childs, D. Critchley, J. C. Head, J. M. Linsley, T. A. Parton, M. K. Robinson, A. Shock, R. J. Taylor, G. J. Warrellow, R. P. Alexander, B. Langham, Bioorg. Med. Chem. Lett. 2002, 12, 7, 1051-1054.
11. J. Xie, A. B. Comeau, C. T. Seto, Org. Lett. 2004, 6, 83-86.
12. (a) A. T. Blomquist, E. A. L. Lancette, J. Am. Chem. Soc. 1961, 83, 1387-1391; (b) S. Cohen, S. G. Cohen, J. Am. Chem. Soc. 1966, 88, 1533-1536.

13. (a) L. Beverina, P. Salice, Eur. J. Org. Chem. 2010, 1207–1225; (b) U. Mayerhöfer, M. Gänger, M. Stolte, B. Fimmel, F. Wüthner, Chem. Eur. J. 2013, 19, 218–232; (c) K.-Y. Law, B. F. Court, Can. J. Chem. **1986**, 64, 2267–2273; (d) A. Treibs, K. Jacob, Liebigs Ann. Chem. **1966**, 699, 153–167.
14. D. Yang, Z. Guan, L. Yang, Y. Huang, Q. Wei, Z. Lu, J. Yu, Solar Energy Materials & Solar Cells, **2012**, 105, 220–228.
15. R. R. Avirah, K. Jyothish, C. H. Suresh, E. Suresh, D. Ramaiah. Chem. Commun. **2011**, 47, 12822–12824.
16. E. Ronchi, R. Ruffo, S. A. Rizzato, Albinati, L. Beverina, G. A. Pagani, Org. Lett. **2011**, 13, 3166–3169.
17. Gaussian 09, Revision A02, M. J. Frisch, G. W. Trucks, H. B. Schlegel, G. E. Scuseria, M. A. Robb, J. R. Cheeseman, G. Scalmani, V. Barone, B. Mennucci, G. A. Petersson, H. Nakatsuji, M. Caricato, X. Li, H. P. Hratchian, A. F. Izmaylov, J. Bloino, G. Zheng, J. L. Sonnenberg, M. Hada, M. Ehara, K. Toyota, R. Fukuda, J. Hasegawa, M. Ishida, T. Nakajima, Y. Honda, O. Kitao, H. Nakai, T. Vreven, J. A. Montgomery, Jr., J. E. Peralta, F. Ogliaro, M. Bearpark, J. J. Heyd, E. Brothers, K. N. Kudin, V. N. Staroverov, T. Keith, R. Kobayashi, J. Normand, K. Raghavachari, A. Rendell, J. C. Burant, S. S. Iyengar, J. Tomasi, M. Cossi, N. Rega, J. M. Millam, M. Klene, J. E. Knox, J. B. Cross, V. Bakken, C. Adamo, J. Jaramillo, R. Gomperts, R. E. Stratmann, O. Yazyev, A. J. Austin, R. Cammi, C. Pomelli, J. W. Ochterski, R. L. Martin, K. Morokuma, V. G. Zakrzewski, G. A. Voth, P. Salvador, J. J. Dannenberg, S. Dapprich, A. D. Daniels, O. Farkas, J. B. Foresman, J. V. Ortiz, J. Cioslowski and D. J. Fox, Gaussian 09, Wallingford CT, **2009**.
18. (a) A. D. Becke, J. Chem. Phys. **1993**, 98, 5648. (b) C. Lee, W. Yang, R. Parr, Phys. Rev. B **1988**, 37, 785.
19. J. W. Mciver, Acc. Chem. Res. 1974, 7, 72.
20. (a) M. Cossi, V. Barone, R. Cammi, Tomasi, J. Chem. Phys. Lett. **1996**, 255, 327. (b) M. T. Cancès, V. Mennucci, J. Tomasi, J. Chem. Phys. **1997**, 107, 3032. (c) V. Barone, M. Cossi, J. Tomasi, J. Comput. Chem. **1998**, 19, 404.
21. K. F. Ansari, C. Lal, European Journal of Medicinal Chemistry. 2009, 44, 4028–4033.
22. Z. Shi, J. L. Qiang, Z. Li, Chinese Science Bulletin. 2001, 46, 393–395.
23. N. A. Zakhavova, B. A. Poral-Koshlts, L. S. E-Fros, Zhur Obshei Khim, 1953, 23, 1225–1230.



