

The Rabe amination after a century: direct addition of *N*-heterocycles to carbonyl compounds†‡

Daniele M. Scarpino Schietroma, Mattia R. Monaco, Valerio Bucalossi, Philipp E. Walter, Patrizia Gentili and Marco Bella\*

Received 21st March 2012, Accepted 10th May 2012

DOI: 10.1039/c2ob25595b

A catalytic version of the Rabe electrophilic amination is presented. This kind of reaction was originally employed in 1918 in a key step for the conversion of quinotoxine to quinine. Ketones and  $\alpha$ -substituted aldehydes give the corresponding  $\alpha$ -aminated carbonyl compounds in moderate yield.  $\alpha,\alpha$ -Unsubstituted aldehydes give rise to amino ketones *via* a novel rearrangement.

During the past ten years, the electrophilic amination reaction of carbonyl compounds<sup>1</sup> witnessed enormous advancement. In particular, one of the most widely exploited strategies has been the introduction of a nitrogen moiety *via* the addition of carbonyl compounds to diazodicarboxylates (carbamate-protected diimides), thanks to the pivotal contributions of Evans, List and Jørgensen and co-workers.<sup>2</sup>

The adducts obtained can be converted to biologically and pharmaceutically relevant molecules, such as  $\alpha$ -amino acids and  $\beta$ -amino alcohols, and have also been employed in the industrial scale preparation of some pharmaceutically relevant molecules.<sup>3</sup> Whilst these reactions present generally good yields and, in a number of cases, exceptional enantioselectivity, they are not as effective in terms of atom economy. The cleavage of the N–N bond and the carbamate moiety deprotection is not always straightforward.<sup>2b–c</sup>

Amino acids and amino alcohols presenting an *N*-heterocycle moiety are important molecules employed as constrained peptidomimetics and analogues of epinephrine-type drugs.<sup>4</sup>

An attractive complementary strategy to prepare these classes of compounds could be the direct introduction of a heterocyclic moiety in the  $\alpha$ -position of carbonyl compounds, *via* the reactive electrophiles *N*-halo amines. These reactive species have sporadically been employed in organic synthesis. Their usage can be traced back to the beginning of the twentieth century in the papers of Rabe.<sup>5</sup> In particular, *N*-bromoquinotoxine was one of

the crucial intermediates in Rabe's conversion of quinotoxine to quinine in 1918.<sup>6</sup> This transformation was later exploited by Woodward to claim the first formal synthesis of this natural substance,<sup>7</sup> which has generated significant debate during the recent years. The electrophilic amination step of the Rabe synthesis raised concern among scientists not fully convinced that the authors actually prepared quinine in 1932.<sup>8</sup> In 2001, Stork reported the first “non-formal” synthesis of quinine,<sup>9</sup> casting the doubt that this could indeed be its first total synthesis at all, since not enough experimental detail was provided in the Woodward synthesis. One of his main concerns was the transformation of quinotoxine into quinine.<sup>10</sup> In 2007, Smith and Williams repeated and verified the Rabe synthesis, employing the original conditions of that time, to specifically address this issue.<sup>11</sup> Regardless of the final outcome of this fascinating scientific debate, the amination reaction of carbonyl compounds is an intriguing transformation, which could surely be revisited in modern (asymmetric) synthesis.<sup>12</sup>

We decided to investigate the addition of simply carbonyl compounds to *N*-heterocycles as electrophiles more in detail, since it could also be an effective strategy to access valuable products. We were also interested to test if the “modern” organocatalytic activation mode of carbonyl compounds *via* enamine<sup>13</sup> would allow us to obtain the desired products under transition metal-free and mild conditions and, possibly, in an asymmetric way.


Initially, we tested the addition of acetone **1a** to the cyclic six-membered chlorinated heterocycle **2a**, prepared *via* the addition of NCS to piperidine.<sup>14–15</sup> Interestingly, with the combination of a catalytic amount of L-proline **1** and excess of triethylamine, the reaction proceeded smoothly at rt, giving the desired product **3a** in 63% yield after 24 h (Table 1, entry 1).

The enhanced reactivity of proline-derived enamines in the presence of bases has been documented by us<sup>16a–b</sup> and other authors.<sup>16c–e</sup> Encouraged by this result, we decided to extend the scope of the reaction to other substrates. We selected three classes of ketones: linear ketones **1b–d**, hindered ketones **1e–f** and cyclic ketones **1g–i**. The reaction proceeded in moderate yield (45–58%) with ketones **1b–d** (Table 1, entries 2–4). It should be noticed that the *N*-chloro amines **2a–b** employed as reactants as well as compounds **3** are thermally unstable. Therefore, this can affect the yield of these transformations. The

Dipartimento di Chimica, “Sapienza” Università di Roma, P.le Aldo Moro 5, 00185 Roma, Italy. E-mail: marco.bella@uniroma1.it

† The authors wish to dedicate this paper to Professor Alfredo Ricci, University of Bologna, for his pioneering and inspiring contribution for the development of amination reactions.

‡ Electronic supplementary information (ESI) available: Experimental procedures and compounds characterization. See DOI: 10.1039/c2ob25595b

**Table 1** Organocatalyzed addition of ketones **1a–i** to *N*-chloro heterocycles **2a–b**

| Entry <sup>a</sup> | Ketone <b>1</b> | Chloro amine ( <i>n</i> ), <b>2</b> | Amino ketone R <sup>1</sup> , R <sup>2</sup> , <b>3</b>                       | Yield <sup>b</sup> (%) |
|--------------------|-----------------|-------------------------------------|---|------------------------|
| 1                  | <b>1a</b>       | (2), <b>2a</b>                      | Me, H, <b>3a</b>  | 63                     |
| 2                  | <b>1b</b>       | (1), <b>2b</b>                      | Me, Me, <b>3b</b> Et, H, <b>3b'</b> <b>3b/3b'</b> = 1.4 : 1 <sup>c</sup>      | 58                     |
| 3                  | <b>1c</b>       | (1), <b>2b</b>                      | Et, Me, <b>3c</b>   | 35                     |
| 4                  | <b>1d</b>       | (1), <b>2b</b>                      | Me, <i>n</i> -Pr, <b>3d</b> <i>n</i> -Bu, H, <b>3d'</b> <b>3d/3d'</b> = 1 : 1 | 45                     |
| 5                  | <b>1e</b>       | (1), <b>2b</b>                      | <i>i</i> -Pr, H, <b>3e</b>  | 25                     |
| 6                  | <b>1f</b>       | (1), <b>2b</b>                      | Ph, Me, <b>3f</b>   | 22                     |
| 7                  | <b>1g</b>       | (2), <b>2a</b>                      | -(CH <sub>2</sub> ) <sub>3</sub> -, <b>3g</b>                                 | 38                     |
| 8                  | <b>1h</b>       | (1), <b>2b</b>                      | -(CH <sub>2</sub> ) <sub>4</sub> -, <b>3h</b>                                 | 36                     |
| 9                  | <b>1i</b>       | (1), <b>2b</b>                      | -(CH <sub>2</sub> ) <sub>5</sub> -, <b>3i</b>                                 | 25                     |

<sup>a</sup> Reactions are run employing 200 mg of *N*-heterocyclic amines **2a–b**, 1 mL of ketones **1a–i**, 0.3 equiv. of L-proline **1** and 2 equiv. of Et<sub>3</sub>N; reaction times: from 1 to 26 d (see ESI† for details). <sup>b</sup> Isolated yield of **3a–i** determined after purification *via* FC. <sup>c</sup> Determined by <sup>1</sup>H-NMR.

reported values of reaction times represent for each compound a compromise between conversion and yield. The transformations were left at rt for prolonged reaction times and the temperature has not been raised due to safety concerns.<sup>15</sup> Less satisfactory results were obtained with bulky ketones **1e** and **1f** (25% and 22% respectively, entries 5–6). In these examples, the lower yield is also due to the difficulty of forming the enamine intermediate on such hindered ketones. Finally, we tested cyclic ketones **1g–i** and we were also able to isolate the desired adducts (entries 7–9). In general, most of the examples (entries 6,8,9) were conducted with *N*-chloro pyrrolidine **2b** since it is much more reactive with respect to *N*-chloro piperidine **2a**.

Aldehyde substrates **4a–d** presented a different reactivity.  $\alpha,\alpha$ -Disubstituted aldehydes such as hydratropaldehyde **4a** afforded the desired adducts **5a–b** bearing a quaternary stereocenter in moderate to good yield (43–87%) together with the corresponding  $\alpha$ -chlorinated products **6a–b** (Table 2, entries 1–2). In this case, the *N*-chloro amines **2a–b** act also as a source of electrophilic chlorine.<sup>17</sup> Isobutyl aldehyde **4b** afforded exclusively the  $\alpha$ -aminated compound **5c** with no trace of the  $\alpha$ -chlorinated adduct **6c** (entry 3).<sup>18–19</sup>

Aldehydes **4c–d**, presenting a “free” CH<sub>2</sub> moiety in the  $\alpha$ -position, showed a different reactivity, giving rise to ketones **3e**, **3j–k** *via* a novel rearrangement (entries 4–6).

We rationalize the formation of ketones **3e**, **3j–k** *via* an intramolecular epoxide **9** formation (*via* adduct **8**) and subsequent


1,2 hydride migration (Scheme 1). The instability of enamine-derived epoxides has been documented.<sup>20</sup>

Clearly, developing an asymmetric catalytic version of the Rabe amination would be a challenging but important target. Any attempt to obtain a measurable enantiomeric excess we performed on cyclohexanone **1h** was met with failure. A possible explanation can be the configurational instability of tertiary stereocenters in the  $\alpha$ -position with respect to amines and ketones, as reported by Valentin *et al.*<sup>21</sup>

The only substrate that could give products in an enantiomerically enriched form is the  $\alpha$ -substituted aldehyde **4a** (Scheme 2).



Despite a vast screening of conditions and catalysts (see ESI†), we could not achieve any measurable enantioselectivity, except in the case when we conducted the reaction in presence of 4 Å molecular sieves. A similar effect has also been reported by Jacobsen and Yoon<sup>22</sup> and it could be the subject of further studies in our laboratory.

In conclusion, the rediscovery of the Rabe reaction allowed us to report our preliminary results on the direct functionalization of ketones and aldehydes with *N*-chloro amines in mild conditions. In particular, for aldehydes bearing a CH<sub>2</sub> moiety in the  $\alpha$ -position, we observed a new rearrangement leading to  $\alpha$ -amino ketones. Minimal but measurable enantioselectivity is observed with aldehydes bearing a quaternary stereocenter. Surely, with more extensive screening conditions, the reaction yield could be improved and the reaction time shortened. However, safety

**Table 2** Organocatalyzed addition of aldehydes **4a–d** to *N*-chloro heterocycles **2a–b**

| Entry <sup>a</sup> | Aldehyde <b>4</b> | Chloro amine ( <i>n</i> ), <b>2</b> | Product R <sup>1</sup> , R <sup>2</sup> , <b>5</b> , <b>6</b> or <b>3</b> | Yield <b>5</b> <sup>b</sup> (%) | Yield <b>6</b> <sup>b</sup> (%) | Yield <b>3</b> <sup>b</sup> (%) |
|--------------------|-------------------|-------------------------------------|---|---------------------------------|---------------------------------|---------------------------------|
| 1                  | <b>4a</b>         | <b>2</b> , <b>2a</b>                | Me, Ph, <b>5a–6a</b>  | 43                              | 25                              | —                               |
| 2                  | <b>4a</b>         | <b>1</b> , <b>2b</b>                | Me, Ph, <b>5b–6b</b>  | 87                              | 10                              | —                               |
| 3                  | <b>4b</b>         | <b>2</b> , <b>2a</b>                | Me, Me, <b>5c</b>   | 25                              | <5                              | —                               |
| 4                  | <b>4c</b>         | <b>1</b> , <b>2a</b>                | H, <i>i</i> -Pr, <b>3j</b>  | —                               | —                               | 46                              |
| 5                  | <b>4c</b>         | <b>1</b> , <b>2b</b>                | H, <i>i</i> -Pr, <b>3e</b>  | —                               | —                               | 17                              |
| 6                  | <b>4d</b>         | <b>2</b> , <b>2a</b>                | H, <i>n</i> -Bu, <b>3k</b>  | —                               | —                               | 38                              |

<sup>a</sup> Reactions are run employing 50 mg of aldehydes **4a–d**, 1.5 equiv. of *N*-heterocyclic amines **2a–b**, 0.30 equiv. of L-Pro **I** and 1.5 mL of DCM as the solvent; reaction times: 16–240 h (see ESI† for details). <sup>b</sup> Isolated yield of **5a–c**, **6a–c**, and **3e**, **3j–k** determined after purification *via* FC.

**Scheme 1** Proposed mechanism for the addition of *N*-chloro amines to  $\alpha,\alpha$ -unsubstituted aldehydes and their rearrangement to  $\alpha$ -amino ketones.**Scheme 2** Enantioselective addition of hydratropaldehyde **4a** to *N*-chloro pyrrolidine.

results can also be exploited to develop a more efficient amination reaction with maximum atom economy.

Financial support to this work was given by University of Roma “Sapienza” (“Progetto di Ateneo 2010 and 2011”). The authors thank Ms Susy Piovesana, for performing some preliminary experiments for this project.

## Notes and references

- 1 *Modern Amination Methods*, ed. A. Ricci, Wiley-VCH Verlag GmbH, Weinheim, Germany, 2007.

concerns prevented us from further investigations. We believe that the proof-of-concept contained in this work could be beneficial to the scientific community and in particular to those operating with unstable chloro amines. Hopefully, these preliminary

- 2 Leading references on the usage of diazodicarboxylates in organic synthesis. Enantioselective amination of amides: (a) D. A. Evans and S. G. Nelson, *J. Am. Chem. Soc.*, 1997, **119**, 6452. Amination of aldehydes: (b) A. Øgevig, K. Juhl, N. Kumaragurubaran, W. Zhuang and K. A. Jørgensen, *Angew. Chem., Int. Ed.*, 2002, **41**, 1790; (c) B. List, *J. Am. Chem. Soc.*, 2002, **124**, 5656. Ketones: (d) N. Kumaragurubaran, K. Juhl, W. Zhuang, A. Øgevig and K. A. Jørgensen, *J. Am. Chem. Soc.*, 2002, **124**, 6254.  $\alpha$ -Ketoesters: (e) K. Juhl and K. A. Jørgensen, *J. Am. Chem. Soc.*, 2002, **124**, 2420.  $\beta$ -Ketoesters: (f) M. Marigo, K. Juhl and K. A. Jørgensen, *Angew. Chem., Int. Ed.*, 2003, **42**, 1367; (g) S. Saaby, M. Bella and K. A. Jørgensen, *J. Am. Chem. Soc.*, 2004, **126**, 8120; (h) X. Liu, H. Li and L. Deng, *Org. Lett.*, 2005, **7**, 167. Aldehydes with the formation of quaternary stereocenters: (i) H. Vogt, S. Vanderheiden and S. Bräse, *Chem. Commun.*, 2003, 2448; (j) T. Baumann, H. Vogt and S. Bräse, *Eur. J. Org. Chem.*, 2007, 266.  $\alpha,\beta$ -Unsaturated aldehydes: (k) S. Bertelsen, M. Marigo, S. Brandes, P. Dinér and K. A. Jørgensen, *J. Am. Chem. Soc.*, 2006, **128**, 12973. Oxindoles: (l) T. Bui, M. Borregan and C. F. Barbas III, *J. Org. Chem.*, 2009, **74**, 8935; (m) T. Bui, N. R. Candeias and C. F. Barbas III, *J. Am. Chem. Soc.*, 2010, **132**, 5574. Naphthols: (n) S. Brandes, M. Bella, A. Kjærsgaard and K. A. Jørgensen, *Angew. Chem., Int. Ed.*, 2006, **45**, 1147. For selected reviews see: (o) R. O. Duthaler, *Angew. Chem., Int. Ed.*, 2003, **42**, 975; (p) J. M. Janey, *Angew. Chem., Int. Ed.*, 2005, **44**, 4292; (q) V. Nair, A. T. Bijur, S. C. Mathew and B. P. Babu, *Chem.–Asian J.*, 2008, **3**, 810; (r) C. Cativiela and M. Ordóñez, *Tetrahedron: Asymmetry*, 2009, **20**, 1; (s) T. Vilaivan and W. Bhanthumnavin, *Molecules*, 2010, **15**, 917; (t) A. Vallribera, R. M. S. Sebastián and A. Shafir, *Curr. Org. Chem.*, 2011, **15**, 1539.
- 3 R. M. De Figueiredo and M. Christmann, *Eur. J. Org. Chem.*, 2007, 2575.
- 4 W. Sneader, *Drug News Perspect.*, 2001, **14**, 539.
- 5 P. Rabe, *Ber. Dtsch. Chem. Ges.*, 1912, **45**, 2163.
- 6 P. Rabe and K. Kindler, *Ber. Dtsch. Chem. Ges.*, 1918, **51**, 466.
- 7 R. B. Woodward and W. E. Doering, *J. Am. Chem. Soc.*, 1945, **67**, 860.
- 8 P. Rabe, *Justus Liebigs Ann. Chem.*, 1932, **492**, 242.
- 9 G. Stork, D. Niu, A. Fujimoto, E. R. Koft, J. M. Balkovec, J. R. Tata and G. R. Duke, *J. Am. Chem. Soc.*, 2001, **123**, 3239.
- 10 For an overview on quinine synthesis see: J. I. Seeman, *Angew. Chem., Int. Ed.*, 2007, **46**, 1378.
- 11 A. C. Smith and R. M. Williams, *Angew. Chem., Int. Ed.*, 2008, **47**, 1736.
- 12 *N*-chloro amines have successfully been exploited by the research group of Jarvo, Miura and others for the direct amination of aromatic and heteroaromatic compounds. (a) T. J. Barker and E. R. Jarvo, *Angew. Chem., Int. Ed.*, 2011, **50**, 8325; (b) T. Kawano, K. Hirano, T. Satoh and M. Miura, *J. Am. Chem. Soc.*, 2010, **132**, 6900; (c) C. He, C. Chen, J. Cheng, C. Liu, W. Liu, Q. Li and A. Lei, *Angew. Chem., Int. Ed.*, 2008, **47**, 6414; (d) T. J. Barker and E. R. Jarvo, *J. Am. Chem. Soc.*, 2009, **131**, 15598; (e) T. Hatakeyama, Y. Yoshimoto, S. K. Ghorai and M. Nakamura, *Org. Lett.*, 2010, **12**, 1516.
- 13 (a) A. Berkessel and H. Gröger, *Asymmetric Organocatalysis: From Biomimetic Concepts to Applications in Asymmetric Synthesis*, Wiley-VCH, Weinheim, 2005; (b) J. Seayad and B. List, *Org. Biomol. Chem.*, 2005, **3**, 719; (c) P. Melchiorre, M. Marigo, A. Carlone and G. Bartoli, *Angew. Chem., Int. Ed.*, 2008, **47**, 6138.
- 14 For the preparation of *N*-chlorinated amines see: J. Quick and R. Oterson, *Synthesis*, 1976, 745.
- 15 CAUTION: twice during this research project we witnessed the violent decomposition of *N*-chloro pyrrolidine **2b** with generation of heat if kept neat at a temperature higher than 10 °C. In one instance, a sealed vial of *N*-chloro pyrrolidine **2b** exploded once removed from the freezer. *N*-chloro piperidine **2a** is less reactive and more stable. Extreme attention should be paid when handling *N*-chlorinated amines and during their preparation warming should be absolutely avoided.
- 16 (a) M. Bella, D. M. Scarpino Schietroma, P. P. Cusella, T. Gasperi and V. Visca, *Chem. Commun.*, 2009, 597; (b) D. M. Scarpino Schietroma, M. R. Monaco, V. Visca, S. Insogna, J. Overgaard and M. Bella, *Adv. Synth. Catal.*, 2011, **53**, 2648; (c) D. G. Blackmond, A. Moran, M. Hughes and A. Armstrong, *J. Am. Chem. Soc.*, 2010, **132**, 7598; (d) J. E. Hein, A. Armstrong and D. G. Blackmond, *Org. Lett.*, 2011, **13**, 4300; (e) M. B. Schmid, K. Zeitler and R. M. Gschwind, *Chem.–Eur. J.*, 2012, **18**, 3362.
- 17 When compound **6a** was treated with a large excess of piperidine, it gave in full conversion and moderate isolated yield the corresponding aminated compound **5a**, possibly via an  $S_N2$  reaction pathway.
- 18 For the electrophilic chlorination of aldehydes see: (a) N. Halland, A. Branton, S. Bachmann, M. Marigo and K. A. Jørgensen, *J. Am. Chem. Soc.*, 2004, **126**, 4790; (b) M. P. Brochu, S. P. Brown and D. W. C. MacMillan, *J. Am. Chem. Soc.*, 2004, **126**, 4108.
- 19 This experiment excludes the  $\alpha$ -aminated compounds **5a–c** forming exclusively via an  $S_N2$  mechanism on the corresponding chlorinated adducts (ref. 17), rather suggesting that two independent reaction pathways give rise to the formation of adducts **5** and **6**.
- 20 W. Adam, M. Ahrweiler, K. Paulini, H. U. Reißig and V. Vörckel, *Chem. Ber.*, 1992, **125**, 2719. An alternative mechanistic pathway could be a base-mediated quasi-Favorskii rearrangement of the chlorocarbonyl intermediate **6**, as suggested by one of the referees.
- 21 S. Bozzini, F. Felluga, G. Nardin, A. Pizzioli, G. Pitacco and E. Valentin, *J. Chem. Soc., Perkin Trans. 1*, 1996, 1961.
- 22 T. P. Yoon and E. N. Jacobsen, *Angew. Chem., Int. Ed.*, 2005, **44**, 466.