ChemComm

COMMUNICATION

Check for updates

Cite this: Chem. Commun., 2019, 55, 13303

Received 1st October 2019, Accepted 11th October 2019

DOI: 10.1039/c9cc07692a

rsc.li/chemcomm

Lewis acid-catalyzed synthesis of silafluorene derivatives from biphenyls and dihydrosilanes *via* a double sila-Friedel–Crafts reaction[†]

R^{3-Si-H}

(c) Intramolecular C-H silylation

b3

R^{3∽ Ŋi-}

R³

Yafang Dong,^a Yuta Takata,^a Yusuke Yoshigoe,^a Kohei Sekine^{ab} and Yoichiro Kuninobu *^{ab}

The synthesis of silafluorene derivatives from aminobiphenyl compounds and dihydrosilanes *via* a double sila-Friedel–Crafts reaction using a borane catalyst has been achieved. This method is applicable to the synthesis of a variety of silafluorene derivatives, such as multisubstituted silafluorenes, spirosilabifluorenes, and silicon-bridged terphenyl compounds, which are not readily obtained using conventional synthetic methods. In addition, we have demonstrated the transformation of the amino groups in these silafluorene derivatives into other substituents.

Silafluorene (dibenzosilole) derivatives are one of the most important compounds applied as organic functional materials, such as organic electroluminescent compounds,¹ light emitting materials,² field effect transistors,³ and photovoltaics.⁴ Therefore, several useful methods for the synthesis of silafluorene derivatives have been developed. As the conventional method, dilithiation of 2,2'-dibromobiphenyls and successive transmetalation of dilithiated biphenyls with dichlorosilanes is well known (Scheme 1a).⁵ Although this method is useful, it has some drawbacks including: (1) multiple reaction steps are required to synthesize the target silafluorene derivatives; (2) it is difficult to synthesize silafluorene derivatives with functional group(s), which react with organolithium reagents; and (3) the synthesis of multisubstituted silafluorene derivatives is difficult because the preparation of their multisubstituted dibromobiphenyl precursors is troublesome. Hence, a straightforward synthetic method for the preparation of multisubstituted silafluorenes is still desirable.

Recently, the synthesis of silafluorene derivatives using an intramolecular sila-Friedel–Crafts reaction of biphenylhydrosilanes has been reported (Scheme 1b).^{6–8} In addition, we have successfully

(a) Transmetalation of dilithiated biphenyls with dichlorosilanes



я^{́3}

R³

R cat. $B(C_6F_5)_3$

(d) This work: Double sila-Friedel-Crafts reaction

cat. RhCl(PPh₃)₃

Scheme 1 Several synthetic methods of silafluorene derivatives.

prepared a variety of silafluorene derivatives *via* a rhodiumcatalyzed intramolecular C–H silylation of biphenylhydrosilanes (Scheme 1c).^{9–11} Several transition metal-catalyzed synthetic methods to prepare silafluorenes have also been reported by other research groups.¹²

Based on the above-mentioned synthetic strategies, the reaction between biphenyl compounds and dihydrosilanes is a more ideal and promising approach to synthesize silafluorene derivatives. Recently, Ingleson^{7b} and Hou¹³ independently reported an effective borane-catalyzed C–H silylation of aromatic compounds *via* intra- and intermolecular sila-Friedel–Crafts reactions.¹⁴ We then considered that the synthesis of silafluorene derivatives was feasible based on a double sila-Friedel–Crafts reaction. Although

View Article Online

^a Department of Molecular and Material Sciences Interdisciplinary Graduate School of Engineering Sciences Kyushu University, 6-1 Kasugakoen, Kasuga-shi, Fukuoka 816-8580, Japan

^b Institute for Materials Chemistry and Engineering Kyushu University, 6-1 Kasugakoen, Kasuga-shi, Fukuoka 816-8580, Japan. E-mail: kuninobu@cm.kyushu-u.ac.jp

[†] Electronic supplementary information (ESI) available. CCDC 1950091 (3b). For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/ c9cc07692a

a ruthenium-catalyzed reaction between 2-phenylindoles and dihydrosilanes *via* sequential C–H silylation was reported, there is no example of the synthesis of silafluorenes.¹⁵ Herein, we report a direct borane-catalyzed synthesis of silafluorene derivatives from biphenyls and dihydrosilanes (Scheme 1d). This is the first example of the synthesis of silafluorene derivatives from biphenyls and dihydrosilanes *via* a double C–H silylation reaction.

We initially investigated the feasibility of our synthetic approach using diaminobiphenyl **1a** and dihydrodiphenylsilane **2a** as model substrates (Scheme 2). As a result, the double sila-Friedel–Crafts reaction of **1a** with **2a** in the presence of a catalytic amount of $B(C_6F_5)_3$ in chlorobenzene heated at 100 °C for 24 h afforded silafluorene **3a** in 73% yield. Ingleson reported that 2,6-dichloropyridine is crucial during the catalytic cycle of the intramolecular sila-Friedel–Crafts reaction of biphenylhydrosilanes. 2,6-Dichloropyridine is probably effective for the deprotonation of the silylated arenium cation.^{7b} Therefore, we investigated several additives as a base. The yield of **3a** was improved to 80% in the presence of 5.0 mol% 2,6-dichloropyridine and **3a** was obtained in 93% yield (87% isolated yield) using 7.5 mol% 2,6-lutidine.

We then investigated the scope of the biaryl compounds used in the reaction (Scheme 3). Biphenyls bearing acyclic or cyclic N,N-dialkylamino groups 1b-e gave their corresponding silafluorene derivatives (3b-e) in 81-95% yield.¹⁶ Silafluorenes 3f and 3g were formed using N-benzyl-N-methylamino and N-aryl-N-methylamino group-substituted biphenyls 1f and 1g, respectively. The previous synthetic methods used to prepare multisubstituted silafluorene derivatives have drawbacks due to the difficulties in preparing their multisubstituted starting materials. In contrast, trisubstituted silafluorene derivatives 3h-k were obtained in good to excellent yield from their corresponding biphenyls (1h-k), which can be readily synthesized. Notably, tetrasubstituted silafluorene 3l can be obtained in 52% yield. The reaction system is also applicable to monoaminobiphenyls 1m and 1n, which gave their corresponding silafluorenes 3m and 3n in 40 and 53% yields, respectively.

Subsequently, the scope of the dihydrosilane substrate (2) was investigated (Scheme 4). Diaryldihydrosilanes bearing electrondonating and electron-withdrawing groups gave silafluorene derivatives **30**, **3p**, and **3q** in 73, 87, and 91% yield, respectively. Silafluorene **3r** was obtained using di(2-naphthyl)silane (**2e**) in



Scheme 2 Synthesis of silafluorene derivative **3a** from diaminobiphenyl **1a** and dihydrodiphenylsilane **(2a)**.



high yield. Phenylmethylsilane and diethyldihydrosilane were also transformed into silafluorene derivatives **3s** and **3t** in high yield.

Spirosilabifluorenes are an attractive scaffold for organic optoelectronic materials due to their decreased HOMO–LUMO energy gap and fluorescent properties.^{9b,12a,17} It is notable that spirosilabifluorene derivative 5 was obtained from biphenyl **1d** and 9,9-dihydro-5-silafluorene (**4**) in 42% yield (Scheme 5).

Silicon-bridged terphenyl compounds have attracted a lot of attention due to their optoelectronic properties. Although silicon-bridged *para*-terphenyl compounds are well known,^{7a,9a,12a,17e,18}





examples of the synthesis of silicon-bridged *meta*-terphenyl compounds are quite rare.¹⁹ Hence, we investigated the synthesis of a silicon-bridged *meta*-terphenyl compound using our synthetic method. A quadruple sila-Friedel–Crafts reaction of *meta*-terphenyl derivative **6** with diphenylsilane (**2a**) afforded silicon-bridged *meta*terphenyl compound 7 in only 9% yield (Scheme 6). After the reaction, a mixture of biphenylhydrosilanes was observed using ¹H NMR spectroscopy. This result indicates that the intramolecular sila-Friedel–Crafts reaction on the central benzene ring did not proceed effectively. Therefore, after the intermolecular boranecatalyzed sila-Friedel–Crafts reaction, a successive rhodiumcatalyzed intramolecular C–H silylation⁹ was carried out. As a result, the yield of 7 was improved to 28% (Scheme 5).

Silafluorene **3b** was obtained in good yield, even on a gramscale (Scheme 7). Treating 1.00 g of biphenyl **1b** with 2.33 g of dihydrodiphenylsilane (**2a**) in the presence of catalytic amounts



Scheme 6 Synthesis of silicon-bridged terphenyl molecule 7.

of $B(C_6F_5)_3$ and 2,6-lutidine in chlorobenzene afforded 1.68 g of silafluorene **3b** (96% yield).

The amino groups on the silafluorene derivatives can be converted into other functional groups (Scheme 8).²⁰ Initially, the amino groups in **3b** were converted into their ammonium salts upon treatment with MeOTf and gave **8** in 97% yield. Ammonium salt **8** was then treated with a Grignard reagent (PhMgBr) in the presence of a palladium catalyst (PdCl₂(PPh₃)₂) to give the cross-coupling product (9) in 96% yield. Other Grignard reagents including 4-(MeO)C₆H₄MgBr, 4-FC₆H₄MgBr and (CH₂==CH-CH₂)MgBr also gave their corresponding silafluorenes (**10–12**) in moderate to good yield.

In summary, we have successfully prepared a variety of silafluorene derivatives from electron-rich biaryls and dihydrosilanes using a double sila-Friedel–Crafts reaction. The silafluorene derivatives were obtained in good to excellent yield, even on a gram-scale. Multisubstituted silafluorene derivatives



Scheme 7 Gram-scale synthesis of silafluorene derivative 3b.



Scheme 8 Conversions of amino groups.

were afforded from their corresponding biphenyl derivatives, which can be readily prepared. This new method is also applicable to the synthesis of spirosilabifluorenes and siliconbridged terphenyl derivatives. In addition, we have demonstrated the transformation of the amino groups in the silafluorene derivatives into several other substituents. We hope that the developed reaction will become a useful and effective protocol to synthesize a variety of silafluorene derivatives.

This work was supported in part by JSPS KAKENHI Grant Numbers JP 17H03016 and 18H04656, Iketani Science and Technology Foundation, The Sumitomo Foundation, and A-STEP (VP30218088652) from JST.

Conflicts of interest

There are no conflicts to declare.

Notes and references

- (a) R.-F. Chen, Q.-L. Fan, S.-J. Liu, R. Zhu, K.-Y. Pu and W. Huang, Synth. Met., 2006, 156, 1161; (b) K. Mouri, A. Wakamiya, H. Yamada, T. Kajiwara and S. Yamaguchi, Org. Lett., 2007, 9, 93; (c) J. C. Sanchez, A. G. DiPasquale, A. L. Rheingold and W. C. Trogler, Chem. Mater., 2007, 19, 6459; (d) E. Wang, C. Li, W. Zhuang, J. Peng and Y. Cao, J. Mater. Chem., 2008, 18, 797; (e) J. C. Sanchez and W. C. Trogler, J. Mater. Chem., 2008, 18, 3143; (f) Y.-Q. Mo, X.-Y. Deng, X. Jiang and Q.-H. Cui, J. Polym. Sci., Part A: Polym. Chem., 2009, 47, 3286.
- 2 (a) S. Beaupre, P.-L. T. Boudreault and M. Leclerc, Adv. Mater., 2010, 22, E6; (b) J. Y. Corey, Adv. Organomet. Chem., 2011, 59, 1.
- 3 (a) H. Usta, G. Lu, A. Facchetti and T. J. Marks, J. Am. Chem. Soc., 2006, **128**, 9034; (b) G. Lu, H. Usta, C. Risko, L. Wang, A. Facchetti, M. A. Ratner and T. J. Marks, J. Am. Chem. Soc., 2008, **130**, 7670.
- 4 (a) J. Hou, H.-Y. Chen, S. Zhang, G. Li and Y. Yang, *J. Am. Chem. Soc.*, 2008, **130**, 16144; (b) Y. Zou, D. Gendron, R. Neagu-Plesu and M. Leclerc, *Macromolecules*, 2009, **42**, 6361.
- 5 H. Gilman and R. D. Gorsich, J. Am. Chem. Soc., 1955, 77, 6380.
- 6 For ruthenium-catalyzed reactions, see: (a) L. Omann and M. Oestreich, Angew. Chem., Int. Ed., 2015, 54, 10276; (b) S. Bähr, A. Simonneau, E. Irran and M. Oestreich, Organometallics, 2016, 35, 925.
- 7 An example of a stoichiometric reaction: (a) S. Furukawa, J. Kobayashi and T. Kawashima, *J. Am. Chem. Soc.*, 2009, 131, 14192. An example of a catalytic reaction: (b) L. D. Curless and M. J. Ingleson, *Organometallics*, 2014, 33, 7241.

- 8 For examples of the synthesis of silafluorene derivatives by intramolecular cyclization via silyl radicals, see: (a) D. Leifert and A. Studer, Org. Lett., 2015, 17, 386; (b) L. Xu, S. Zhang and P. Li, Org. Chem. Front., 2015, 2, 459.
- 9 (a) T. Ureshino, T. Yoshida, Y. Kuninobu and K. Takai, J. Am. Chem. Soc., 2010, 132, 14324; (b) Y. Kuninobu, K. Yamauchi, N. Tamura, T. Seiki and K. Takai, Angew. Chem., Int. Ed., 2013, 52, 1520.
- 10 M. Murata, M. Takizawa, H. Sasaki, Y. Kohari, H. Sakagami, T. Namikoshi and S. Watanabe, *Chem. Lett.*, 2016, **45**, 857.
- 11 S. C. Richter and M. Oestreich, *Trends Chem.*, DOI: 10.1016/ j.trechm.2019.07.003.
- 12 (a) T. Matsuda, S. Kadowaki, T. Goya and M. Murakami, Org. Lett., 2007,
 9, 133; (b) M. Shimizu, K. Mochida and T. Hiyama, Angew. Chem., Int. Ed., 2008, 47, 9760; (c) Y. Yabusaki, N. Ohshima, H. Kondo, T. Kusamoto, Y. Yamanoi and H. Nishihara, Chem. Eur. J., 2010, 16, 5581; (d) J. M. Breunig, P. Gupta, A. Das, S. Tussupbayev, M. Diefenbach, M. Bolte, M. Wagner, M. C. Holthausen and H.-W. Lerner, Chem. Asian J., 2014, 9, 3163.
- 13 Y. Ma, B. Wang, L. Zhang and Z. Hou, J. Am. Chem. Soc., 2016, 138, 3663.
- 14 For other C-H silylation via sila-Friedel-Crafts reaction, see;
 (a) S. Bähr and M. Oestreich, Angew. Chem., Int. Ed., 2017, 56, 52.
 For an example of Brønsted or Lewis acid-catalyzed sila-Friedel-Crafts reaction, see: (b) Q.-A. Chen, H. F. T. Klare and M. Oestreich, J. Am. Chem. Soc., 2016, 138, 7868; (c) Q. Yin, H. F. T. Klare and M. Oestreich, Angew. Chem., Int. Ed., 2016, 55, 3204.
- 15 L. Omann and M. Oestreich, Organometallics, 2017, 36, 767.
- 16 The structure of **3b** was determined by single crystal X-ray structure analysis. For the details, see the ESI.[†] CCDC 1950091 (**3b**).
- 17 For selected papers for the synthesis and optoelectronic properties of spirosilabifluorenes, see: (a) A. G. Russell, N. S. Spencer, D. Philp, B. M. Kariuki and J. S. Snaith, Organometallics, 2003, 22, 5589; (b) S. H. Lee, B.-B. Jang and Z. H. Kafafi, J. Am. Chem. Soc., 2005, 127, 9071; (c) H. Xiao, B. Leng and H. Tian, Polymer, 2005, 46, 5707; (d) T. Agou, M. D. Hossain and T. Kawashima, Chem. Eur. J., 2010, 16, 368; (e) S. Furakawa, J. Kobayashi and T. Kawashima, Dalton Trans., 2010, 39, 9329; (f) E. Pusztai, I. S. Toulokhonova, N. Temple, H. Albright, U. I. Zakai, S. Guo, I. A. Guzei, R. Hu and R. West, Organometallics, 2013, 32, 2529; (g) A. V. Zabula, A. Y. Rogachev, I. A. Guzei and R. West, Organometallics, 2013, 32, 3760; (h) M. Murai, Y. Takeuchi, K. Yamauchi, Y. Kuninobu and K. Takai, Chem. Eur. J., 2016, 22, 6048.
- 18 L. Li, J. Xiang and C. Xu, Org. Lett., 2007, 9, 4877.
- 19 (a) H. N. Shin, C. S. Kim, Y. J. Cho, H. J. Kwon, B. O. Kim, S. M. Kim and S. S. Yoon, WO 2010114243, Dow Advanced Display Materials, Ltd, 2010; (b) H. Yang, CN 105985367, EverDisplay Optronics (Shanghai) Limited, 2016.
- 20 J. T. Reeves, D. R. Fandrick, Z. Tan, J. J. Song, H. Lee, N. K. Yee and C. H. Senanayake, *Org. Lett.*, 2015, **14**, 4388.