

189. Structure Elucidation and Synthesis of Flavonol Acylglycosides. III¹⁾. The Synthesis of Tiliroside

by Barbara Vermes

Forschungsgruppe für Alkaloidchemie der Ungarischen Akademie der Wissenschaften,
Budapest, Gellert 4

and Vedanta Mohan Chari and Hildebert Wagner

Institut für Pharmazeutische Biologie der Universität München, Karlstr. 29, D-8000 München 2

(11.VI.81)

Summary


Naturally occurring kaempferol 3-*O*- β -D-(6''-*O*-coumaroyl)glucopyranoside (tiliroside) has been synthesized thereby confirming its structure. This is the first acylated flavonoid glycoside to be synthesized.

Introduction. – In the plant kingdom tiliroside seems to have the widest distribution among the flavonol acylglycosides hitherto found. It was first isolated by Hörhammer *et al.* [1] from the leaves and flowers of *Tilia argentea* and has been subsequently reported to occur in *Tilia cordata* [2], *Anaphalis contorta* [3], *Helianthemum ovatum* [4], *Pinus contorta* [5], *Thymelea hirsuta* [6], *Pteridium aquilinum* [7] and some other plants.


The first structure proposed for tiliroside was a 7-*p*-coumarate of kaempferol 3-*O*- β -D-glucoside (astragalin). Harborne [8] showed by UV. spectroscopic investigations that tiliroside was a 3-*O*-(*p*-coumaroyl)glucoside of kaempferol. The exact position of the *p*-coumaroyl residue was shown by us to be on C(6'') of the glucose moiety on the basis of ¹³C-NMR. spectroscopy. Thus tiliroside was assigned the structure kaempferol 3-*O*- β -D-(6''-*O*-*p*-coumaroyl)glucopyranoside [9] (1).

Meanwhile it was independently established that the compounds isolated from *Helianthemum ovatum* [4] and *Pteridium aquilinum* [7] were identical with tiliroside from *Tilia argentea*. The same structure was postulated for a *p*-coumarate of astragalin (tribuloside) isolated from *Tribulus terrestris* [10]. An actual comparison [4] of the samples of tiliroside and tribuloside however showed their non-identity. As a part of our program on the structure elucidation and synthesis of naturally occurring flavonol acylglycosides we undertook the synthesis of tiliroside.

¹⁾ Part II, see [9].



1

2 $R^1 = \text{CH}_2\text{C}_6\text{H}_5$; $R^2 = \text{H}$; $R^3 = \text{COCH}_2\text{Cl}$; $R^4 = \text{OAc}$ 3 $R^1 = R^2 = \text{H}$; $R^3 = \text{COCH}_2\text{Cl}$; $R^4 = \text{OAc}$ 4 $R^1 = R^2 = R^4 = \text{OAc}$; $R^3 = \text{COCH}_2\text{Cl}$ 5 $R^1 = R^2 = R^4 = \text{OAc}$; $R^3 = \text{H}$ 6 $R^1 = R^2 = R^4 = \text{OAc}$; $R^3 = p\text{-(O-Ac)coumaroyl}$

Results and discussion. - Condensation of 7,4'-di-*O*-benzylkaempferol [11] with 2,3,4-tri-*O*-acetyl-6-*O*-chloroacetyl- α -D-glucopyranosylbromide²⁾ in pyridine solution in presence of Ag_2CO_3 and subsequent chromatographic purification gave **2** in 54% yield. As expected, glycosidation gave the 3-*O*- β -D-glucosyl derivative [13]. Catalytic debenzoylation of **2** and subsequent careful acetylation resulted in the product **4**. The chloroacetyl group was not affected during this sequence. The selective removal of the *O*-chloroacetyl group was effected by treatment of **4** with thiourea in methanol at RT. for 60 h yielding the compound **5**.

For coupling **5** with *p*-(*O*-acetyl)coumaroyl chloride [15], pyridine/dichloromethane was preferred. These conditions were applied by *Chittenden & Buchanan* [16] for the partial benzylation of sugars. A long reaction time (96 h at RT.) and a large excess of the acid chloride was necessary. Higher temperature caused partial change of **5** presumably due to a C(4'') to C(6'') acetyl migration. This facile migration was even observed during attempts to purify **4** by column chromatography on silica gel. The reaction product **6** was identical with tiliroside heptaacetate [4] [7]. The crucial point of the synthesis was the removal of the acetyl groups in **6** leaving the *p*-coumaroyl moiety intact. *Zemplen* deacetylation [17] using catalytic amounts of NaOMe and the calculated quantity of MeOH was performed on 6-*O*-cinnamoylglucose tetraacetate as a model. This resulted in selective cleavage of the acetyl groups in presence of the cinnamoyl residue and the method was applied to tilirosideheptaacetate [4] [7]. Tiliroside was formed in a 67% overall yield along with minor amounts of partially deacetylated products and kaempferol 3-*O*- β -D-glucoside (astragalin). The purified synthetic tiliroside was identical with the natural product (mixed m.p., Rf, UV., IR., ^1H -NMR. and ^{13}C -NMR.) [9].

We are indebted to Dr. *L. Pijewska* and *J. Kameczki* for a sample of tiliroside, and Drs. *A. Neszmélyi*, *O. Seligmann* and *P. Kolonits* for NMR. spectra. Financial support enabling Dr. *B. Vermes* to stay as a visiting scientist at the Institut für Pharmazeutische Biologie, München, was kindly provided by the *Deutsche Forschungsgemeinschaft* on a contract with the *Institute for Cultural Relations*, Budapest.

²⁾ The acylated glucopyranosylbromide was synthesized according to the method of *Gagnaire & Vottero* [12] starting with the acylation of β -D-1,2,3,4-tetra-*O*-acetylglucopyranose with chloroacetyl chloride in ether/pyridine followed by bromination with HBr/AcOH.

Experimental part

General remarks. M.p. were determined on a Kofler block and are uncorrected. UV. spectra (nm) were run in a) EtOH, b) with NaOAc, c) with AlCl_3 , d) with NaOMe. IR. spectra (cm^{-1}) were recorded as KBr pellets; NMR. spectra (ppm, J in Hz) were recorded at 60 and 100 MHz for ^1H and 23.5 MHz for ^{13}C with TMS as internal standard. Column chromatography was performed on silica gel and TLC. on Merck-6F₂₅₄-plates. Solvent systems: A=toluene/EtOH 9:1; B=toluene/EtOH 9:2; C=toluene/ Me_2CO 5:4; D=EtOAc/MeOH/ H_2O 100:16.5:13.5.

1. *Synthesis of 7,4'-Dibenzyl-3,5-dihydroxyflavone-3-O- β -D-(2'',3'',4''-tri-O-acetyl-6''-O-chloroacetyl)glucopyranoside (2).* To a solution of 7,4'-O-dibenzyl kaempferol [11] (466 mg) in pyridine (10 ml), Drierite (800 mg), Ag_2CO_3 (230 mg) and *a*-D-2,3,4-tri-O-acetyl-6-O-chloroacetylglucopyranosyl bromide [12] (750 mg) were added at 0°. After stirring for 1.5 h in the dark another portion (250 mg) of the bromide was added and stirring continued for another 1.5 h. The mixture was diluted with CHCl_3 (80 ml), filtered, extracted with 3% aq. sulfuric acid and washed with sat. aq. NaHCO_3 -solution and water. After evaporation of the solvent the residue was purified by column chromatography (A, Rf 0.64) to yield the glucoside **2** (450 mg, 54%) as a colorless, amorphous powder (from EtOH/ Me_2CO 9:1), m.p. 84–88°. – UV. (EtOH): (a) 262, 325; (b) 262, 325; (c) 275, 297, 337, 398; (d) 284, 380. – IR.: 1610, 1640, 1745 (CO). – ^1H -NMR. (CDCl_3): 2.01, 2.03, 2.12 (s, 3 H each, 3 OAc(glucose)); 3.82 (s, 2 H, CH_2Cl); 4.08 (m, 3 H, $\text{H}-\text{C}(5'',6'',6'')$); 5.13 and 5.17 (s, 2 H each, 2 CH_2Ph); 5.18–5.40 (3 H, $\text{H}-\text{C}(2'',3'',4'')$); 5.62 (d, $J=8.5$, 1H, $\text{H}-\text{C}(1'')$); 6.46 (d, $J=2.0$, 1H, $\text{H}-\text{C}(6)$); 6.52 (d, $J=2.0$, 1H, $\text{H}-\text{C}(8)$); 7.08 (d, $J=8.5$, 2 H, $\text{H}-\text{C}(3',5')$); 7.45 (10 H, arom.); 8.01 (d, $J=8.5$, 2 H, $\text{H}-\text{C}(2',6')$); 12.5 (s, 1 H, $\text{HO}-\text{C}(5)$).

$\text{C}_{43}\text{H}_{39}\text{ClO}_{15}$ (831.2) Calc. C 62.13 H 4.73 Cl 4.27% Found C 62.52 H 4.44 Cl 4.25%

2. *Synthesis of 5,7,4'-Tri-O-acetyl-3-O- β -D-(2'',3'',4''-tri-O-acetyl-6''-O-chloroacetyl)glucopyranoside (4).* By catalytic hydrogenation **2** (450 mg) was debenzylated in EtOH to yield **3** (390 mg, 90%) as a colourless powder, Rf 0.49 (B).

$\text{C}_{29}\text{H}_{27}\text{ClO}_{15}$ (651.0) Calc. C 53.50 H 4.18 Cl 5.44% Found C 53.92 H 4.35 Cl 5.74%

Compound **3** (390 mg) was immediately acetylated with acetic anhydride (2 ml) in presence of pyridine (2 ml) and gave after the usual work-up a grey precipitate which was purified by column chromatography (C, Rf 0.5, 400 mg, 88%), white needles, m.p. 196–199° (EtOH). – UV. (EtOH): (a) 248, 306. – IR.: 1620, 1640, 1760 (CO). – ^1H -NMR. (CDCl_3): 2.01 (s, 6 H, 2 OAc(glucose)); 2.12 (s, 3 H, OAc(glucose)); 2.28 (s, 6 H, 2 OAc(flavone)); 2.42 (s, 3 H, OAc(flavone)); 3.95 (s, 2 H, CH_2Cl); 4.10 (m, 3 H, $\text{H}-\text{C}(5'',6'',6'')$); 4.87–5.07 (m, 3 H, $\text{H}-\text{C}(2'',3'',4'')$); 5.56 (d, $J=8.0$, 1H, $\text{H}-\text{C}(1'')$); 6.88 (d, $J=2$, 1H, $\text{H}-\text{C}(6)$); 7.15–7.40 (m, 3 H, $\text{H}-\text{C}(8,3',5')$); 8.1 (d, $J=8.0$, 2 H, $\text{H}-\text{C}(2',6')$).

$\text{C}_{35}\text{H}_{33}\text{ClO}_{18}$ (777.1) Calc. C 54.09 H 4.28 Cl 4.56% Found C 53.60 H 4.72 Cl 4.67%

3. *Synthesis of 5,7,4'-Tri-O-acetyl-3-O- β -D-(2'',3'',4''-tri-O-acetyl)glucopyranoside (5).* To a solution of **4** (100 mg) in methanol (18 ml), thiourea (100 mg) in MeOH (5 ml) was added. After standing at RT. for 60 h the solvent was evaporated and the residue was extracted with CHCl_3 , filtered and washed with water. Evaporation of solvent yielded a white microcrystalline material (80 mg, 90%), m.p. 118–125°; Rf 0.26 (C). This compound was pure enough for the next stage as attempts at further purification on silica gel or repeated crystallization caused partial degradation of the product. – UV. (EtOH): (a) 250, 310. – IR.: 1645, 1750 (CO).

$\text{C}_{33}\text{H}_{32}\text{O}_{17}$ (700.6) Calc. C 56.57 H 4.60% Found C 56.30 H 4.68%

4. *Synthesis of 3,5,7,4'-Tetrahydroxyflavone-3-O- β -D-(6''-O-p-coumaroylglucopyranoside)-heptaacetate (tilirosideheptaacetate) (6).* To a solution of **5** (140 mg) and *p*-(O-acetyl)coumaroyl chloride [15] (160 mg) in CH_2Cl_2 , pyridine (2 ml) was added at 0° and the mixture left at RT. for 96 h. During this period a further 160 mg *p*-(O-acetyl)coumaroyl chloride was added in 4 portions. The mixture was diluted with CH_2Cl_2 (30 ml), extracted with 3% aq. sulfuric acid and washed with sat. aq. NaHCO_3 -solution and water. After evaporation of the solvent the residue was chromatographed (C, Rf 0.39) to yield tilirosideheptaacetate (**6**) (110 mg, 62%) as colorless needles from EtOH/ CHCl_3 20:1, m.p. 185–188° ([4]: 187–189). – IR. (KBr): 1620, 1700, 1760 (CO) (superimposable with that of an authentic sample). – ^1H -NMR. (CDCl_3): 2.01 (s, 6 H, 2 OAc(glucose)); 2.13 (s, 3 H, OAc(glucose)); 2.20 (s, 3 H,

OAc); 2.33 (*s*, 6 H, 2 OAc); 2.45 (*s*, 3 H, OAc); 3.65 (*m*, 1 H, H-C(5'')); 4.10 (*m*, 2 H, H-C(6'', 6'')); 4.9-5.1 (*m*, 3 H, H-C(2'', 3'', 4'')); 5.6 (*d*, 1 H, H-C(1'')); 6.27 (*d*, *J* = 13, 1 H, -CH=); 6.80 (*d*, *J* = 2, 1 H, H-C(6)); 7.12 and 7.18 (*d*, *J* = 7, 2 H each, H-C(3', 5', 3'', 5'')); 7.24 (*d*, *J* = 2, 1 H, H-C(8)); 7.52 (*d*, *J* = 7, 2 H, H-C(2''', 6'')); 7.54 (*d*, *J* = 13, 1 H, -CH=); 8.01 (*d*, *J* = 7, 2 H, H-C(2', 6')).

C₄₄H₄₀O₂₀ (888.8) Calc. C 59.46 H 4.54% Found C 59.10 H 4.77%

5. *Synthesis of 3,5,6,4'-Tetrahydroxyflavone-3-O-β-D-(6''-O-p-coumaroyl)glucopyranoside (tiliroside)* (1). To a solution of **6** (180 mg) in CHCl₃ (10 ml), 1N NaOMe in MeOH (0.03 ml) was added and left at RT. for 2 h. After acidification with AcOH and evaporation of the solvent the residue was purified by chromatography (D, Rf 0.47) to yield 80 mg (67%) of tiliroside as pale yellow needles from MeOH/H₂O 3:2, m.p. 267-269° (undepressed on admixture with an authentic sample) ([4]: 269-271°, [1]: 247-256°); [α]_D²⁵ = +67.8 (*c* = 0.12, MeOH); [α]_D²⁵ = +69.9 (*c* = 0.1001, MeOH) [4]. - UV. (EtOH): (a) 265, 320; (b) 273, 320; (c) 275, 305, 400; (d) 275, 380. - ¹³C-NMR. (DMSO-d₆, kaempferol moiety): 177.5 (C(4)); 164.1 (C(7)); 161.2 (C(5)); 159.9 (C(4'')); 156.4, 156.5 (C(9,2)); 133.1 (C(3)); 129.9, 130.7 (C(2', 6')); 120.9 (C(1')); 115.1, 115.8 (C(3', 5')); 104.0 (C(10)); 98.8 (C(6)); 93.6 (C(8)). - ¹³C-NMR. (DMSO-d₆, *p*-coumarate moiety): 166.1 (C(9'')); 159.9 (C(4'')); 144.4 (C(7'')); 130.7, 129.9 (C(2'', 6'')); 125.1 (C(1'')); 115.1, 115.8 (C(3'', 5'')); 113.9 (C(8'')). - ¹³C-NMR. (DMSO-d₆, β-D-glucopyranoside moiety): 101.3 (C(1'')); 76.5 (C(3'')); 74.3, 74.2 (C(2'', 5'')); 70.2 (C(4'')); 63.0 (C(6'')).

C₃₀H₂₆O₁₃ (594.5) Calc. C 60.61 H 4.41% Found C 60.95 H 4.71%

REFERENCES

- [1] L. Hörhammer, L. Stich & H. Wagner, *Naturwissenschaften* 46, 358 (1959).
- [2] R. P. Paris & J. P. Theallet, *Ann. Pharm. Fr.* 19, 850 (1961).
- [3] J. H. Lin, J. M. Lin & F. C. Chem, *J. Chinese Chem. Soc.* 23, 57 (1976).
- [4] L. Pijewska & J. Kamecki, *Roczniki Chem.* 51, 291 (1977).
- [5] R. Higuchi & D. M. X. Donnelly, *Phytochemistry* 17, 787 (1978).
- [6] S. I. Ismail, *Fitotherapie* 1978, 156.
- [7] M. Kuroyanagi, M. Fukuoka, K. Yoshihira, S. Natori & K. Yamazaki, *Chem. Pharm. Bull.* 26, 3594 (1978).
- [8] J. B. Harborne, *Phytochemistry* 1, 151 (1961).
- [9] V. M. Chari, M. Jordan & H. Wagner, *Planta Medica* 34, 93 (1978).
- [10] S. P. Bhutani, S. S. Chibber & T. R. Seshadri, *Phytochemistry* 8, 299 (1969).
- [11] H. Wagner, H. Danninger, O. Seligmann, M. Nogradi, L. Farkas & N. Farnsworth, *Chem. Ber.* 103, 3678 (1970).
- [12] D. Y. Gagnaire & P. J. A. Vottero, *Carbohydr. Res.* 28, 165 (1973).
- [13] H. Wagner, in *Zechmeister's «Prog. Chem. Org. Nat. Prod.»*, ed. W. Herz, H. Griesebach & G. W. Kirby, vol. 31, pp. 155, Springer-Verlag, Wien 1974.
- [14] M. Bertolini & C. P. J. Glaudemans, *Carbohydr. Res.* 15, 263 (1970).
- [15] S. Ogawa, *Bull. Chem. Soc. Jpn* 2, 20 (1927).
- [16] G. J. F. Chittenden & J. G. Buchanan, *Carbohydrate Res.* 11, 379 (1969).
- [17] G. Zemplén, *Ber. Deutsch. Chem. Ges.* 59, 12, 58 (1926).