## COMMUNICATION

## Preparation and relevance of a cross-coupling product between sinapyl alcohol and sinapyl *p*-hydroxybenzoate

Fachuang Lu,<sup>*a,b*</sup> John Ralph,<sup>\**a,b*</sup> Kris Morreel,<sup>*c*</sup> Eric Messens<sup>*c*</sup> and Wout Boerjan<sup>*c*</sup>

- <sup>a</sup> US Dairy Forage Research Center, USDA-ARS, 1925 Linden Drive West, Madison, WI 53706, USA
- <sup>b</sup> Dept. Forestry, University of Wisconsin-Madison, Madison, WI 53706, USA.
  E-mail: jralph@wisc.edu; Fax: +1 608 264 5147; Tel: +1 608 264 5407
- <sup>c</sup> Dept. Plant Systems Biology, Flanders Interuniversity Institute for Biotechnology, Ghent University, B-9000 Gent, Belgium

Received 27th July 2004, Accepted 10th September 2004 First published as an Advance Article on the web 28th September 2004

Cross-coupling of sinapyl *p*-hydroxybenzoate and sinapyl alcohol produces an 8-8-cross-coupled product that is also detected in lignifying poplar tissues, implicating sinapyl *p*-hydroxybenzoate as a lignin precursor.

The primary lignin monomers are the three monolignols, pcoumaryl, coniferyl, and sinapyl alcohols (4-hydroxycinnamyl, 4-hydroxy-3-methoxycinnamyl, and 3,5-dimethoxy-4-hydroxycinnamyl alcohols). Lignins of many agriculturally important crops and woody plants have their sidechain primary alcohol (the so-called 9- or  $\gamma$ -OH) acetylated, p-hydroxybenzoylated or *p*-coumarovlated.<sup>1-3</sup> The biochemistry of such acylation remains unresolved and the genes associated with the presumably involved transferase enzymes are unknown. As genes controlling various processes and the functions of such acylation are sought in order to improve the utilization of plant resources, it has become important to resolve whether monolignols are first acylated to produce ester conjugates which are then incorporated by coupling and cross-coupling into lignin by the conventional free-radical coupling reactions, or whether acylation occurs *following* the monolignol radical coupling reactions or on the lignin polymer itself.<sup>4,5</sup> We have provided preliminary evidence (to be more fully documented this year, Lu and Ralph

Peroxidase

unpublished) that the high level of 9-acetylation in kenaf lignins arises, at least in significant part, from pre-acetylated sinapyl alcohol,<sup>6</sup> and have provided at least good circumstantial evidence for monolignol *p*-coumarates in grasses.<sup>7,8</sup>

During recent studies profiling phenolic metabolite differences between normal poplar plants and transgenics deficient in genes/enzymes on the lignin biosynthetic pathway,<sup>9,10</sup> we discovered a novel 8-8-coupling product **2ab** bearing a *p*-hydroxybenzoate substituent.<sup>9</sup> This product was readily detected in normal plants but decreased to below detectable levels in plants heavily down-regulated by caffeic acid *O*-methyltransferase (COMT). The structural proof of the crucial product is *via* the independent preparation of the compound here.

Coupling reactions involving monolignols strongly favor homo-dehydrodimerization; cross-coupling reactions can be difficult to achieve.<sup>2,11</sup> Nevertheless, peroxidase–H<sub>2</sub>O<sub>2</sub> oxidation of equimolar amounts of sinapyl *p*-hydroxybenzoate and sinapyl alcohol produced a satisfactory 25% isolated yield of the cross-coupled product **2ab**, Scheme 1.† The ratio of products **2aa**: **2ab**: **2bb** was essentially 1:2:1 suggesting that the coupling reactions were insensitive to the acylation of the 9-OH.

Sinapyl alcohol **1a** radical 8-8-coupling produces an intermediate *bis*-quinone methide **QMaa**, Scheme 1, which







View Article Online

rearomatizes to afford syringaresinol 2aa by internal trapping *via* the two 9-OH groups.<sup>13</sup><sup>‡</sup> When the 9-OH group is acylated, as in the "dimerization" of sinapyl *p*-hydroxybenzoate **1b**, however, it is obviously incapable of trapping the quinone methide. Consequently, one quinone methide rearomatizes by external water addition, as is typically seen following 8-O-4-dehydrodimerization.<sup>13</sup> At this point there is an internal OH capable of trapping the other quinone methide moiety. The resulting product is therefore not a dehydrodimer, but the product 2bb, with a molecular mass 16 units higher. The cross-coupling reaction between 1a and 1b also produces an intermediary bis-quinone methide, but one such moiety is internally trapped by the single 9-OH. Rearomatization of the other requires water addition, and produces product 2ab. Finding the cross-coupling product 2ab in plants is compelling evidence that sinapyl p-hydroxybenzoate must therefore be an authentic "monomer" in the lignin (or lignan) pathway; it is unlikely to have arisen from the dehydrodimer of sinapyl alcohol, 2aa.

Indeed product **2ab** was identified in the actively lignifying xylem tissue of 3 month old poplar.<sup>9</sup> The product synthesized here was shown to be identical by its UV and mass (including MS-MS) spectral data, and HPLC retention time (including by co-injection). Its detection in actively lignifying tissue suggests that it may be a compound destined for the lignin polymer. More importantly, it demonstrates that sinapyl *p*-hydroxybenzoate is the "monomer" in the coupling reaction *in planta*. A *p*-hydroxybenzoyl transferase is therefore implicated for acylating the monolignol **1a** to feed the acylated substrate **1b** into the lignification process in poplar, where the lignin contains *p*-hydroxybenzoylated units.<sup>15</sup> The product **2ab** has also been isolated previously, without comment, from *Salix sachalinensis*.<sup>16</sup>

One of the more intriguing and unexpected chemical aspects of the homo-coupling reaction using sinapyl p-hydroxybenzoate 1b in peroxidase-H<sub>2</sub>O<sub>2</sub> at pH 5 to independently generate compound 2bb was that the intermediary quinone methide QMbb was isolable and stable.§ A pale yellow precipitate formed that was difficult to dissolve in common organic solvents. NMR in DMSO-d<sub>6</sub> revealed it to be the quinone methide QMbb. Although certain syringyl and guaiacyl quinone methides were sufficiently stable in solution to allow NMR spectra to be recorded in 1983,17 they have generally been considered unstable, although crystalline bromo analogs have been long known.<sup>18</sup> QMbb in the solid state did not degrade after over 8 months in a freezer. Regrettably, it has not yet been possible to obtain sufficiently good crystals for an X-ray crystal structure. Extracting the entire reaction product into EtOAc and washing with saturated aqueous NH<sub>4</sub>Cl acidified with HCl produces mainly compound 2bb (along with the tetrahydronaphthalene 2'bb) demonstrating that compound 2bb derives from this quinone methide intermediate.

In conclusion, demonstration that sinapyl *p*-hydroxybenzoate **1b** is an authentic precursor of lignification augments the finding that sinapyl acetate is implicated in kenaf bast fiber lignification<sup>6</sup> and the evidence for sinapyl and coniferyl *p*-coumarate in maize (and other grasses).<sup>7,8</sup> The mechanism therefore appears to be a general one for the three types of natural lignin acylation observed in nature. Future research will be aimed at determining if low levels of structures resulting from incorporation of **2ab** and **2bb** can be detected in lignins. The ultimate goals will be to find the transferases and their genes, and to attempt to understand the role of lignin acylation in these plants.

## Acknowledgements

We gratefully acknowledge partial funding through the DOE Energy Biosciences program (#DE-AI02-00ER15067) and the USDA-CSREES National Research Initiatives (Improved Utilization of Wood and Wood Fiber #2001-02176).

## Notes and references

\* Synthesis of the cross-coupled product from sinapyl alcohol and sinapyl p-hydroxybenzoate: sinapyl p-hydroxybenzoate5 (700 mg, 2.12 mmol) and sinapyl alcohol<sup>12</sup> (460 mg, 2.19 mmol) were dissolved in acetone (100 ml) and added into phosphate buffer (20 mM sodium dihydrogen phosphate, pH 4.5, 400 ml) in 500 ml flask. Horseradish peroxidase (Sigma, Type II, 150-250 units mg<sup>-1</sup>, 8 mg) was added followed by addition of the hydrogen peroxide urea complex (200 mg, 2.13 mmol) dissolved in 10 ml water. This mixture was stirred for 1 h when TLC indicated complete starting material disappearance. The mixture was saturated with NH4Cl and acidified with dilute aqueous HCl (3%) to pH around 3–4 and extracted with EtOAc (300 ml  $\times$  2). The organic phase was dried over MgSO4 and filtered. The crude product (pale white foam) was obtained after evaporation of the organic phase. The cross-coupled compound 2ab (280 mg, 25% yield) was isolated by silica gel chromatography with cyclohexane-EtOAc, 1:1 as eluting solvent. UV and MS data have been reported separately.9 NMR (acetone-d<sub>6</sub>)  $\delta_{\rm H}$  4.96 (1H, d, J = 5.0 Hz, A7), 3.99 (1H, m, A8), 4.14 (2H, m, A9s), 3.78\* (6H, s, A3/5-OMe), 6.72 (2H, s, A2/6); 4.90 (1H, d, J = 6.3 Hz, B7), 2.55 (1H, m, B8), 4.41 (1H, m, B9a), 4.68 (1H, m, B9b), 3.74\* (6H, s, B3/5-OMe), 6.66 (2H, s. B2/6); 7.77 (2H, m, C2/6), 6.87 (2H, m, C3/5) (\* assignments may be interchanged);  $\delta_{\rm C}$  136.0 (A1), 104.53 (A2/6), 148.5 (A3/5), 135.9 (A4), 72.6 (A7), 48.4 (A8), 70.0 (A9); 134.5 (B1), 104.45 (B2/6), 148.5 (B3/5), 136.0 (B4), 85.3 (B7), 49.8 (B8), 64.0 (B9); 122.4 (C1), 132.5 (C2/6), 116.0 (C3/5), 162.6 (C4), 166.4 (C7). Small amounts of other products such as 8-O-4-dehydrodimers and aryltetralin 8-8-products were not further examined.

<sup>‡</sup> In reactions with sinapyl alcohol alone, the 8-*O*-4-dehydrodimer is also produced but typically at less than 9% yield.<sup>14</sup>

 $\$  **QMbb**: NMR (DMSO-d<sub>6</sub>)  $\delta_{\rm H}$  3.66 (3H, s, S3-OMe), 3.70 (3H, s, S5-OMe), 4.02 (1H, m, S8), 4.42 (2H, m, S9), 6.56 (1H, d, *J* = 10.3, S7), 6.56 (1H, d, *J* = 1.9, S6), 6.82 (2H, m, P3/5), 6.93 (1H, d, *J* = 1.9, S2), 7.82 (2H, m, P2/6);  $\delta_{\rm C}$  40.7 (S8), 55.55 (S5-OMe), 55.62 (S3-OMe), 65.9 (S9), 104.8 (S6), 112.9 (S2), 115.9 (P3/5), 121.1 (P1), 132.3 (P2/6), 133.5 (S1), 140.4 (S7), 152.2 (S5), 153.7 (S3), 163.3 (P4), 166.2 (P7), 175.29 (S4).

- 1 W. Boerjan, J. Ralph and M. Baucher, Lignin Biosynthesis, Annu. Rev. Plant Biol., 2003, 54, 519.
- 2 J. Ralph, K. Lundquist, G. Brunow, F. Lu, H. Kim, P. F. Schatz, J. M. Marita, R. D. Hatfield, S. A. Ralph, J. H. Christensen and W. Boerjan, Lignins: natural polymers from oxidative coupling of 4-hydroxyphenylpropanoids, *Phytochem. Rev.*, 2004, 3, in press.
- 3 J. C. del Rio, A. Gutierrez and A. T. Martinez, Identifying acetylated lignin units in non-wood fibers using pyrolysis-gas chromatography/mass spectrometry, *Rapid Commun. Mass Spectrom.*, 2004, **18**, 1181.
- 4 J. Ralph and F. Lu, The DFRC method for lignin analysis. Part 6. A modified method to determine acetate regiochemistry on native and isolated lignins, *J. Agric. Food Chem.*, 1998, **46**, 4616.
- 5 Y. Nakamura and T. Higuchi, Ester linkage of *p*-coumaric acid in bamboo lignin. II. Syntheses of coniferyl *p*-hydroxybenzoate and coniferyl *p*-coumarate as possible precursors of aromatic acid esters in lignin, *Cellul. Chem. Technol.*, 1978, **12**, 199.
- 6 F. Lu and J. Ralph, Preliminary evidence for sinapyl acetate as a lignin monomer in kenaf, *Chem. Commun.*, 2002, 90.
- 7 J. Ralph, R. D. Hatfield, S. Quideau, R. F. Helm, J. H. Grabber and H.-J. G. Jung, Pathway of *p*-coumaric acid incorporation into maize lignin as revealed by NMR, *J. Am. Chem. Soc.*, 1994, **116**, 9448.
- 8 F. Lu and J. Ralph, Detection and determination of *p*-coumaroylated units in lignins, *J. Agric. Food Chem.*, 1999, **47**, 1988.
- 9 K. Morreel, J. Ralph, H. Kim, F. Lu, G. Goeminne, S. A. Ralph, E. Messens and W. Boerjan, Profiling of oligolignols reveals monolignol coupling conditions in lignifying poplar xylem, *Plant Physiol.*, 2004, in press.
- 10 K. Morreel, J. Ralph, F. Lu, G. Goeminne, R. Busson, P. Herdewijn, J. L. Goeman, J. Van der Eycken, W. Boerjan and E. Messens, Phenolic profiling of COMT-deficient poplar reveals novel benzodioxane oligolignols, *Plant Physiol.*, 2004, in press.
- 11 K. Syrjanen and G. Brunow, Regioselectivity in lignin biosynthesis. The influence of dimerization and cross-coupling, *J. Chem. Soc., Perkin Trans. 1*, 2000, 183.
- 12 F. Lu and J. Ralph, Highly selective syntheses of coniferyl and sinapyl alcohols, *J. Agric. Food Chem.*, 1998, **46**, 1794.
- K. Freudenberg and A. C. Neish. Constitution and Biosynthesis of Lignin, Springer-Verlag, Berlin-Heidelberg-New York, 1968.
   M. Tanahashi, H. Takeuchi and T. Higuchi, Dehydrogenative
- 14 NI. Iananashi, H. Takeuchi and T. Higuchi, Dehydrogenative polymerization of 3,5-disubstituted *p*-coumaryl alcohols, *Wood Res.*, 1976, **61**, 44.
- 15 H. Meyermans, K. Morreel, C. Lapierre, B. Pollet, A. De Bruyn, R. Busson, P. Herdewijn, B. Devreese, J. Van Beeumen, J. M. Marita, J. Ralph, C. Chen, B. Burggraeve, M. Van Montagu, E. Messens

- 16 H. Lee, N. Watanabe, T. Sasaya and S. Ozawa, Extractives of short-rotation hardwood species. I. Phenolics of the wood of Salix sachalinensis Fr. Schm, *Mokuzai Gakkaishi*, 1993, **39**, 1409.
- 17 J. Ralph and B. R. Adams, Determination of the conformation and isomeric composition of lignin model quinone methides by NMR, J. Wood Chem. Technol., 1983, 3, 183.
- 18 T. Zinke and O. Hahn, Über die Einwirkung von Brom und von Chlor auf Phenole: Substitutionsprodukten, Pseudobromide and Pseudochloride; IX. Über die Einwirkung von Brom auf Isoeugenol, Justus Liebigs Ann. Chem., 1903, 329, 1.