This article was downloaded by: [University of Haifa Library] On: 23 June 2013, At: 05:15 Publisher: Taylor & Francis Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Journal of Asian Natural Products Research

Publication details, including instructions for authors and subscription information: http://www.tandfonline.com/loi/ganp20

Two new lignan glycosides from the seeds of Cuscuta chinensis

Xiang-Hui He $^{\rm a}$, Wen-Zhi Yang $^{\rm a}$, A-Hui Meng $^{\rm a}$, Wen-Ni He $^{\rm a}$, De-An Guo $^{\rm a}$ & Min Ye $^{\rm a}$

^a The State Key Laboratory of Natural and Biomimetic Drugs, School of Pharmaceutical Sciences, Peking University, Beijing, 100191, China Published online: 08 Nov 2010.

To cite this article: Xiang-Hui He , Wen-Zhi Yang , A-Hui Meng , Wen-Ni He , De-An Guo & Min Ye (2010): Two new lignan glycosides from the seeds of Cuscuta chinensis , Journal of Asian Natural Products Research, 12:11, 934-939

To link to this article: <u>http://dx.doi.org/10.1080/10286020.2010.506434</u>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: http://www.tandfonline.com/page/terms-and-conditions

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae, and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand, or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.



Two new lignan glycosides from the seeds of Cuscuta chinensis

Xiang-Hui He, Wen-Zhi Yang, A-Hui Meng, Wen-Ni He, De-An Guo and Min Ye*

The State Key Laboratory of Natural and Biomimetic Drugs, School of Pharmaceutical Sciences, Peking University, Beijing 100191, China

(Received 23 April 2010; final version received 1 July 2010)

Two new lignan glycosides, 2'-hydroxyl asarinin 2'-O- β -D-glucopyranoside (cuscutoside C, 1) and 2'-hydroxyl asarinin 2'-O- β -D-apiofuranosyl- $(1 \rightarrow 2)$ -[β -D-glucopyranosyl- $(1 \rightarrow 6)$]- β -D-glucopyranoside (cuscutoside D, 2), were isolated from the seeds of *Cuscuta chinensis* Lam., along with six known compounds, 2'-hydroxyl asarinin 2'-O- β -D-glucopyranosyl- $(1 \rightarrow 6)$ - β -D-glucopyranoside (3), 2'-hydroxyl asarinin 2'-O- β -D-apiofuranosyl- $(1 \rightarrow 2)$ - β -D-glucopyranoside (cuscutoside A, 4), kaempferol 3,7-di-O- β -D-glucopyranoside (5), 5-caffeoyl quinic acid (6), 4-caffeoyl quinic acid (7), and cinnamic acid (8). Their structures were elucidated on the basis of spectroscopic analyses including HR-ESI-MS, ESI-MS/MS, ¹H and ¹³C NMR, HSQC, HMBC, and TOCSY.

Keywords: Cuscuta chinensis Lam.; lignan; cuscutoside C; cuscutoside D

1. Introduction

TuSiZi, which is derived from the seeds of Cuscuta chinensis Lam., is a commonly used traditional Chinese medicine (TCM). It was first recorded in the ancient Chinese herbal classic ShenNong's Herbal around 2000 years ago. TuSiZi was used as a tonic by TCM practitioners for the treatment of deficiency in kidney and liver. Modern researches have shown that it can improve sexual function, prevent and treat cardiovascular diseases, regulate the body's immune system, treat osteoporosis, and prevent senescence [1-9]. All these activities were hitherto considered to be due to flavonoids [10]. Recently, our phytochemical screening of C. chinensis by liquid chromatography coupled with mass spectrometry discovered a series of lignans and quinic acid derivatives [11]. Although a number of lignans had been isolated from this plant [12-14], many compounds remain to be unknown. In order to find out the relationship between bioactivities and chemical constituents, we conducted the phytochemical studies on the seeds of C. chinensis, particularly aiming to obtain new bioactive lignans. As a result, two new furofuran-type lignan glycosides, cuscutoside C (1) and cuscutoside D (2) (Figure 1), were isolated from an ethanol extract, together with two known lignans (2'-hydroxyl asarinin 2'-O- β -D-glucopyranosyl- $(1 \rightarrow 6)$ - β -D-glucopyranoside (3) [13] and 2'-hydroxyl asarinin 2'-O- β -D-apiofuranosyl- $(1 \rightarrow 2)$ - β -D-glucopyranoside or cuscutoside A (4) [12]), one flavonoid (kaempferol 3,7-di-O- β -D-glucopyranoside (5) [15]), two quinic acid derivatives (5-caffeoyl quinic acid (6) and 4-caffeoyl quinic acid (7) [16]), and one phenolic acid (cinnamic acid, 8) [17]. Among them, 5, 6, 7, and 8 were reported from this plant for the first time. The structures of 1 and 2 were elucidated on the basis of spectroscopic analyses including

ISSN 1028-6020 print/ISSN 1477-2213 online © 2010 Taylor & Francis DOI: 10.1080/10286020.2010.506434 http://www.informaworld.com

^{*}Corresponding author. Email: yemin@bjmu.edu.cn



Figure 1. Structures of cuscutoside C (1) and D (2).

HR-ESI-MS, ESI-MS/MS, ¹H and ¹³C NMR, HSQC, HMBC, and TOCSY.

2. Results and discussion

Compound 1 was obtained as a white amorphous powder. The HR-ESI-MS spectrum showed the $[M + Na]^+$ ion at m/z 555.1481, thus its molecular formula was established as C₂₆H₂₈O₁₂. In the negative ESI-MS/MS spectrum, the $[M - H]^{-}$ ion at m/z 531 yielded a product ion at m/z 369 by losing 162 Da, suggesting that compound 1 contained one hexosyl residue. The ¹H NMR spectrum showed signals due to five aromatic protons in the region of $\delta_{\rm H}$ 6.81–6.91, signals due to four methylenedioxy protons in the region of $\delta_{\rm H}$ 5.90-6.10, and signals due to protons of ditetrahydrofurofuran ($\delta_{\rm H}$ 2.87 (m), 3.81 (dd, J = 3.4, 9.0 Hz), 4.01 (dd, J = 4.0,9.1 Hz), 4.14 (dd, J = 6.4, 6.6 Hz), 4.57 (d, J = 5.2 Hz), 5.08 (d, J = 4.2 Hz)). These signals were very similar to those of 2'hydroxyl asarinin-type furofuran lignans [12,13]. Indeed, several 2'-hydroxyl asarinin glycosides had been reported from the seeds of C. chinensis [12,13]. Thus, the aglycone of compound 1 was deduced to be 2'-hydroxyl asarinin. The ¹³C NMR spectrum also showed signals due to 2'-hydroxyl asarinin (Table 1). In addition, signals (δ_C 60.8, 69.9, 76.7, 77.1, 73.4, 101.9) due to a hexosyl residue were observed. By comparing with reported NMR spectral data [12] and by acid hydrolysis-TLC analysis, the hexose was identified as glucose. The coupling constant of the anomeric proton at $\delta_{\rm H}$ 4.71 (d, J = 7.3 Hz) revealed that the glucosyl residue should be oriented in the β configuration. The location of the sugar moiety was deduced to be at C-2' (δ_{C} 148.8), according to the HMBC correlation between H-1" ($\delta_{\rm H}$ 4.71) of Glc and C-2' of the aglycone (Figure 2). Based on the above analyses, the structure of 1 was elucidated as 2'-hydroxyl asarinin 2'-O-β-D-glucopyranoside, and was named cuscutoside C.

Compound **2** was obtained as a white amorphous powder. Based on the HR-ESI-MS data, its molecular formula was deduced as $C_{37}H_{46}O_{21}$. The ¹H and ¹³C NMR spectral data indicated that **2** possessed the same 2'-hydroxyl asarinin aglycone as **1**. The only difference was the sugar moiety. In the negative ESI-MS/MS spectrum, an $[M - H]^-$ ion at *m/z* 825 and



Figure 2. Key HMBC correlations of cuscutoside C (1) and D (2).

fragment ions $[M - H - hexose]^{-}$ at m/z663, $[M - H - hexose - pentose]^{-1}$ at m/z 531, and [M - H - 2hexose pentose]⁻ at m/z 369 were observed. It suggested that the sugar moiety of 2 should contain two hexosyl residues and one pentosyl residue. The exact sugar types and sequence were determined by detailed analyses of the 1D and 2D NMR spectra. Comparison of the carbon chemical shifts with reported ¹³C NMR spectral data [12,13] and TLC analysis after acid hydrolysis allowed the identification of sugar types as Glc and Api. Proton signals due to the sugar residues were assigned according to HSQC and TOCSY spectra (Table 1). In the HMBC spectrum, correlations between H-1" ($\delta_{\rm H}$ 5.00) and C-2' ($\delta_{\rm C}$ 148.1), between H-1^{'''} ($\delta_{\rm H}$ 4.15) and C-6" ($\delta_{\rm C}$ 69.0), and between H-1"" ($\delta_{\rm H}$ 5.35) and C-2" ($\delta_{\rm C}$ 78.8) indicated the attachment positions of sugars to sugars and sugars to the aglycone (Figure 2). From the J values of H-1" ($\delta_{\rm H}$ 5.00, d, $J = 8.0 \,\mathrm{Hz}$, H-1^{'''} (δ_{H} 4.15, d. J = 8.0 Hz), and H-1^{////} (δ_{H} 5.35, d, J = 4.0 Hz), the β -configurations of Glc and Api were suggested [18]. From the above evidences, the structure of 2 was elucidated as 2'-hydroxyl asarinin 2'-O- β -D-apiofuranosyl- $(1 \rightarrow 2)$ - $[\beta$ -D-glucopyranosyl- $(1 \rightarrow 6)$]- β -D-glucopyranoside, and was named cuscutoside D.

The structures of known compounds 3-8 were identified by comparing with the literature data.

3. Experimental

3.1 General experimental procedures

The optical rotations were measured on a Rudolph Research Analytical Autopol III automatic polarimeter. Melting point were measured with an uncorrected X-4A microscope melting point apparatus (CBIO Bioscience & Technologies Co. Ltd, Beijing, China). UV spectra were recorded on a TU1901 UV-vis spectrophotometer (Beijing Purkinje General Instrument Co. Ltd, Beijing, China). IR spectra were recorded on a Nicolet NEXUS-470 spectrometer (Madison, WI, USA). 1D and 2D NMR spectra were obtained on a Bruker Avance III 400 spectrometer. HR-ESI-MS spectra were obtained on a Bruker Apex IV FT-MS spectrometer (Billerica, MA, USA). ESI-MS/MS spectra were recorded on a Finnigan LCQ Advantage ion trap mass

$\begin{tabular}{ c c c c c c c c c c c c c c c c c c c$		1		2	
$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$		$\delta_{ m H}$	δ_{C}	$\delta_{ m H}$	δ_{C}
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	Aglycone				
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	C(1)		135.5		135.5
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	CH(2)	6.91 (s)	106.5	6.92 (s)	106.5
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	C(3)		146.5		146.5
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	C(4)		147.4		146.5
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	CH(5)	6.85 (overlapped)	107.9	6.86 (d, $J = 8.0$)	107.9
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	CH(6)	6.84 (overlapped)	119.3	6.83 (d, $J = 8.0$)	119.3
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	CH(7)	4.57 (d, $J = 5.2$)	84.3	4.54 (d, J = 4.0)	84.6
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	CH(8)	2.87 (m)	54.0	2.86 (m)	54.1
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	$CH_2(9)$	4.14 (dd, J = 6.6, 6.4) (Ha)	70.8	4.12 (m) (Ha)	70.7
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	aut	3.81 (dd, J = 9.0, 3.4) (He)	1015	3.80 (m) (He)	100 5
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	C(1')		124.5		123.7
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	C(2')	(04/)	148.8		148.1
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	CH(3')	6.84 (s)	98.4	6.92 (s)	97.7
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	C(4')		146.4		146.5
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	C(5')	(01)	141./	(70)	141.5
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	CH(0')	0.81(S)	104.6	0.78(s)	104.6
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	CH(7)	5.08 (d, J = 4.2)	81.1	5.05 (d, J = 4.0)	80.5
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	$CH(\delta')$	2.87 (m)	54.0 72.4	2.80 (m)	54.0
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	$CH_{2}(9')$	4.14 (dd, $J = 0.0, 0.4$) (Ha)	72.4	4.19 (m) (Ha)	12.5
$\begin{array}{cccccccccccccccccccccccccccccccccccc$		4.01 (dd, $J = 9.1, 4.0$) (He) 5.00 (a) 5.05 (d, $I = 11.4$)	100.9	5.95 (m) (He) 5.04 (d. $I = 8.0$) 5.00 (c)	100.9
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	0-CH ₂ -0	5.99 (s), 5.95 (d, $J = 11.4$)	100.8	5.94 (d, J = 8.0), 5.99 (s)	100.8
$\begin{array}{cccc} \mathrm{CH}(1'') & 4.71 & (d, J = 7.3) & 101.9 & 5.00 & (d, J = 8.0) & 98.4 \\ \mathrm{CH}(2'') & 3.21 & (\mathrm{m}) & 73.4 & 3.47 & (\mathrm{m}) & 78.8 \\ \mathrm{CH}(3'') & 3.29 & (\mathrm{m}) & 77.1 & 3.01 & (\mathrm{m}) & 77.0 \\ \mathrm{CH}(4'') & 3.14 & (\mathrm{m}) & 69.9 & 3.17 & (\mathrm{m}) & 70.1 \\ \mathrm{CH}(5'') & 3.21 & (\mathrm{m}) & 76.7 & 3.68 & (\mathrm{m}) & 75.1 \\ \mathrm{CH}_2(6'') & 3.43 & (\mathrm{m}), 3.72 & (d, J = 11.3) & 60.8 & 3.60 & (d, J = 4.0), 3.94 & (d, J = 4.0) & 69.0 \\ 6'' - O - Glc & & & & & & & & & & & & & & & & & & &$	2'-0-Glc				
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	CH(1")	4.71 (d, $J = 7.3$)	101.9	5.00 (d, $J = 8.0$)	98.4
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	CH(2")	3.21 (m)	73.4	3.47 (m)	78.8
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	CH(3")	3.29 (m)	77.1	3.01 (m)	77.0
$\begin{array}{cccc} \mathrm{CH}(5'') & 3.21 \ (\mathrm{m}) & 76.7 & 3.68 \ (\mathrm{m}) & 75.1 \\ \mathrm{CH}_2(6'') & 3.43 \ (\mathrm{m}), 3.72 \ (\mathrm{d}, J=11.3) & 60.8 & 3.60 \ (\mathrm{d}, J=4.0), 3.94 \ (\mathrm{d}, J=4.0) & 69.0 \\ 6''-O-Glc & & & & & \\ \mathrm{CH}(1''') & & 4.15 \ (\mathrm{d}, J=8.0) & 103.5 \\ \mathrm{CH}(2''') & & 2.95 \ (\mathrm{m}) & 73.5 \\ \mathrm{CH}(3''') & & 3.79 \ (\mathrm{m}) & 76.7 \\ \mathrm{CH}(4''') & & 3.02 \ (\mathrm{m}) & 70.1 \\ \mathrm{CH}(5''') & & 3.10 \ (\mathrm{m}) & 76.9 \\ \mathrm{CH}_2(6''') & & 3.64 \ (\mathrm{d}, J=8.0), 3.40 \ (\mathrm{m}) & 61.1 \\ 2''-O-Api & & \\ \mathrm{CH}(1''') & & 5.35 \ (\mathrm{d}, J=4.0) & 109.2 \\ \mathrm{CH}(2''') & & 3.80 \ (\mathrm{m}) & 76.2 \\ \mathrm{CH}(3''') & & 3.31 \ (\mathrm{d}, J=8.0) & 63.4 \\ \mathrm{CH}_2(5''') & & 3.57 \ (\mathrm{m}), 3.86 \ (\mathrm{m}) & 73.5 \\ \end{array}$	CH(4")	3.14 (m)	69.9	3.17 (m)	70.1
$\begin{array}{c} \mathrm{CH}_2(6'') & 3.43 \ \mathrm{(m)}, 3.72 \ \mathrm{(d,}\ J=11.3) & 60.8 & 3.60 \ \mathrm{(d,}\ J=4.0), 3.94 \ \mathrm{(d,}\ J=4.0) & 69.0 \\ 6''-O-Glc & & & & \\ \mathrm{CH}(1''') & & 4.15 \ \mathrm{(d,}\ J=8.0) & 103.5 \\ \mathrm{CH}(2''') & & 2.95 \ \mathrm{(m)} & 73.5 \\ \mathrm{CH}(3''') & & 3.79 \ \mathrm{(m)} & 76.7 \\ \mathrm{CH}(4''') & & 3.02 \ \mathrm{(m)} & 70.1 \\ \mathrm{CH}(5''') & & 3.10 \ \mathrm{(m)} & 76.9 \\ \mathrm{CH}_2(6''') & & 3.64 \ \mathrm{(d,}\ J=8.0), 3.40 \ \mathrm{(m)} & 61.1 \\ 2''-O-Api & & \\ \mathrm{CH}(1''') & & 5.35 \ \mathrm{(d,}\ J=4.0) & 109.2 \\ \mathrm{CH}(2''') & & 3.80 \ \mathrm{(m)} & 76.2 \\ \mathrm{CH}(3''') & & 3.47 \ \mathrm{(m)} & 78.0 \\ \mathrm{CH}_2(4''') & & 3.31 \ \mathrm{(d,}\ J=8.0) & 63.4 \\ \mathrm{CH}_2(5''') & & 3.57 \ \mathrm{(m)}, 3.86 \ \mathrm{(m)} & 73.5 \end{array}$	CH(5")	3.21 (m)	76.7	3.68 (m)	75.1
$\begin{array}{llllllllllllllllllllllllllllllllllll$	CH ₂ (6")	3.43 (m), 3.72 (d, $J = 11.3$)	60.8	3.60 (d, $J = 4.0$), 3.94 (d, $J = 4.0$)	69.0
CH(1''') $4.15 (d, J = 8.0)$ 103.5CH(2''') $2.95 (m)$ 73.5CH(3''') $3.79 (m)$ 76.7CH(4''') $3.02 (m)$ 70.1CH(5''') $3.10 (m)$ 76.9CH_2(6''') $3.64 (d, J = 8.0), 3.40 (m)$ 61.12''-O-Api $CH(1''')$ $5.35 (d, J = 4.0)$ 109.2CH(1''') $3.80 (m)$ 76.2CH(3''') $3.47 (m)$ 78.0CH ₂ (4''') $3.31 (d, J = 8.0)$ 63.4CH ₂ (5''') $3.57 (m), 3.86 (m)$ 73.5	6"-O-Glc				102 5
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	CH(1''')			4.15 (d, J = 8.0)	103.5
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	CH(2")			2.95 (m)	73.5
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	CH(3''')			3.79 (m)	/6./
CH(5") 3.10 (m) 76.9 CH_2(6''') $3.64 \text{ (d, } J = 8.0), 3.40 \text{ (m)}$ 61.1 $2''-O-Api$ $CH(1''')$ $5.35 \text{ (d, } J = 4.0)$ 109.2 CH(2''') 3.80 (m) 76.2 CH(3''') 3.47 (m) 78.0 CH ₂ (4''') $3.31 \text{ (d, } J = 8.0)$ 63.4 CH ₂ (5''') $3.57 \text{ (m)}, 3.86 \text{ (m)}$ 73.5	CH(4''')			3.02 (m)	70.1
$CH_2(6^m)$ $3.64 (d, J = 8.0), 5.40 (m)$ 61.1 $2''-O-Api$ $CH(1''')$ $5.35 (d, J = 4.0)$ 109.2 $CH(2''')$ $3.80 (m)$ 76.2 $CH(3''')$ $3.47 (m)$ 78.0 $CH_2(4''')$ $3.31 (d, J = 8.0)$ 63.4 $CH_2(5''')$ $3.57 (m), 3.86 (m)$ 73.5	CH(5''')			3.10 (m)	/0.9
$2'''-O-Apt$ $5.35 (d, J = 4.0)$ 109.2 $CH(1''')$ $3.80 (m)$ 76.2 $CH(3''')$ $3.47 (m)$ 78.0 $CH_2(4''')$ $3.31 (d, J = 8.0)$ 63.4 $CH_2(5''')$ $3.57 (m), 3.86 (m)$ 73.5	CH ₂ (6 ^m)			3.64 (d, J = 8.0), 3.40 (m)	61.1
$\begin{array}{c} \text{CH}(2^{\prime\prime\prime\prime}) & 3.80 \text{ (m)} & 76.2 \\ \text{CH}(3^{\prime\prime\prime\prime}) & 3.47 \text{ (m)} & 78.0 \\ \text{CH}_2(4^{\prime\prime\prime\prime}) & 3.31 \text{ (d, } J = 8.0) & 63.4 \\ \text{CH}_2(5^{\prime\prime\prime\prime}) & 3.57 \text{ (m)}, 3.86 \text{ (m)} & 73.5 \end{array}$	2" -O-Api CH(1 ^{////})			5.35 (d, $J = 4.0$)	109.2
$CH(3''')$ 3.47 (m) 78.0 $CH_2(4''')$ $3.31 \text{ (d, } J = 8.0)$ 63.4 $CH_2(5''')$ $3.57 \text{ (m)}, 3.86 \text{ (m)}$ 73.5	CH(2"")			3.80 (m)	76.2
$CH_2(4^{\prime\prime\prime\prime})$ $3.31 (d, J = 8.0)$ 63.4 $CH_2(5^{\prime\prime\prime\prime})$ $3.57 (m), 3.86 (m)$ 73.5	CH(3 ^{////})			3.47 (m)	78.0
CH ₂ (5 ^{///}) 3.57 (m), 3.86 (m) 73.5	$CH_{2}(4'''')$			3.31 (d, J = 8.0)	63.4
	CH ₂ (5"")			3.57 (m), 3.86 (m)	73.5

Table 1. ¹H (400 MHz) and ¹³C NMR (100 MHz) spectral data for compounds **1** and **2** (in DMSO- d_6 , δ in ppm, J in Hz).

spectrometer (ThermoFinnigan, San Jose, CA, USA). Semi-preparative HPLC was conducted on a LabAlliance Series 1500 pump equipped with a LabAlliance Model 500 variable wavelength detector and an Agilent Zorbax SB-C18 semi-preparative column ($9.4 \times 250 \text{ mm}$, $5 \mu \text{m}$). Sephadex LH-20 (GE Healthcare Bio-Sciences

Corporation, formerly Amersham Pharmacia Biotechnology Incorporation, NJ, USA), polyamide (80–100 mesh; Linjiang Reagent Chemical Corporation, Jiangsu, China), and ODS (YMC Company, Ltd, Kyoto, Japan) were used for column chromatography (CC). TLC were performed on silica gel plates (Branch of Qingdao Haiyang Chemical Plant, Qingdao, China) and polyamide sheets (Taizhou Si-Jia Biochemical Plastic Company, Zhejiang, China) with spots examined in the UV light at 254 nm.

3.2 Plant material

The seeds of *C. Chinensis* Lam. were collected in Beijing in fall 2000 and were identified by Dr Min Ye. A voucher specimen (No. 20000901) has been deposited at the Pharmacognosy Biotechnology Laboratory, School of Pharmaceutical Sciences, Peking University, China.

3.3 Extraction and isolation

Air-dried seeds of C. Chinensis (5 kg) were pulverized and sieved (40 mesh), defatted with petroleum ether under reflux, and then extracted with 95% ethanol under reflux for three times (2h for each time). The extract was concentrated in a rotary vacuum evaporator to give 600 g of crude residue. The residue was dispersed in hot water and then extracted with CHCl₃. The water layer afforded 500 g of the residue which was subjected to a polyamide column and eluted with aqueous ethanol $(0\% \rightarrow 40\%, v/v)$ to give eight fractions (Fr.1-8). Fr.7 (7.5 g) was subjected to an ODS CC (eluted with gradient aqueous methanol, $10\% \rightarrow 100\%$, v/v) and yielded 10 fractions (Fr.7.1-7.10). Fr.7.8 was then subjected to semi-preparative HPLC and eluted with aqueous methanol (45%, v/v, 2 ml/min) to afford 2 (35 mg), 3 (140 mg), and 4 (36 mg). Fr.7.5 was subjected to Sephadex LH-20 CC and eluted with methanol to afford 5 (30 mg). Fr.7.2 was

subjected to semi-preparative HPLC and eluted with aqueous acetonitrile (10%, v/v, 2 ml/min) to afford **6** (6 mg) and **7** (20 mg). Fr.5 (12 g) was also subjected to ODS CC (eluted with gradient aqueous methanol, $10\% \rightarrow 100\%$, v/v) to give six fractions (Fr.5.1–5.6). Fr.5.6 was then applied to semi-preparative HPLC and eluted with aqueous acetonitrile (33%, v/v, 2 ml/min) to give **1** (60 mg) and **8** (5 mg).

3.3.1 Cuscutoside C(1)

White amorphous powder. mp 112–113°C. $[\alpha]_D^{20}$ – 73.5 (c = 0.067, MeOH). UV (MeOH) λ_{max} (nm): 291.0, 237.0. IR (KBr) ν_{max} (cm⁻¹): 3383, 2883, 1630, 1503, 1485, 1440, 1400, 1347, 1249, 1180, 1074, 1036, 931, 864, 810, 717, 559. ¹H and ¹³C NMR spectral data: see Table 1. HR-ESI-MS: m/z 555.1481 [M + Na]⁺ (calcd for C₂₆H₂₈O₁₂Na, 555.1473).

3.3.2 Cuscutoside D (2)

White amorphous powder. mp 153– 154°C. $[\alpha]_D^{20}$ – 99.0 (c = 0.067, MeOH). UV (MeOH) λ_{max} (nm): 291.5, 237.0. IR (KBr) ν_{max} (cm⁻¹): 3394, 2884, 1705, 1632, 1503, 1485, 1440, 1249, 1179, 1068, 1037, 931, 875, 813, 717, 560. ¹H and ¹³C NMR spectral data: see Table 1. HR-ESI-MS: m/z 849.2417 [M + Na]⁺ (calcd for C₃₇H₄₆O₂₁Na, 849.2424).

3.4 Acid hydrolysis of 1 and 2

Compounds 1 (3 mg) and 2 (5 mg) were heated in 2 ml of 2 M CF₃COOH at 90°C for 12 h, respectively. The reaction mixture was then diluted with 2 ml H₂O and extracted with CHCl₃ (4 ml × 3). The water layer was evaporated to dryness and then dissolved with 1 ml of MeOH. The solutions were then subjected to silica gel TLC together with authentic D-Glc and D-Api samples, and developed using *n*-BuOH–Me₂CO–H₂O (4:5:1) as the solvent. The R_f values of D-Glc and D-Api were 0.63 and 0.80, respectively.

Acknowledgement

We are grateful for the financial support from the Peking University Health Science Center (No. 985-2-119-121).

References

- H.M. Mi, C. Guo, and H.T. Song, *Chin. Tradit. Herb. Drugs* 22, 547 (1991).
- [2] S.J. Peng, R.K. Lu, and L.H. Yu, Chin. J. Integr. Tradit. West. Med. 17, 145 (1997).
- [3] Y.B. Xiong and C.H. Zhou, *Chin. Pharm. J.* 29, 89 (1994).
- [4] Z.R. Liu, P.T. Luo, and T.J. Fu, *Nat. Prod. Res. Dev.* 6, 532 (2004).
- [5] G.S. Li, Y.Y. Chen, and S.C. Li, *Chin. J. Tradit. Med. Sci. Technol.* 4, 256 (1997).
- [6] L.G. Gu, M. Ye, and Y.N. Yan, *Chin. J. Inf. Tradit. Chin. Med.* 8, 42 (2001).
- [7] X.G. Cai and S.X. Zhao, *Pharm. Clin. Chin. Mater. Med.* 23, 27 (2007).
- [8] Y.M. Xie, L.L. Qin, and X.D. Yu, Chin. J. Inf. Tradit. Chin. Med. 12, 22 (2005).
- [9] J. Guo, S.G. Bai, and Y.M. Wang, *Chin. J. Gerontol.* 16, 37 (1996).

- [10] L.P. Liu, J. Liu, and H.F. Chen, J. Chin. Med. Mater. 24, 839 (2001).
- [11] M. Ye, Y.N. Yan, and D.A. Guo, *Rapid Commun. Mass Spectrom.* 19, 1469 (2005).
- [12] S. Yahara, H. Domoto, C. Sugimura, T. Nohara, Y. Niiho, Y. Nakajima, and H. Ito, *Phytochemistry* **37**, 1755 (1994).
- [13] S.X. Xiang, Z.S. He, and Y. Ye, *Chin. J. Chem.* **19**, 282 (2001).
- [14] M. Ye, Y.N. Yan, L. Qiao, and X.M. Ni, *China J. Chin. Mater. Med.* 27, 115 (2002).
- [15] X.H. Xu, B.Q. Ruan, S.H. Jiang, and D.Y. Zhu, *Chin. J. Nat. Med.* 3, 93 (2005).
- [16] K. Iwai, N. Kishimoto, Y. Kakino, K. Mochida, and T. Fujita, J. Agric. Food Chem. 52, 4893 (2004).
- [17] P.A. Davidse, J.L.M. Dillen, A.M. Heyns, T.A. Modro, and P.H. Rooyen, *Can. J. Chem.* 68, 741 (1990).
- [18] N.H. El-Sayed, M. Wojcinska, K. Drost-Karbowska, I. Matlawska, J. Williams, and T.J. Mabry, *Phytochemistry* 60, 835 (2002).