

 Tetrahedron Letters, Vol. 37, No. 25, pp. 4369-4372, 1996

 Copyright © 1996 Elsevier Science Ltd

 Printed in Great Britain. All rights reserved

 0040-4039/96 \$15.00 + 0.00

PII: S0040-4039(96)00819-2

## Asymmetric Synthesis. XLI.<sup>1</sup> Totally Stereoselective Synthesis of 1,3-disubstituted tetrahydroisoquinolines via the CN(R,S) Method.

Grace Gosmann, Dominique Guillaume, Henri-Philippe Husson\*.

Laboratoire de Chimie Thérapeutique associé au CNRS, Faculté des Sciences Pharmaceutiques et Biologiques, Université R. Descartes, 4, Av. de l'Observatoire, 75270 Paris Cedex 06, France.

Abstract : Optically active *cis*- or *trans*- 1,3-disubstituted tetrahydroisoquinolines can be prepared selectively from the same oxazolidine 4. This latter is easily obtained from keto-acid 1 and (R)-(-)-phenylglycinol. Copyright © 1996 Elsevier Science Ltd

Although there are several methods for the asymmetric synthesis of 1-alkyl or 1-aryl tetrahydroisoquinolines,<sup>2</sup> the only example of asymmetric synthesis of all isomers of 1,3-disubstituted tetrahydroisoquinolines has been described by Bringmann *et al.*<sup>3</sup> However, while this method allows a variety of substituents at C-1, it is limited for the substitution at C-3, which requires the preparation of enantiomerically pure 1-alkyl-2-phenylethylamine. Indeed, 1,3-disubstituted tetrahydroisoquinoline is a structural feature found in a large variety of alkaloids<sup>4</sup> and we have initiated a project aimed at the synthesis of natural products in this series and analogues of biological interest.

In continuation of our work on the development of general synthetic methodologies for the construction of a wide diversity of structures, by means of equivalents of non-racemic synthons using phenylglycinol as source of chirality,<sup>5</sup> we present here an efficient approach to the title compounds in enantiomerically pure form.

The condensation of  $\delta$ -ketoacids with optically active  $\beta$ -aminoalcohols is known to afford chiral bicyclic lactams.<sup>2b</sup> More interestingly, we have discovered that the stepwise condensation of 1<sup>6</sup> with (*R*)-(-)-phenylglycinol (Scheme 1) led to amide<sup>7</sup> 2 (yield 70%) which was cyclized to the isoquinolinium salt 3 (90% yield) by treatment with trifluoroacetic acid. The sodium borohydride reduction of 3 under controlled conditions furnished oxazolidine 4 (50% yield) as a 60:40 mixture of stereomers at C-3. The absolute configuration at C-1 was unambiguously established as (S) from the synthesis of the known (-)-norcryptostyline III 6<sup>8</sup> by sodium borohydride treatment of 4 followed by hydrogenolysis of the chiral appendage.

At this stage, it was possible to envisage the stereoselective creation of an R or S chiral centre at C-3 of oxazolidine 4 using the CN(R,S) strategy which has proved to be a powerful tool in this and other laboratories. 5a,d,9



Thus, the single (1S, 3S) aminoalcohols  $7^{10}$  (Scheme 2) were isolated on treatment of oxazolidine 4 with methyl magnesium bromide or benzyl magnesium chloride<sup>11</sup> in the presence of a catalytic amount of aluminium chloride (65% yield).

The access to the (15, 3R) diastereometric C-3 substituted derivative is a less obvious goal. On the basis of observations made during earlier work, <sup>12</sup> we reasoned that it was necessary to transform oxazolidine 4 into aminonitrile 8. Reaction of 4 with trimethylsilyl cyanide in the presence of a catalytic amount of aluminium chloride gave 8 (65% yield). This latter was deprotonated (LDA, 4 equiv.) to give the corresponding anion which could be alkylated with methyl iodide or benzyl bromide to give 9. As the crude product contained significant quantities of the recyclized compounds 10, it was treated by a solution of 5% aqueous hydrofluoric acid in acetonitrile to effect complete cyclization of 9 to 10. Reductive opening of the oxazolidine ring of 10 was performed in the presence of trifluoroacetic acid yielding aminoalcohols 11 (60% yield).<sup>10</sup>

The relative configuration of the newly-created asymmetric centre could be determined by comparison of the <sup>13</sup>C NMR data of 7 and 11: a  $\gamma$ -gauche effect observed for C-3 of 7 is diagnostic of a *trans* relationship between the C-1 and C-3 substituents.<sup>13</sup>

Scheme 2



The remarkable stereocontrol in the reaction leading to diastereomeric compounds 7 and 11 was interpreted in terms of an elimination-addition mechanism wherein Grignard reagents or hydride ion approach the intermediate iminium ion under stereoelectronic control from the axial direction on the less hindered face.

As in the preparation of 6, the chiral appendage of 7 and 11 could be easily removed by hydrogenolysis to give a secondary amine allowing further substitutions or cyclizations.

Using (S)-(+)-phenylglycinol, it should be possible to obtain the enantiomeric C-1 substituted compounds and hence the four isomers of 1,3-disubstituted tetrahydroisoquinolines in optically-pure form.

Acknowledgements : We thank Faculdade de Farmacia, Universidade Federal do Rio Grande do Sul (Porto Alegre, RS, Brasil) and CAPES/COFECUB for a research grant to GG.

## **References** and notes

- 1. For part XL, see Micouin, L.; Quirion, J.-C.; Husson, H.-P. *Tetrahedron Lett.* 1996, 37, 849-852. 2. (a) Carbonnelle, A. C.; Gott, V.; Roussi, G. *Heterocycles*, 1993, 36, 1763-1769; (b) Munchhof, M. J.;
- Meyers, A. I. J. Org. Chem. 1995, 60, 7086-7087 and references cited herein.
- 3. Bringmann, G.; Weirich, R.; Reuscher, H.; Jansen, J. R.; Kinzinger, L.; Ortmann, T. Liebigs Ann. Chem. 1993, 877-888.
- 4. (a) Takahashi, K.; Tomita, F. Antibiot. 1983, 36, 468-470. (b) Bringmann, G. The Alkaloids, A. Brossi
- Ed.; Academic Press: New York 1986, 29, p. 141-184.
  5. (a) Royer, J; Husson, H.-P, Janssen Chim. Acta 1993, 11, 3-8. (b) Lienard, P.; Quirion, J.-C.; Husson, H.-P. Tetrahedron 1993, 49, 3995-4006. (c) Micouin, L.; Varea, T.; Riche, C.; Chiaroni, A.; Quirion, J.-C.; Husson, H.-P. Tetrahedron Lett. 1994, 35, 2529-2532. (d) Husson, H.-P.; Royer, J. Advances in the Use of Synthons in Organic Chemistry, A. Dondoni Ed.; JAI Press: London 1995, 2, p. 1-68.
- 6. Prepared in 70% yield by stirring methyl-3,4-dimethoxyphenylacetate and 3,4,5-trimethoxybenzoyl chloride in the presence of PCl5 and AlCl3 in CH2Cl2 for 2 days then saponification.
- 7. All new compounds were fully characterised by IR, MS, <sup>1</sup>H and <sup>13</sup>C NMR spectroscopy and exhibit satisfactory combustion analyses for C, H, and N.
- 8. Brossi, A.; Teitel, S. Helv. Chim. Acta 1971, 54, 1564-1571.
- 9. Guerrier, L.; Royer, J.; Grierson, D. S.; Husson, H.-P. J. Am. Chem. Soc., 1983, 105, 7754-7755.
- **10.** Compound **7a**, oil:  $[\alpha]_D^{20} = +21$  (c 0.80, CHCl<sub>3</sub>). <sup>1</sup>H NMR (300MHz, CDCl<sub>3</sub>)  $\delta$  7.20 (m, 3H), 7.05 (m, 2H), 6.40 (s, 2H), 6.30 (s, 1H), 6.22 (s, 1H), 5.20 (s, 1H), 4.20 (m, 2H), 3.85 (s, 3H), 3.80 (m, 1H), 3.80 (s, 9H), 3.70 (s, 3H), 3.50 (m, 1H), 2.62 (dd, 1H, J= 17.0, 5.0 Hz), 2.45 (dd, 1H, J=17.0, 10.0 Hz), 1.3 (d, 3H, J = 6.7 Hz); <sup>13</sup>C (75 MHz, CDCl<sub>3</sub>)  $\delta$  152.8, 140.8, 130.3, 128.6, 128.1; 127.7, 127.4,

110.9, 106.5, 61.6, 60.8, 60.6, 60.4, 56.1, 55.9, 55.6, 46.9, 33.8, 19.3. **7b**, oil:  $[\alpha]_D^{20} = +5$  (c 0.94,

CHCl<sub>3</sub>). <sup>1</sup>H NMR (300MHz, CDCl<sub>3</sub>)  $\delta$  7.10 (m, 5H), 6.95 (m, 3H), 6.85 (m, 2H), 6.30 (s, 2H), 6.20 (s, 1H), 6.15 (s, 1H), 5.20 (s, 1H), 4.05 (m, 2H), 3.80 (s, 3H), 3.65 (s, 12H), 3.6 (m, 2H), 3.00 (dd, 1H, 1H), 3.80 (s, 2H), 3.6 (m, 2H), 3.80 (s, 2H) J= 14.4, 8.0 Hz), 2.85 (dd, 1H, J= 14.4, 7.5 Hz), 2.40 (d, 2H, J=7.5 Hz); <sup>13</sup>C (75 MHz, CDCl<sub>3</sub>)  $\delta$  152.7, 147.2, 146.9, 140.5, 139.1, 138.5, 128.8, 128.5, 128.2, 127.6, 126.2, 111.1, 110.8, 106.6, 61.7, 61.1, 60.8, 60.3, 55.9, 55.6, 51.9, 38.4, 31.4.

Compound **11a**, oil:  $[\alpha]_D^{20} = -31$  (c 0.61, CHCl<sub>3</sub>). <sup>1</sup>H NMR (300MHz, CDCl<sub>3</sub>)  $\delta$  7.10 (s, 5H), 6.51 (s, 1H), 6.45 (s, 2H), 6.05 (s, 1H), 4.78 (s, 1H), 3.90 (m, 3H), 3.80 (s, 3H), 3.75 (s, 3H), 3.71 (s, 6H), 3.61 (s, 3H), 3.15 (m, 1H), 2.50 (dd, 1H, J= 15.0, 11.0 Hz), 2.35 (dd, 1H, J=15.0, 4.5 Hz), 1.30 (d, 3H, J= 6.1 Hz); <sup>13</sup>C NMR (75 MHz, CDCl<sub>3</sub>) δ 152.8, 146.6, 141.6, 138.8, 130.4, 128.3, 128.1, 127.5, 110.5, 110.3,

104.0, 66.9, 62.4, 60.8, 60.4, 55.9, 55.8, 54.7, 35.9, 23.5. 11b, oil:  $[\alpha]_D^{20} = -40$  (c 1.04, CHCl3). <sup>1</sup>H

NMR (300MHz, CDCl<sub>3</sub>) § 7.21 (m, 8H), 7.10 (m, 2H), 6.60 (s, 2H), 6.50 (s, 1H), 6.10 (s, 1H), 4.90 (s, 1H), 4.10 (m, 2H), 4.00 (m, 1H), 3.88 (s, 3H), 3.80 (s, 9H), 3.7 (s, 3H), 3.50 (m, 1H), 3.10 (dd, 1H, J= 13.0, 4.0 Hz), 2.55 (m, 2H), 2.30 (dd, 1H, J=15.0, 9.5 Hz); <sup>13</sup>C NMR (75 MHz, CDCl<sub>3</sub>) δ 152.9, 147.5, 146.7, 140.8, 139.3, 139.1, 136.6, 130.6, 129.2, 128.4, 128.2, 127.9, 127.7, 126.2, 110.8, 104.7, 68.0, 62.3, 60.9, 60.7, 60.5, 56.0, 55.8, 44.5, 32.2.

- 11. As part of their work on the development of the Zincke reaction,<sup>5d</sup> Marazano et al. have recently described the reaction of a substrate closely related to 4 with Grignard reagents: Barbier D. Ph.D. Dissertation, Université Paris-Sud, Orsay 1995.
- 12. Grierson, D. S.; Royer, J.; Guerrier, L.; Husson, H.-P. J. Org. Chem. 1986, 51, 4475-4477.

13. Tomioka, K.; Kubota, Y.; Koga, K. Tetrahedron, 1993, 49, 1891-1900.

(Received in France 6 April 1996; accepted 25 April 1996)