


Geminal Silicon/Zinc Reagent as an Equivalent of Difluoromethylene Bis-carbanion


Mikhail D. Kosobokov,[†] Vitalij V. Levin,[†] Artem A. Zemtsov,[†] Marina I. Struchkova,[†] Alexander A. Korlyukov,[‡] Dmitry E. Arkhipov,[‡] and Alexander D. Dilman^{*,†}[†]N. D. Zelinsky Institute of Organic Chemistry, Leninsky prosp. 47, 119991 Moscow, Russian Federation[‡]A. N. Nesmeyanov Institute of Organoelement Compounds, Vavilov str. 28, 119991 Moscow, Russian Federation

Supporting Information

ABSTRACT: A new difluorinated reagent, [difluoro(trimethylsilyl)methyl]zinc bromide, bearing C–Zn and C–Si bonds is described. The reagent is conveniently prepared by cobalt-catalyzed halogen/zinc exchange. It can be coupled with two different C-electrophiles in a stepwise manner (with allylic halides for C–Zn bond and aldehydes for C–Si bond) affording products containing a difluoromethylene fragment.




The ability of fluorine to affect biological properties of organic compounds¹ has led to the discovery and successful use of numerous fluorine-containing drugs.^{1,2} In the majority of these molecules, the fluorine is present either as a substituent or in the form of a trifluoromethyl group. At the same time, compounds bearing difluoromethylene units are less frequent among marketed medicines, but interest in this structural motive is growing. Indeed, it has been noted that the CF₂ fragment is isosteric to ethereal oxygen^{1a,b,3} and, in some cases, even to carbonyl or *gem*-dimethylmethylene groups⁴ (Scheme 1).

Scheme 1. Bioisosterism of CF₂ Fragment


Typically, synthesis of CF₂-containing products either involves a deoxofluorination reaction, which requires harsh reagents,⁵ or relies on a building block approach, which is laborious.^{6,7} Recently, we described a strategy for assembling *gem*-difluorinated compounds from difluorocarbene, nucleophile, and electrophile.⁸ Herein, we report a reagent for coupling of binucleophilic CF₂ unit with two electrophilic species.

Previously, sulfur⁹ and phosphorus¹⁰ reagents were shown to act as equivalents of –CF₂– synthon in reactions with two aldehydes affording 1,3-diols (Scheme 2). However, the use of fluorinated silanes would be desirable, since the carbon–silicon bond is readily cleaved by fluoride anion or even weaker bases.¹¹ To this end, the bis-silicon reagent (TMS)₂CF₂ was reported but failed to serve as the bis-carbanion in reaction with aldehydes.¹² Concerning polar organometallics, preparation of α -lithiated and α -magnesiated silanes was attempted but was unsuccessful.¹³ In this work, we describe the silicon/zinc reagent with C–Zn and

Scheme 2. Equivalents of CF₂ Bis-carbanion Synthon

C–Si bonds being sequentially involved in coupling with suitable electrophiles.

In the initial experiment, (bromodifluoromethyl)trimethylsilane (Me₃SiCF₂Br, **1**)¹⁴ was treated with zinc dust. Reagent **2** was detected in small amounts along with homocoupling product **3**, as determined by ¹⁹F NMR spectroscopy¹⁵ (Scheme 3). Attempts to improve this reductive protocol failed despite extensive variation of reaction conditions.^{16,17}

Scheme 3. Attempted Reductive Zincation of **1**

Then, the generation of organozinc species **2** was evaluated by means of bromine/zinc exchange using isopropylzinc iodide (Table 1). The reaction of silane **1** with 1.6 equiv of *i*-PrZnI was slow at room temperature, providing modest yields of **2** (entries

Received: January 22, 2014

Published: February 19, 2014

Table 1. Preparation of Reagent 2

$ \begin{array}{ccc} \text{Me}_3\text{Si}-\text{C}(\text{F})_2-\text{Br} & \xrightarrow[\text{20 h, solvent, temp}]{i\text{-PrZnI, additive}} & \text{Me}_3\text{Si}-\text{C}(\text{F})_2-\text{ZnBr} \\ \text{1} & & \text{2} \end{array} $						
no.	solvent	RZnI (equiv)	additive (equiv)	temp (°C)	conv of 1 ^b (%)	yield of 2 ^b (%)
1	THF	1.6	none	20	65	30
2	THF/NMP	1.6	none	20	65	15
3	THF	1.6	LiCl (1.6)	20	100	0
4	THF	1.2	Ni(acac) ₂ (0.01)	5	29	12
5	THF	1.2	NiBr ₂ ·(PPh ₃) ₂ (0.01)	5	49	24
6	THF	1.2	CoBr ₂ ·(PPh ₃) ₂ (0.01)	5	62	38
7	THF	1.2	CoBr ₂ ·dppe (0.01)	5	95	30
8	THF/DG	1.2	CoBr ₂ ·dppe (0.01)	5	100	90
9	THF/DG	1.0	CoBr ₂ ·dppe (0.01)	5	95	88
10 ^c	THF/DG	1.0	CoBr ₂ ·dppe (0.01)	5	95	87
11	THF/DG	1.2	CoBr ₂ ·dppe(0.005)	5	97	88
12	THF/DG	1.2	CoBr ₂ ·dppp (0.01)	5	86	60
13	THF/DG	1.2	CoBr ₂ ·dppb (0.01)	5	78	63
14	THF/DG	1.2	CoBr ₂ (0.01)	5	43	28
15 ^d	THF/DG	1.2	CoBr ₂ ·dppe (0.01)	5	69	34

^aTHF: tetrahydrofuran. NMP: *N*-methylpyrrolidone. DG: diglyme. ^bDetermined by ¹⁹F NMR using PhCF₃ as internal standard. ^cReaction time 48 h. ^d*i*-PrZnBr was used.

1 and 2). Then, the effect of various catalysts was evaluated.¹⁸ The significant rate acceleration was observed by adding 1 mol % of CoBr₂·dppe complex (dppe, 1,2-bis(diphenylphosphino)ethane), which allowed the temperature to decrease as well the amount of *i*-PrZnI (to 1.2 equiv, entry 7). Rewardingly, the addition of diglyme as a cosolvent (THF/diglyme, 1/1) led to complete conversion together with 90% yield of 2 (entry 8). The use of a precisely stoichiometric amount of *i*-PrZnI afforded almost the same yield of 2 at 95% conversion (entry 9). Other evaluated cobalt and nickel salts and complexes were notably less efficient. When isopropylzinc bromide was employed instead of isopropylzinc iodide, the reaction was slow (entry 15).

We also investigated the stability, composition, and structure of organozinc reagent 2. The obtained THF/diglyme solutions of 2 are reasonably stable and can even be stored at low temperatures. For example, after 72 h at −30 °C, no decrease of reagent concentration was noted, whereas after the same time at room temperature only 25% of 2 decomposed (¹⁹F NMR control).

¹⁹F NMR analysis of 2 in THF/diglyme solution showed two close signals (−130.4 and −130.7 in a ratio 93:7, respectively), likely owing to Schlenk equilibrium. Interestingly, when isolated crystals of 2·diglyme (vide infra) were dissolved in CD₃CN, only one signal was observed.

It was surprising to find that reagent 2 contains bromide at zinc despite using isopropylzinc iodide for the halogen/zinc exchange. Indeed, such a halogen distribution pattern was established by identification of isopropyl iodide byproduct.¹⁹ Furthermore, we were fortunate to obtain highly air-sensitive crystals of 2 coordinated with diglyme, and X-ray analysis unambiguously demonstrated the presence of bromide (Figure 1).

We suggest a tentative mechanism of bromine/zinc exchange accounting for unusual halogen redistribution (Scheme 4). At first, the cobalt(II) complex is reduced to generate catalytically active Co(I) iodide species.²⁰ Subsequently, oxidative addition leads to Co(III) intermediate,²¹ which serves as iodinating reagent with respect to isopropylzinc iodide. At the latter step the isopropyl iodide is formed along with generation of the Zn–Br





Figure 1. X-ray structure of 2·diglyme. Hydrogen atoms are omitted for clarity.

Scheme 4. Proposed Mechanism of Halogen/Zinc Exchange



bond. The final step involves Co/Zn transmetalation providing reagent 2.

Having developed a method for the preparation of reagent 2, we proceeded with exploration of its properties as equivalent of difluoromethylene bis-carbanion. Given the greater polarity of C–Zn bond compared to that of C–Si bond, reactions of 2 as organozinc species were first evaluated. For fluorinated organozinc reagents, copper(I)-catalyzed arylation²² and allylation^{8b,23} have been reported. Since arylations are performed under harsh

■ ASSOCIATED CONTENT

■ Supporting Information

Experimental procedures, compound characterization data, copies of NMR spectra for all compounds, and X-ray data for 2-diglyme (CIF). This material is available free of charge via the Internet at <http://pubs.acs.org>.

■ AUTHOR INFORMATION

Corresponding Author

*E-mail: adil25@mail.ru.

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

This work was supported by the Ministry of Science (project MD-1151.2011.3), Russian Foundation for Basic Research (projects 13-03-12074, 14-03-00293, 14-03-31253_mol_a, 14-03-31265_mol_a), and the Russian Academy of Sciences. We are grateful to R. Novikov (N. D. Zelinsky Institute of Organic Chemistry) for 2D-NMR measurements and to G. Ivanov (Higher Chemical College, Russian Academy of Sciences) for experimental assistance.

■ REFERENCES

- (1) (a) *Fluorine in Medicinal Chemistry and Chemical Biology*; Ojima, I., Ed.; John Wiley & Sons: Chichester, 2009. (b) Begue, J.-P.; Bonnet-Delpont, D. *Bioorganic and Medicinal Chemistry of Fluorine*; Wiley-VCH: Weinheim, 2008. (c) O'Hagan, D.; S. Rzepa, H. *Chem. Commun.* **1997**, 645–652. (d) Purser, S.; Moore, P. R.; Swallow, S.; Gouverneur, V. *Chem. Soc. Rev.* **2008**, 37, 320–330. (e) Müller, K.; Faeh, C.; Diederich, F. *Science* **2007**, 317, 1881–1886.
- (2) (a) O'Hagan, D. *J. Fluorine Chem.* **2010**, 131, 1071–1081. (b) Isanbor, C.; O'Hagan, D. *J. Fluorine Chem.* **2006**, 127, 303–319.
- (3) (a) Blackburn, G. M.; England, D. A.; Kolkman, F. *J. Chem. Soc., Chem. Commun.* **1981**, 930–932. (b) Hakogi, T.; Yamamoto, T.; Fujii, S.; Ikeda, K.; Katsumura, S. *Tetrahedron Lett.* **2006**, 47, 2627–2630. (c) Xu, Y.; Aoki, J.; Shimizu, K.; Umezū-Goto, M.; Hama, K.; Takanezawa, Y.; Yu, S.; Mills, G. B.; Arai, H.; Qian, L.; Prestwich, G. D. *J. Med. Chem.* **2005**, 48, 3319–3327.
- (4) Meanwell, N. A. *J. Med. Chem.* **2011**, 54, 2529–2591.
- (5) Al-Maharik, N.; O'Hagan, D. *Aldrichimica Acta* **2011**, 44, 65–75.
- (6) (a) Tozer, M. J.; Herpin, T. F. *Tetrahedron* **1996**, 52, 8619–8683. (b) Qing, F.-L.; Zheng, F. *Synlett* **2011**, 1052–1072.
- (7) *gem*-Difluorocyclopropanes constitute a specific class, which is generally accessed by difluorocarbene addition; see: Dolbier, W. R.; Battiste, M. A. *Chem. Rev.* **2003**, 103, 1071–1098.
- (8) (a) Levin, V. V.; Zemtsov, A. A.; Struchkova, M. I.; Dilman, A. D. *Org. Lett.* **2013**, 15, 917–919. (b) Zemtsov, A. A.; Kondratyev, N. S.; Levin, V. V.; Struchkova, M. I.; Dilman, A. D. *J. Org. Chem.* **2014**, 79, 818–822.
- (9) (a) Prakash, G. K. S.; Hu, J.; Mathew, T.; Olah, G. A. *Angew. Chem., Int. Ed.* **2003**, 42, 5216–5219. (b) Prakash, G. K. S.; Hu, J. *Acc. Chem. Res.* **2007**, 40, 921–930.
- (10) Beier, P.; Alexandrova, A. V.; Zibinsky, M.; Surya Prakash, G. K. *Tetrahedron* **2008**, 64, 10977–10985.
- (11) Prakash, G. K. S.; Yudin, A. K. *Chem. Rev.* **1997**, 97, 757–786.
- (12) (a) Yudin, A. K.; Prakash, G. K. S.; Deffieux, D.; Bradley, M.; Bau, R.; Olah, G. A. *J. Am. Chem. Soc.* **1997**, 119, 1572–1581. (b) For a homologous vicinal bis-carbanion synthon $-\text{CF}_2\text{C}(\equiv\text{NR})-$, see: Kobayashi, T.; Nakagawa, T.; Amii, H.; Uneyama, K. *Org. Lett.* **2003**, 5, 4297–4300.
- (13) Fuchikami, T.; Ojima, I. *J. Organomet. Chem.* **1981**, 212, 145–153.
- (14) $\text{Me}_3\text{SiCF}_2\text{Br}$ can be readily obtained from Me_3SiCF_3 ; see: (a) Kosobokov, M. D.; Dilman, A. D.; Levin, V. V.; Struchkova, M. I. *J. Org. Chem.* **2012**, 77, 5850–5855. (b) Li, L.; Wang, F.; Ni, C.; Hu, J. *Angew. Chem., Int. Ed.* **2013**, 52, 12390–12394.
- (15) Compound 3 was previously described; see ref 12a.
- (16) Various additives, which were expected to affect reductive metalation (such as LiCl or copper(I) salts), had an even greater deleterious effect. See: (a) Krasovskii, A.; Malakhov, V.; Gavryushin, A.; Knochel, P. *Angew. Chem., Int. Ed.* **2006**, 45, 6040–6044. (b) Metzger, A.; Schade, M. A.; Knochel, P. *Org. Lett.* **2008**, 10, 1107–1110.
- (17) For preparation of nonfluorinated reagent $\text{Me}_3\text{SiCH}_2\text{ZnX}$, see: (a) Jones, P.; Reddy, C. K.; Knochel, P. *Tetrahedron* **1998**, 54, 1471–1490. (b) Charette, A. B.; Beauchemin, A.; Marcoux, J.-F. *Tetrahedron Lett.* **1999**, 40, 33–36. (c) Hatakeyama, T.; Nakagawa, N.; Nakamura, M. *Org. Lett.* **2009**, 11, 4496–4499.
- (18) Transition-metal catalysts are used for accelerating halogen/zinc exchange using diethylzinc or triorganozincate as metallating reagents; see: (a) Stadtmüller, H.; Lentz, R.; Tucker, C. E.; Stuedemann, T.; Doerner, W.; Knochel, P. *J. Am. Chem. Soc.* **1993**, 115, 7027–7028. (b) Stadtmüller, H.; Vaupel, A.; Tucker, C. E.; Stuedemann, T.; Knochel, P. *Chem.—Eur. J.* **1996**, 2, 1204–1220. (c) Vaupel, A.; Knochel, P. *J. Org. Chem.* **1996**, 61, 5743–5753. (d) Melzig, L.; Diene, C. R.; Rohbogner, C. J.; Knochel, P. *Org. Lett.* **2011**, 13, 3174–3177.
- (19) A solution of **2** was concentrated under vacuum collecting the volatiles in a cold trap. NMR analysis of condensed material indicated *i*-PrI and *i*-PrBr in a ratio 94:6.
- (20) (a) Cahiez, G.; Moyeux, A. *Chem. Rev.* **2010**, 110, 1435–1462. (b) Gosmini, C.; Bégouin, J.-M.; Moncomble, A. *Chem. Commun.* **2008**, 3221–3233.
- (21) Reactions of Co(I) complexes with R_4I to give Co(III) species are known; see: Harrison, D. J.; Gorelsky, S. I.; Lee, G. M.; Korobkov, I.; Baker, R. T. *Organometallics* **2013**, 32, 12–15 and reference cited therein.
- (22) Popov, I.; Lindeman, S.; Daugulis, O. *J. Am. Chem. Soc.* **2011**, 133, 9286–9289.
- (23) Burton, D. J.; Hartgraves, G. A. *J. Fluorine Chem.* **2007**, 128, 1198–1215.
- (24) The use of a previously developed $\text{CuI}/1,10\text{-phenanthroline}$ catalytic system for allylation of fluorinated organozincs (ref 8b) gave slow reaction.
- (25) For the synthesis of silanes **5**, we employed reagent **2** prepared with 1.2 equiv of *i*-PrZnI (entry 8, Table 1). If reagent **2** prepared under the conditions of entry 9 is employed, the residual starting silane $\text{Me}_3\text{SiCF}_2\text{Br}$ hampers purification of silanes **5**.
- (26) Mogi, R.; Morisaki, K.; Hu, J.; Prakash, G. K. S.; Olah, G. A. *J. Fluorine Chem.* **2007**, 128, 1098–1103.