C—F Activation Hot Paper

International Edition: DOI: 10.1002/anie.201604776 German Edition: DOI: 10.1002/ange.201604776

Single C-F Bond Cleavage of Trifluoromethylarenes with an *ortho*-Silyl Group

Suguru Yoshida,* Ken Shimomori, Youngchan Kim, and Takamitsu Hosoya*

Abstract: The transformation of a single C–F bond of trifluoromethylarenes bearing a hydrosilyl group at the ortho position was achieved. The activation of the hydrosilyl group with a trityl cation in the presence of nucleophiles allowed for selective C–F bond functionalization, for example, by allylation, carboxylation, or chlorination. Further derivatization of the resulting fluorosilylarenes afforded various aromatic difluoromethylene compounds.

Organofluorine compounds are an important class of molecules that are widely used in various fields, including the pharmaceutical sciences, agrochemistry, and materials science.^[1] The increasing demand for fluorinated compounds has prompted the development of a variety of methods for introducing a fluorine atom or a fluorinated functional group into organic frameworks. In particular, with the increasing availability of convenient reagents in recent years, various methods for fluorination and trifluoromethylation have been developed.^[2,3] However, methods for synthesizing complex fluorinated compounds are still limited, and new approaches are highly sought-after.

In principle, a diverse range of fluorinated compounds can be straightforwardly prepared from trifluoromethylated compounds by the selective transformation of one of the three C-F bonds of the trifluoromethyl group. However, the high stability of the C-F bond has rendered this transformation difficult to achieve (Figure 1 A).^[4-7] Furthermore, previous studies regarding the transformation of C-F bonds stressed the difficulty of performing single C-F bond cleavage with an aromatic trifluoromethyl group; all of the three C-F bonds of the trifluoromethyl group were cleaved in these cases.^[8-10] The reason behind this difficulty is that the C-F bonds of the diand monofluorinated intermediates are likely to be more reactive than those of the trifluoromethyl-substituted starting material (Figure 1B). For example, Ozerov and co-workers reported the reduction of an aromatic trifluoromethyl group to the corresponding methyl group by C-F bond cleavage with a silvlium species (Figure 1C).^[8b] We assumed that a pinpoint transformation of the aromatic trifluoromethyl group could be achieved through a single C-F bond cleavage

 [*] Dr. S. Yoshida, K. Shimomori, Y. Kim, Prof. Dr. T. Hosoya Laboratory of Chemical Bioscience Institute of Biomaterials and Bioengineering Tokyo Medical and Dental University 2-3-10 Kanda-Surugadai, Chiyoda-ku, Tokyo 101-0062 (Japan) E-mail: s-yoshida.cb@tmd.ac.jp thosoya.cb@tmd.ac.jp

Supporting information for this article can be found under: http://dx.doi.org/10.1002/anie.201604776.

Angew. Chem. Int. Ed. 2016, 55, 1–5

Α в energy energy length (Å) (kcal/mol) length (Å) (kcal/mol) C-H 1.10 105 H₃C-F 1.38 110 C–F 1.38 110 H₂FC-F 1.36 120 HF₂C-F C–CI 1.78 84 1.33 128 C–Br 1.94 70 1.32 131 C Ozerov et al., 2008 Et₂SiH cat. Ph₃C⁺[HCB₁₁H₅Cl₆] D This work transformation fluoride migration & CF₃ Nu of the silyl group nucleophilic attack nucleophile Si Si Ph₃CX Ŕ R' R R'R generation of silyl cation easily activatable with a base role switching

Figure 1. Transformation of an aromatic trifluoromethyl group by C–F bond cleavage. A) Bond lengths and dissociation energies of C–X bonds for homolytic bond cleavage.^[4] B) Bond lengths and dissociation energies of C–F bonds in H_{4–n}CF_n (n = 1-4) for homolytic bond cleavage.^[5] C) Reported transformation of an aromatic trifluoromethyl group with a silylium intermediate.^[8a] D) Strategy proposed herein.

that is triggered by the generation of a silyl cation at the *ortho* position in the presence of an appropriate nucleophile (Figure 1 D). We also anticipated that the resulting products could be transformed into a wide range of difluoromethylene compounds because the hydrosilyl group of the starting material is converted into a base-activatable fluorosilyl group during this transformation.^[11]

After an extensive screening of reaction conditions for the transformation of trifluoromethylarenes by single C-F bond cleavage, we found that defluorinative mono-allylation of 2-(diphenylsilyl)benzotrifluoride (1a)^[12] proceeded upon treatment with a trityl cation source in the presence of allyltrimethylsilane (3a) to afford gem-difluorinated 2a (Table 1). The best result was obtained when a mixture of 1a and 3a was treated with triphenylmethylium tetrafluoroborate in a 1:1 mixture of dichloromethane and 1,1,1,3,3,3hexafluoro-2-propanol (HFIP) at 0°C for 10 minutes, which afforded 2a in 86% yield (entry 6). The diallylated product, which would result from further C-F bond cleavage, was not detected. Product 2a contains a fluorosilyl group, which indicates that the reaction proceeded by hydride abstraction from the silyl group in **1a** with the trityl cation,^[13] followed by migration of a fluoride to the silvl group. Performing the reaction in dichloromethane, toluene, acetonitrile, or nitromethane furnished a complex mixture that included triphenylmethane (entries 1-4). These results indicate that using

© 2016 Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim

Wiley Online Library

These are not the final page numbers!





En	try Solvent		2a [%] ^[a]
1	CH ₂ Cl ₂		trace
2	toluene		0
3	CH₃CN		0
4	CH₃NC	2	0
5	CH ₂ Cl ₂	(CF ₃) ₂ CHOH (9:1)	34
6	CH ₂ Cl ₂	(CF ₃) ₂ CHOH (1:1)	86 ^[b]
7	CH ₂ Cl ₂	(CF ₃) ₂ CHOH (1:9)	58

[a] Yields determined by ¹H NMR analysis, unless otherwise noted. [b] Yield of isolated product.

a solvent such as HFIP that strongly stabilizes cationic intermediates was crucial to the efficiency of this transformation.^[14]

The optimized conditions were applicable to the defluorinative allylation of various benzotrifluorides 1 (Table 2). Benzotrifluorides bearing electron-rich substituents, such as

Table 2: C-F bond allylation of various benzotrifluorides.



[[]a] Allylsilane (10 equiv) was used.

2

www.angewandte.org

© 2016 Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim

These are not the final page numbers!

methoxy, *para*-anisyl, *para*-tolyl, or 3-thienyl groups, were smoothly allylated to afford the corresponding products **2b**– **2e** in high yields. The allylation of substrates bearing electrondeficient groups also afforded difluoromethylated products **2f–2j**, some of which contained chloro, bromo, or ester moieties for further transformation. Notably, as demonstrated by the synthesis of **2i** and **2j**, the C–F bond cleavage of substrates having two trifluoromethyl groups occurred only at the trifluoromethyl group adjacent to the silyl group. 2-(Dimethylsilyl)benzotrifluorides also participated in the reaction, providing the allylated products, such as **2k** and **2l**, in moderate yields.

A variety of substituted allylsilanes, such as **3b–3e**, were also suitable for this transformation (Table 3). Methallylation and 2-bromoallylation afforded the corresponding products

Table 3: C-F bond allylation using various allylsilanes.



2m and **2n** in moderate yields (entries 1 and 2). In the case of γ -substituted allylsilanes **3d** and **3e**, C–C bond formation selectively occurred at the γ -position (entries 3 and 4).^[15] Notably, 4,4-difluoro-1-butene derivative **2p** was obtained by C–F bond cleavage of the trifluoromethyl group of benzotrifluoride **1a** without damaging the trifluoromethyl group of allylsilane **3e**.

The fluorodiphenylsilyl and fluorodimethylsilyl groups in the allylated products are more reactive than the corresponding hydrosilyl groups, thereby allowing for further transformations into a diverse range of difluorinated compounds (Scheme 1). For example, desilylprotonation of **2c** or **2l** with tetra(*n*-butyl)ammonium fluoride (TBAF) smoothly proceeded to afford **4** in high yields. C–C bond formation by Hiyama cross-coupling with an iodoarene using silver(I) oxide as an activator afforded *para*-terphenyl derivative **5**.^[16,17] Modification of the allyl group under Heck arylation conditions in the presence of TBAF afforded desilylprotonated styrene derivative **6**.



Scheme 1. Further transformations of the fluorosilyl group.

To gain insight into the mechanism of the defluorinative allylation reaction, we performed several experiments. The reaction of benzotrifluoride 1 m, which bears a hydrosilyl group at the *para* position, did not afford the desired allylated product 2 q; a complex mixture was obtained, and the formation of triphenylmethane was observed (Scheme 2A). This result indicates the importance of the proximity of the hydrosilyl and trifluoromethyl groups to achieve the transformation, which most likely proceeds by intramolecular fluoride migration. Furthermore, a competition experiment using equimolar amounts of 1 a, 1 j, and a trityl cation source resulted in the preferential production of 2 a over 2 j, indicating that more electron-rich substrates are favored by this reaction (Scheme 2B).

On the basis of the experimental results, we currently believe that the reaction is triggered by hydride abstraction from the hydrosilyl group of benzotrifluoride 1a with a trityl cation, followed by the transfer of one fluoride of the trifluoromethyl group to generate carbocationic intermediate **II** (Scheme 3 A). Subsequent allylation of carbocation **II** with allylsilane 3a would then furnish the desired product 2a by a mechanism similar to that of the Hosomi–Sakurai allylation.





Angew. Chem. Int. Ed. 2016, 55, 1-5

© 2016 Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim

einheim www.angewandte.org
These are not the final page numbers!



Scheme 3. Plausible reaction mechanisms.

However, a concerted mechanism involving hydride abstraction and C–F bond cleavage without generation of silylium intermediate I cannot be excluded (Scheme 3B).

Nucleophiles other than allylsilane were also applicable to the defluorinative transformation (Scheme 4). For example, using benzoic acid 7 or anhydrous hydrogen chloride as the nucleophile, carboxylation and chlorination of benzotrifluoride 1a took place to afford ester 8 and chlorodifluoromethylarene 9, respectively.



Scheme 4. C-F bond carboxylation and chlorination.

In summary, we have described the transformation of a single C–F bond of benzotrifluorides with an *ortho*-hydrosilyl group upon treatment with a trityl cation in the presence of various nucleophiles. Through further transformations of the resulting fluorosilylarenes, this method allows for the synthesis of various difluoromethylene compounds. Further studies to expand the scope of this reaction are currently underway.

Acknowledgements

This work was supported by the JSPS KAKENHI Grants 24310164 (T.H.) and 26350971 (S.Y.) and the Platform for Drug Discovery, Informatics, and Structural Life Science from MEXT and AMED, Japan.



Keywords: allylation \cdot C–F Activation \cdot fluorine \cdot silanes \cdot trityl cation

- a) S. Swallow in *Fluorine in Pharmaceutical and Medicinal Chemistry: From Biophysical Aspects to Clinical Applications* (Eds.: V. Gouverneur, K. Müller), Imperial College Press, London, **2012**, pp. 141–174; b) K. Müller, C. Faeh, F. Diederich, *Science* **2007**, *317*, 1881–1886; c) S. Purser, P. R. Moore, S. Swallow, V. Gouverneur, *Chem. Soc. Rev.* **2008**, *37*, 320–330; d) J. Wang, M. Sánchez-Roselló, J. L. Aceña, C. del Pozo, A. E. Sorochinsky, S. Fustero, V. A. Soloshonok, H. Liu, *Chem. Rev.* **2014**, *114*, 2432–2506; e) E. P. Gillis, K. J. Eastman, M. D. Hill, D. J. Donnelly, N. A. Meanwell, *J. Med. Chem.* **2015**, *58*, 8315–8359.
- [2] a) T. Liang, C. Neumann, T. Ritter, Angew. Chem. Int. Ed. 2013, 52, 8214–8264; Angew. Chem. 2013, 125, 8372–8423; b) M. G. Campbell, T. Ritter, Chem. Rev. 2015, 115, 612–633.
- [3] a) J. Charpentier, N. Früh, A. Togni, *Chem. Rev.* 2015, *115*, 650–682; b) X. Liu, C. Xu, M. Wang, Q. Liu, *Chem. Rev.* 2015, *115*, 683–730.
- [4] F. A. Carey, R. J. Sandberg in Advanced Organic Chemistry Part A: Structure and Mechanisms, 5th Ed., Springer, New York, 2007, p. 258.
- [5] D. M. Lemal, J. Org. Chem. 2004, 69, 1-11.
- [6] a) W. D. Jones, *Dalton Trans.* 2003, 3991–3995; b) H. Amii, K. Uneyama, *Chem. Rev.* 2009, 109, 2119–2183; c) R. P. Hughes, *Eur. J. Inorg. Chem.* 2009, 4591–4606; d) T. Stahl, H. F. T. Klare, M. Oestreich, *ACS Catal.* 2013, *3*, 1578–1587; e) T. Ahrens, J. Kohlmann, M. Ahrens, T. Braun, *Chem. Rev.* 2015, 115, 931–972.
- [7] For selected examples, see: a) R. Panisch, M. Bolte, T. Müller, J. Am. Chem. Soc. 2006, 128, 9676–9682; b) M. Ohashi, T. Kambara, T. Hatanaka, H. Saijo, R. Doi, S. Ogoshi, J. Am. Chem. Soc. 2011, 133, 3256–3259; c) O. Allemann, S. Duttwyler, P. Romanato, K. K. Baldridge, J. S. Siegel, Science 2011, 332, 574–577; d) C. B. Caputo, D. W. Stephan, Organometallics 2012, 31, 27–30; e) X.-W. Liu, J. Echavarren, C. Zarate, R. Martin, J. Am. Chem. Soc. 2015, 137, 12470–12473; f) T. Niwa, H. Ochiai, Y. Watanabe, T. Hosoya, J. Am. Chem. Soc. 2015, 137, 14313–14318; g) P. Tian, C. Feng, T.-P. Loh, Nat. Commun. 2015, 6, 7472; h) T. Ichitsuka, T. Fujita, J. Ichikawa, ACS Catal. 2015, 5, 5947–5950.

- [8] a) V. J. Scott, R. Çelenligil-Çetin, O. V. Ozerov, J. Am. Chem. Soc. 2005, 127, 2852–2853; b) C. Douvris, O. V. Ozerov, Science 2008, 321, 1188–1190; c) T. Stahl, H. F. T. Klare, M. Oestreich, J. Am. Chem. Soc. 2013, 135, 1248–1251.
- [9] a) K. Fuchibe, T. Akiyama, J. Am. Chem. Soc. 2006, 128, 1434–1435; b) J. Terao, M. Nakamura, N. Kambe, Chem. Commun. 2009, 6011–6013; c) T. Akiyama, K. Atobe, M. Shibata, K. Mori, J. Fluorine Chem. 2013, 152, 81–83.
- [10] For examples of the successful transformation of a single C–F bond of trifluoromethylarenes, see: a) D. Ferraris, C. Cox, R. Anand, T. Lectka, *J. Am. Chem. Soc.* 1997, *119*, 4319–4320; b) H. Amii, Y. Hatamoto, M. Seo, K. Uneyama, *J. Org. Chem.* 2001, *66*, 7216–7218; c) H. Dang, A. M. Whittaker, G. Lalic, *Chem. Sci.* 2016, *7*, 505–509; d) R. Doi, M. Ohashi, S. Ogoshi, *Angew. Chem. Int. Ed.* 2016, *55*, 341–344; *Angew. Chem.* 2016, *128*, 349–352.
- [11] a) Y. Hatanaka, T. Hiyama, J. Org. Chem. 1989, 54, 268–270;
 b) H. F. Sore, W. R. J. D. Galloway, D. R. Spring, Chem. Soc. Rev. 2012, 41, 1845–1866.
- [12] The benzotrifluorides 1 with an *ortho*-hydrosilyl group that were used in this study were easily prepared from 2-iodobenzotrifluorides. Arylated substrates were prepared by Suzuki–Miyaura cross-couplings of the corresponding bromo-substituted benzotrifluorides with aryl boronic acids. See the Supporting Information for details.
- [13] a) J. Y. Corey, J. Am. Chem. Soc. 1975, 97, 3237–3238;
 b) G. K. S. Prakash, C. Bae, G. Rasul, G. A. Olah, J. Org. Chem. 2002, 67, 1297–1301.
- [14] The product that would result from nucleophilic addition of HFIP to the carbocationic intermediate was not detected, probably owing to the low nucleophilicity of HFIP; see: a) J. Ichikawa, S. Miyazaki, M. Fujiwara, T. Minami, J. Org. Chem. 1995, 60, 2320–2321; b) J.-P. Bégué, D. Bonnet-Delpon, Synlett 2004, 18–29; c) T. Dohi, N. Yamaoka, Y. Kita, Tetrahedron 2010, 66, 5775–5785.
- [15] A. Hosomi, M. Endo, H. Sakurai, Chem. Lett. 1976, 5, 941-942.
- [16] K. Hirabayashi, J. Kawashima, Y. Nishihara, A. Mori, T. Hiyama, Org. Lett. 1999, 1, 299–302.
- [17] Hiyama cross-coupling with 2-diphenylsilyl-4-(4-methoxyphenyl)benzotrifluoride instead of 2c under the same conditions did not afford 5.

Received: May 16, 2016 Published online: ■■ ■■, ■■■

www.angewandte.org

These are not the final page numbers!



Communications

Communications

C-F Activation

S. Yoshida,* K. Shimomori, Y. Kim, T. Hosoya* _____ IIII--IIII

Single C-F Bond Cleavage of Trifluoromethylarenes with an *ortho*-Silyl Group



One of the three: The transformation of a single C–F bond of trifluoromethylarenes bearing an *ortho*-hydrosilyl group is described. The activation of the hydrosilyl group with a trityl cation in the presence of nucleophiles enabled selective C–F bond functionalization, for example, by allylation, carboxylation, or chlorination. Further derivatization of the resulting fluorosilylarenes afforded various difluoromethylated arenes.

