


Synthesis and Characterization of an Isolable Base-Stabilized Silacycloprop-1-ylidene**

Ricardo Rodriguez, Thibault Troadec, Tsuyoshi Kato,* Nathalie Saffon-Merceron, Jean-Marc Sotiropoulos, and Antoine Baceiredo*


The chemistry of stable silylenes has been the subject of considerable research since the isolation of silicocene^[1] and N-heterocyclic silylene.^[2] To date, several stable silylenes,^[3–5] including kinetically stabilized dialkylsilylene,^[6] have been synthesized and their chemical properties have been extensively investigated.^[7] A novel silylene stabilized by ylidic carbon centers has also recently been reported.^[8] However, the structural modifications of stable silylenes are still far more limited than those of their lighter analogues, the carbenes.^[9]



Recently, silylenes **I** stabilized by the coordination of a donating ligand to the silicon(II) atom have attracted much attention owing to their high reactivity, which is strongly related to the nature of the ligands.^[10–13] In particular, this stabilization method allows for easy functionalization of the Si(II) atom. Indeed, silylenes with reactive Si–X bonds **II** (X = Cl, Br)^[11,14] and even dihalogenosilylene complexes **III**^[12] have been isolated as stable crystalline materials and have been used as precursors for preparing unique molecules.^[14,15] We have also successfully synthesized the first base-stabilized silicon(II) hydride species **IV**,^[16] which are difficult to prepare in the absence of steric protection.^[17] In this series, the small highly strained cyclic silylenes, such as the silacycloprop-1-ylidenes **V**, remain challenging target molecules.^[18]

Herein, we report the first synthesis of a base-stabilized silacycloprop-1-ylidene **3**.

We recently reported the synthesis of the first phosphine-stabilized silicon(II) hydride **1** and studied the reactivity of the species with olefins.^[16] Compound **1** similarly reacts at room temperature with diphenylacetylene through a [2+1] cycloaddition reaction to give the corresponding pentacoordinate silirene **2**, which was fully characterized by NMR spectroscopy in solution. A ²⁹Si NMR analysis of silirene **2** shows a high-field doublet (–135.2 ppm, ¹J_{SiP} = 58.9 Hz), which is characteristic of this type of structure.^[19] In the ¹H NMR spectrum, the Si–H fragment appears as a doublet at 6.25 ppm (²J_{PH} = 96.4 Hz) with the characteristic ²⁹Si satellites (ca. 4.7%, ¹J_{HSi} = 286.1 Hz), confirming a direct silicon–hydrogen bond in **2** (Scheme 1).



Scheme 1. Synthesis of base-stabilized silacycloprop-1-ylidene **3**.

Silirene **2** is thermally labile and quantitatively isomerizes in toluene at 80 °C into silacycloprop-1-ylidene **3**, in which the Si^{II} atom is stabilized by the intramolecular coordination of the imine fragment. Compound **3** was successfully isolated as colorless crystals in 92% yield from pentane at room temperature (Scheme 1). Although the mechanism of the reaction is still unclear, this transformation formally involves two 1,2-shifts of silicon ligands, hydride and phosphine, to each of the carbon atoms of the silirene ring. Contrary to the previous cases, neither an alkyne-insertion into the Si^{II}–H bond^[15] nor tricyclic phosphine formation^[19] was observed. In the ³¹P NMR spectrum, **3** displays two singlets (46.5 and 42.8 ppm), which is in agreement with the presence of two diastereomers (92:8). In the ²⁹Si NMR spectrum, two high-field doublets were observed in the same ratio (–87.5 ppm, ²J_{SiP} = 2.9 Hz, for the major one), which is in agreement with a silicon atom in a small three-membered ring. The cyclopropylidene ring proton appears as doublets (3.57 ppm, ³J_{PH} = 30.8 Hz, and 3.02 ppm, ³J_{PH} = 31.1 Hz) in the ¹H NMR spectrum.

The crystal structure of **3** reveals a cyclopropylidene ring featuring a divalent silicon atom complexed by the imine

[*] Dr. R. Rodriguez, T. Troadec, Dr. T. Kato, Dr. A. Baceiredo
Université de Toulouse, UPS, and CNRS, LHFA
118 route de Narbonne, 31062 Toulouse (France)
E-mail: kato@chimie.ups-tlse.fr
baceired@chimie.ups-tlse.fr
Homepage: http://hfa.ups-tlse.fr

Dr. N. Saffon-Merceron
Université de Toulouse, UPS, and CNRS, ICT
118 route de Narbonne, 31062 Toulouse (France)

Dr. J.-M. Sotiropoulos
Université de Pau et des Pays de l'Adour, IPREM
Technopôle Hélioparc
2 avenue du Président Angot, 64053 Pau (France)

[**] We are grateful to the CNRS and the ANR (NOPROBLEM, LEGO) for financial support of this work.

Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/anie.201201581>.

fragment (Figure 1).^[20] The three-membered ring has a small bond angle around the silicon center (C1–Si1–C2 43.33°) and the two Si–C bonds (Si1–C1 1.993 Å, Si1–C2 1.920 Å) are




Figure 1. Crystal structure of **3**. Thermal ellipsoids set at 30% probability. H atoms omitted for clarity. Selected bond lengths [Å] and angles [°]: Si1–C1 1.993(3), Si1–C2 1.920(3), C1–C2 1.572(3), Si1–N1 1.857(2), C1–P1 1.763(2), P1–C3 1.728(3), C3–C4 1.382(3), C4–N1 1.343(3); C1–Si1–C2 47.33(10), Si1–C1–C2 63.90(13), Si1–C2–C1 68.77(13), N1–Si1–C1 102.02(10), N1–Si1–C2 98.96(10).

longer than usual for Si–C bonds (ca. 1.89 Å).^[21] The silicon–nitrogen distance is also quite long (Si–N1 1.857 Å). The long bond distances around the tri-coordinate silicon center can probably be rationalized by the enhanced p character of these bonds as a consequence of the strong pyramidalization of the silicon center ($\Sigma^\circ_{\text{Si}} = 248.3^\circ$). The P–C bond (P1–C1 1.763 Å) is on the short end of the range expected for phosphoniocyclopropanes (1.759–1.780 Å).^[22] The geometry of the α -iminophosphorus ylide fragment (P1–C1–C2–N1) complexing the Si(II) center is similar to that observed in the corresponding transition-metal complexes.^[23]

To obtain more information about **3**, calculations were performed at the B3LYP/6-31G(d,p) level of theory (Figure 2). The optimized structure of **3** agrees quite well with the experimental data.^[24] The HOMO of **3** corresponds mainly to the lone pair orbital on the silicon atom. Natural bonding orbital (NBO) analysis reveals a very high level of s character for the nonbonding orbital localized on the silicon




Figure 2. Calculated molecular orbitals of **3** at a 0.05 cut-off level. Energy levels of orbitals are given in parentheses.

atom (s: 75.4%, p: 24.6%). Despite this result, the energy level of the HOMO (–4.49 eV) remains quite high, suggesting a strong nucleophilic character. The LUMO does not localize on the Si^{II} atom, but rather corresponds to the π conjugate system of the α -iminophosphorus ylide fragment. The vacant orbital of the Si^{II} atom is much higher in energy (LUMO + 7), resulting in a large energy gap between LUMO + 7 and HOMO (5.45 eV), which suggests a decreasing silylenoid character for **3**.

In contrast to the phosphine-stabilized silylene **1**, which shows a silylene-like reactivity,^[16,19,25] **3** does not react with olefins or alkynes. However, it readily reacts with the Lewis acid BH₃·THF to give the corresponding borane complex **4**, which was isolated as colorless crystals in 93% yield. The crystal structure of **4** reveals the coordination of BH₃ to the Si^{II} atom^[26] (Figure 3).^[20] The three bonds around the silicon





Figure 3. Crystal structure of **4**. Thermal ellipsoids set at 30% probability. H atoms (except for H atoms on B1) are omitted for clarity. Selected bond lengths [Å] and angles [°]: Si1–B1 1.987(2), Si1–C1 1.932(2), Si1–C2 1.868(2), C1–C2 1.610(2), Si1–N1 1.796(2), C1–P1 1.778(2), P1–C3 1.733(2), C3–C4 1.370(2), C4–N1 1.363(2); C1–Si1–C2 50.11(7), C2–C1–Si1 62.89(9), C1–C2–Si1 67.00(9), N1–Si1–C1 107.32(7), N1–Si1–C2 106.93(7).

atom are significantly shorter (Si1–C1: 1.932 Å, Si1–C2: 1.868 Å, Si1–N1: 1.796 Å) than those in **3**. These shorter bond lengths are probably due to the reduced p character of these bonds, which is induced by the coordination of the lone pair. Indeed, the sum of the bond angles for the SiX₃ subunit (X = N1, C1, and C2) in **4** (264.4°) is much larger than that observed for **3** (248.3°). Silacycloprop-1-ylidene **3** also reacts with polarized unsaturated molecules, such as ethyl vinyl ketone, to give spirocyclic silane **5** in 92% yield (Scheme 2). The reaction probably involves an initial [4+1] cycloaddition reaction at the Si^{II} atom of **3** followed by a ring opening reaction of the three-membered silacyclopropane ring to afford **5**. This system features a bicyclic structure with a novel seven-membered ring that includes a phosphonium ylide fragment. The structure of this molecule was unambiguously confirmed by X-ray diffraction analysis.^[20,27]



Scheme 2. Reaction of **3** with $\text{BH}_3\cdot\text{THF}$ and ethyl vinyl ketone.

The isomerization of silirene **2** into silacycloprop-1-ylidene **3** involves a reduction of the silicon center from Si^{IV} to Si^{II} (Figure 4). Such an isomerization to give a stable lower-




Figure 4. Relative energies (kcal mol^{-1}) of isomers **2** and **3**.

valent silicon species is, to the best of our knowledge, unprecedented. Calculations indicate that isomers **2** and **3** are very close in energy ($\Delta E_{3,2} = +3.2 \text{ kcal mol}^{-1}$). This is probably due to the efficient stabilization of the Si^{II} center in **3** by coordination of the iminophosphorus ylide fragment.

In conclusion, we have successfully synthesized an N-donor-stabilized silacycloprop-1-ylidene **3** through the isomerization of tetravalent silirene **2** under mild conditions. This process involves an unprecedented $\text{Si}^{\text{IV}} \rightarrow \text{Si}^{\text{II}}$ rearrangement. Further investigations on the reactivity of this new species are underway.

Received: February 27, 2012

Published online: ■■■ ■■■, ■■■■■

Keywords: complexes · phosphorus · silicon · silylenes · small ring systems

- [1] P. Jutzi, D. Kanne, C. Krüger, *Angew. Chem.* **1986**, *98*, 163; *Angew. Chem. Int. Ed. Engl.* **1986**, *25*, 164.
- [2] M. Denk, R. Lennon, R. Hayashi, R. West, A. V. Belyakov, H. P. Verne, A. Haaland, M. Wagner, N. Metzler, *J. Am. Chem. Soc.* **1994**, *116*, 2691.
- [3] a) R. West, M. Denk, *Pure Appl. Chem.* **1996**, *68*, 785; b) W. Li, N. J. Hill, A. C. Tomasik, G. Bikzhanova, R. West, *Organometallics* **2006**, *25*, 3802; c) A. C. Tomasik, A. Mitra, R. West, *Organometallics* **2009**, *28*, 378.
- [4] a) B. Gehrus, M. F. Lappert, J. Heinicke, R. Boese, D. Bläser, *J. Chem. Soc. Chem. Commun.* **1995**, 1931; b) J. Heinicke, A. Oprea, M. K. Kindermann, T. Karpati, L. Nyulaszi, T. Veszpremi, *Chem. Eur. J.* **1998**, *4*, 541; c) B. Gehrus, P. B. Hitchcock, M. F. Lappert, *Z. Anorg. Allg. Chem.* **2005**, *631*, 1383.
- [5] M. Driess, S. Yao, M. Brym, C. van Wüllen, D. Lentz, *J. Am. Chem. Soc.* **2006**, *128*, 9628.
- [6] M. Kira, S. Ishida, T. Iwamoto, C. Kabuto, *J. Am. Chem. Soc.* **1999**, *121*, 9722.
- [7] For reviews on stable silylenes and their reactivities, see: a) M. Asay, C. Jones, M. Driess, *Chem. Rev.* **2011**, *111*, 354; b) S. Yao, Y. Xiong, M. Driess, *Organometallics* **2011**, *30*, 1748; c) M. Kira, T. Iwamoto, S. Ishida, *Bull. Chem. Soc. Jpn.* **2007**, *80*, 258; d) N. Takeda, N. Tokitoh, *Synlett* **2007**, 2483; e) H. Ottosson, P. G. Steel, *Eur. J. Inorg. Chem.* **2006**, 1577; f) N. J. Hill, R. West, *J. Organomet. Chem.* **2004**, *689*, 4165; g) M. Kira, *J. Organomet. Chem.* **2004**, *689*, 4475; h) M. Okazaki, H. Tobita, H. Ogino, *Dalton Trans.* **2003**, 493; i) B. Gehrus, M. F. Lappert, *J. Organomet. Chem.* **2001**, *617*, 209; j) M. Haaf, T. A. Schmedake, R. West, *Acc. Chem. Res.* **2000**, *33*, 704.
- [8] M. Asay, S. Inoue, M. Driess, *Angew. Chem.* **2011**, *123*, 9763; *Angew. Chem. Int. Ed.* **2011**, *50*, 9589.
- [9] a) M. Melaimi, M. Soleilhavoup, G. Bertrand, *Angew. Chem.* **2010**, *122*, 8992; *Angew. Chem. Int. Ed.* **2010**, *49*, 8810; b) W. Kirmse, *Angew. Chem.* **2004**, *116*, 1799; *Angew. Chem. Int. Ed.* **2004**, *43*, 1767; c) J. Vignolle, X. Cattoën, D. Bourissou, *Chem. Rev.* **2009**, *109*, 3333; d) D. Bourissou, O. Guerret, F. P. Gabbai, G. Bertrand, *Chem. Rev.* **2000**, *100*, 39; e) T. Kato, E. Maerten, A. Baceiredo, in *Transition Metal Complexes of Neutral η^1 -Carbon Ligands*, Vol. 30, XI ed. (Eds.: R. Chauvin, Y. Canac), Springer, Berlin, **2010**, p. 131.
- [10] D. Gau, T. Kato, N. Saffon-Merceron, F. P. Cossio, A. Baceiredo, *J. Am. Chem. Soc.* **2009**, *131*, 8762.
- [11] a) C. W. So, H. W. Roesky, J. Magull, E. B. Oswald, *Angew. Chem.* **2006**, *118*, 4052; *Angew. Chem. Int. Ed.* **2006**, *45*, 3948; b) C. W. So, H. W. Roesky, P. M. Gurubasavaraj, R. B. Oswald, M. T. Gamer, P. G. Jones, S. Blaurock, *J. Am. Chem. Soc.* **2007**, *129*, 12049.
- [12] a) R. S. Ghadwal, H. W. Roesky, S. Merkel, J. Henn, D. Stalke, *Angew. Chem.* **2009**, *121*, 5793; *Angew. Chem. Int. Ed.* **2009**, *48*, 5683; b) A. C. Filippou, O. Chernov, G. Schnakenburg, *Angew. Chem.* **2009**, *121*, 5797; *Angew. Chem. Int. Ed.* **2009**, *48*, 5687.
- [13] a) N. Takeda, H. Suzuki, N. Tokitoh, R. Okazaki, *J. Am. Chem. Soc.* **1997**, *119*, 1456; b) N. Takeda, T. Kajiwara, H. Suzuki, R. Okazaki, N. Tokitoh, *Chem. Eur. J.* **2003**, *9*, 3530.
- [14] D. Gau, T. Kato, N. Saffon-Merceron, F. P. Cossio, A. Baceiredo, *Angew. Chem.* **2010**, *122*, 6735; *Angew. Chem. Int. Ed.* **2010**, *49*, 6585.
- [15] a) H. Tanaka, S. Inoue, M. Ichinohe, M. Driess, A. Sekiguchi, *Organometallics* **2011**, *30*, 3475; b) R. S. Ghadwal, H. W. Roesky, K. Pröpper, B. Dittrich, S. Klein, G. Frenking, *Angew. Chem.* **2011**, *123*, 5486; *Angew. Chem. Int. Ed.* **2011**, *50*, 5374; c) S. M. I. Al-Rafia, A. C. Malcolm, R. McDonald, M. J. Ferguson, E. Rivard, *Angew. Chem.* **2011**, *123*, 8504; *Angew. Chem. Int. Ed.* **2011**, *50*, 8354.
- [16] R. Rodriguez, D. Gau, Y. Contie, T. Kato, N. Saffon-Merceron, A. Baceiredo, *Angew. Chem.* **2011**, *123*, 11694; *Angew. Chem. Int. Ed.* **2011**, *50*, 11492.
- [17] S.-H. Zhang, H.-X. Yeong, H.-W. Xi, K. H. Lim, C.-W. So, *Chem. Eur. J.* **2010**, *16*, 10250.
- [18] Characterization of sila-cyclopropylidene by IR spectroscopy in an Ar matrix at 10 K: G. Maier, H. P. Reisenauer, H. Egenolf, *Eur. J. Org. Chem.* **1998**, 1313.
- [19] D. Gau, R. Rodriguez, T. Kato, N. Saffon-Merceron, A. Baceiredo, *J. Am. Chem. Soc.* **2010**, *132*, 12841.
- [20] CCDC 868349 (**3**), 868350 (**4**), and 868351 (**5**) contain the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif.
- [21] M. Kaftory, M. Kapon, M. Botoshansky, *The Chemistry of Organic Silicon Compounds*, Vol. 2 (Eds.: Z. Rappoport, Y. Apeloig), Wiley, Chichester, **1998**, p. 181.
- [22] a) C. Puke, G. Erker, B. Wibbeling, R. Fröhlich, *Eur. J. Org. Chem.* **1999**, 1831; b) C. P. Casey, S. Kraft, D. R. Powell, M. Kavana, *J. Organomet. Chem.* **2001**, *617*, 723; c) U. Kleinitz, R. Mattes, *Chem. Ber.* **1994**, *127*, 605.

- [23] P. Braunstein, J. Pietsch, Y. Chauvin, S. Mercier, L. Saussine, A. DeCian, J. Fischer, *J. Chem. Soc. Dalton Trans.* **1996**, 3571.
- [24] Selected bond lengths [Å] and angles [°] calculated at B3LYP/6-31G(d,p) for **3**: Si1–C1 2.035, Si1–C2 1.946, Si1–N1 1.905, C1–C2 1.564; C1–Si1–C2 46.20, N1–Si1–C1 101.92, N1–Si1–C2 98.74. For more computational details, see the Supporting Information.
- [25] a) D. Gau, R. Rodriguez, T. Kato, N. Saffon-Merceron, F. P. Cossío, A. Baceiredo, *Chem. Eur. J.* **2010**, *16*, 8255; b) R. Rodriguez, D. Gau, T. Kato, N. Saffon-Merceron, A. De Cózar, F. P. Cossío, A. Baceiredo, *Angew. Chem.* **2011**, *123*, 10598; *Angew. Chem. Int. Ed.* **2011**, *50*, 10414.
- [26] BH₃ complexed silicon(II) species: a) M. Y. Abraham, Y. Wang, Y. Xie, P. Wei, H. F. Schaefer III, P. von R. Schleyer, G. H. Robinson, *J. Am. Chem. Soc.* **2011**, *133*, 8874; b) A. Jana, D. Leusser, I. Objartel, H. W. Roesky, D. Stalke, *Dalton Trans.* **2011**, *40*, 5458.
- [27] See the Supporting Information.
-


Communications



Silicon Chemistry

R. Rodriguez, T. Troadec, T. Kato,*
N. Saffon-Merceron, J.-M. Sotiropoulos,
A. Baceiredo* ————— ■■■■—■■■■

Synthesis and Characterization of an
Isolable Base-Stabilized Silacycloprop-1-
ylidene



Strained but stable: An isolable silacycloprop-1-ylidene stabilized by intramolecular complexation with an iminophosphorus ylide fragment was successfully synthesized and fully characterized. The formation of this small highly strained cyclic silylene involves an unprecedented $\text{Si}^{\text{IV}} \rightarrow \text{Si}^{\text{II}}$ rearrangement under very mild conditions.