This article was downloaded by: [University of Chicago Library] On: 10 October 2014, At: 15:51 Publisher: Taylor & Francis Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



# Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information: <u>http://www.tandfonline.com/loi/gpss20</u>

Synthesis and Optical Properties of a Bis(Benzothiaborino)Carbazole, a Thiaborin-Carbazole Mixed Ladder-Type π-Conjugated Molecule

Tomohiro Agou<sup>a</sup>, Junji Kobayashi<sup>a</sup> & Takayuki Kawashima<sup>a</sup> <sup>a</sup> Department of Chemistry, Graduate School of Science, The University of Tokyo, Tokyo, Japan Published online: 27 May 2010.

To cite this article: Tomohiro Agou , Junji Kobayashi & Takayuki Kawashima (2010) Synthesis and Optical Properties of a Bis(Benzothiaborino)Carbazole, a Thiaborin-Carbazole Mixed Ladder-Type  $\pi$ -Conjugated Molecule, Phosphorus, Sulfur, and Silicon and the Related Elements, 185:5-6, 947-951, DOI: <u>10.1080/10426501003769736</u>

To link to this article: http://dx.doi.org/10.1080/10426501003769736

# PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms &

Conditions of access and use can be found at <u>http://www.tandfonline.com/page/terms-and-conditions</u>



Phosphorus, Sulfur, and Silicon, 185:947–951, 2010 Copyright © Taylor & Francis Group, LLC ISSN: 1042-6507 print / 1563-5325 online DOI: 10.1080/10426501003769736

# SYNTHESIS AND OPTICAL PROPERTIES OF A BIS(BENZOTHIABORINO)CARBAZOLE, A THIABORIN-CARBAZOLE MIXED LADDER-TYPE $\pi$ -CONJUGATED MOLECULE

**Tomohiro Agou, Junji Kobayashi, and Takayuki Kawashima** Department of Chemistry, Graduate School of Science, The University of Tolyo

Department of Chemistry, Graduate School of Science, The University of Tokyo, Tokyo, Japan

A ladder-type  $\pi$ -conjugated molecule containing both thiaborin and carbazole units exhibited strong light absorption and emission like other ladder-type  $\pi$ -conjugated molecules based on heteraborins.

Keywords Carbazole; ladder-type  $\pi$ -conjugated molecule; optical properties; thiaborin

#### INTRODUCTION

Ladder-type  $\pi$ -conjugated compounds are promising candidates for organic functional materials, including electron- or hole-transporting materials and light emitting materials. Their planar structures can enhance delocalization of  $\pi$ -orbitals and decrease HOMO–LUMO energy gaps, resulting in high electron or hole affinity. In addition, as the  $\pi$ -conjugated structures are fixed tightly by the rigid framework, intramolecular motions that are related to nonradiation decay of the excited state are suppressed, and thus photoluminescence properties of the ladder-type compounds are improved.<sup>1</sup> Recently, the introduction of various main group elements into ladder-type conjugated molecules proved to be a very effective way to develop new characteristics of  $\pi$ -conjugated molecules.<sup>2</sup>

In previous reports, we have described the optical properties of ladder-type azaborines and thiaborins.<sup>3</sup> However, their analogues containing both azaborine and thiaborin units have not been synthesized yet, and thus the effect of the mixing of these two different heteraborins in one molecule is still unclear. In this article, we report the attempted synthesis of a mixed-type, ladder-type heteraborin bearing azaborine and thiaborin units in a hexahydroheptacene framework, which resulted in the formation of a bis(benzothiaborino)carbazole instead of the expected ladder-type heteraborin (Figure 1).

Dedicated to Professor Naomichi Furukawa on the occasion of his 70th birthday.

Address correspondence to Takayuki Kawashima, Department of Chemistry, Graduate School of Science, The University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-0033, Japan. E-mail: takayuki@chem.s.utokyo.ac.jp

Received 9 November 2008; accepted 20 December 2008.

This work was partly supported by Grants-in-Aid for The Global COE program for Chemical Innovation (T.K.) and for Scientific Research (T.K.) from MEXT. We also thank Tosoh Finechem Corporation for the generous gifts of alkyllithium compounds.



Figure 1 Ladder-type mixed heteraborin 1 and bis(benzothiaborino)carbazole 2.

# **RESULTS AND DISCUSSION**

#### Synthesis

The synthesis of hexabromide **6** was accomplished by taking advantage of Buchwald–Hartwig amination protocols (Scheme 1).<sup>4,5</sup>

The synthesis of mixed ladder-type heteraborin **1** was attempted. However, instead of **1**, bis(benzothiaborino)carbazole **2** was obtained. A similar reaction using the corresponding sulfide gave bis(benzothiaborino)dibenzothiophene **7**, a sulfur analogue of **2**, together with the desired **8** (Figure 2).<sup>3a</sup> On the other hand, in the case of the synthesis of ladder-type azaborines, carbazole derivative was not obtained at all.<sup>3b,3c</sup> Therefore, terminal thiaborins formed by a twofold ring closure should be essential for these oxidative C–C coupling reactions. In fact, an electron transfer from lithio-derivatives to the thiaborin moiety may occur to give **2** or **7**, because the thiaborin has a LUMO of lower energy compared to that of the azaborine, as revealed by theoretical calculations on heteraborins by us,<sup>3a</sup> and seems to act as a one-electron oxidant (Scheme 2).<sup>6</sup>

# **Optical Properties**

The optical data for 2 as well as for other thiaborins are summarized in Table I.

The absorption maximum at longest wavelength of 2 lies between those of 7 and 8. The electronic effect of the insertion of a carbazole moiety is weaker as compared with that of the insertion of a dibenzothiaborin ring. Compared to 7, compound 2 showed red-shifted absorption maxima, probably due to the higher electron-donating ability of the nitrogen atom.



Scheme 1 The synthesis of hexabromide 6.



Scheme 2 The formation of bis(benzothiaborino)carbazole 2.

Compound 2 exhibited green-colored photoluminescence with small Stokes shift, indicating that it maintains a planar and rigid framework upon photo-excitation, just like the other heteraborin-based ladder-type molecules. In addition, compound 2 can emit a green-colored luminescence also in the solid state due to the inhibition of intermolecular interaction by bulky mesityl groups, and thus a carbazole-thiaborin mixed ladder-type structure will be useful to develop strong light-emissive materials.

#### **EXPERIMENTAL**

Generally chemicals were used as received. For optical measurements, commercial spectrochemical- or fluorometric-grade solvents were used as received. All manipulations were performed under argon atmosphere using standard Schlenck technique. Dry solvents (THF, Et<sub>2</sub>O, hexane, and toluene) were purchased from Kanto Chemicals and further purified by an MBRAUN MB-SPS system equipped with activated alumina and molecular sieve columns before use. Column chromatography was carried out with Kanto Silica Gel 60N. Gel permeation liquid chromatography was performed using Japan Analytical Industry LC-918 and LC-908 with JAIGEL 1H+2H columns and chloroform as a solvent. <sup>1</sup>H and <sup>13</sup>C NMR spectra were recorded with a Bruker DRX-500 spectrometer. Low resolution mass spectra were measured with a JEOL JMS-700P (FAB). High resolution mass spectra were obtained with a JEOL JMS-700P. UV-vis spectra were recorded with a JASCO V-670



Figure 2 Thiaborin and ladder-type thiaborins.

	$\lambda_{\text{max}}/\text{nm} (\log \varepsilon)$	$\lambda_{em}/nm$	Stokes shift <sup>c</sup> /cm <sup>-1</sup>
2	521 (3.79)	540	$6.8 \times 10^{2}$
$7^{b}$	482 (3.60)	495	$5.4 \times 10^{2}$
<b>8</b> <sup>a</sup>	558 (3.95)	580	$6.8 \times 10^{2}$
<b>9</b> <sup>a</sup>	387 (3.94)	403	$1.0 \times 10^{3}$
<b>10</b> <sup><i>a</i></sup>	499 (4.28)	517	$7.0 \times 10^2$

Table I Optical properties of bis(benzothiaborino)carbazole 2 and ladder-type thiaborins 7–10 in cyclohexane

<sup>a</sup>Ref. 1.

<sup>b</sup>Unpublished result.

<sup>c</sup>Stokes shift =  $1/\lambda_{max}$ — $1/\lambda_{em}$ .

spectrophotometer, and fluorescence spectra were measured with a JASCO FP-6500 fluorescence spectrophotometer. All melting points were determined on a Yanaco MP-S3 micro melting point apparatus and are uncorrected. Elemental analyses were performed by the Microanalytical Laboratory of the Department of Chemistry, Faculty of Science, University of Tokyo. The preparation of (2-bromo-4-butylphenyl) (2,5-dibromo-4-iodophenyl) sulfide (3) was already reported.<sup>3a</sup>

#### Synthesis of 2,5-Dibromo-4-(2-bromo-4-butylphenylthio)aniline (4)

A mixture of 3(2.0 g, 3.3 mmol),  $Ph_2C = NH(0.61 \text{ mL}, 3.6 \text{ mmol})$ ,  $Pd(dba)_2(0.10 \text{ g}, 3.3 \text{ mmol})$ 0.17 mmol), 1,1'-bis(diphenylphophino)ferrocene (DPPF) (0.10 g, 0.17 mmol), t-BuONa (0.50 g, 5.2 mmol), and toluene (50 mL) was stirred at 100°C for 45 h.<sup>4</sup> To the mixture, aq. NH<sub>4</sub>Cl was added, and the mixture was extracted with Et<sub>2</sub>O. The organic layer was dried over  $Na_2SO_4$ , and the solvents were removed under reduced pressure. The crude material was dissolved in THF (10 mL) and MeOH (40 mL), and to this mixture  $NH_2OH \cdot HCl$  (5.0 g, 72 mmol) and AcONa (9.0 g, 0.11 mol) were added. The mixture was stirred at room temperature for 19 h, and the solvent was removed under reduced pressure. The resulting material was extracted with Et<sub>2</sub>O and the organic layer was dried over Na<sub>2</sub>SO<sub>4</sub>. The solvent was evaporated, and the crude material was chromatographed  $(Al_2O_3, CHCl_3)$  and further purified by GPLC to give 4 as colorless viscous oil (1.3 g, 82%). 4: colorless oil. <sup>1</sup>H NMR  $(500 \text{ MHz}, \text{CDCl}_3)$ :  $\delta = 0.90$  (t, J = 7.3 Hz, 3H), 1.32 (sext, J = 7.5 Hz, 2H), 1.55 (quint, J = 7.7 Hz, 2H), 2.53 (t, J = 7.7 Hz, 2H), 6.72 (d, J = 8.1 Hz, 1H), 6.98 (dd, J = 8.1, 1.6 Hz, 1H), 7.18 (s, 1H), 7.38 (d, J = 1.6 Hz, 1H), 7.49 (s, 1H); <sup>13</sup>C NMR (126 MHz,  $CDCl_3$ ):  $\delta = 13.9, 22.2, 33.2, 34.7, 119.3, 121.5, 122.0, 128.1, 129.4, 129.5, 129.7, 132.9, 129.7, 129.7, 132.9, 129.7, 132.9, 129.7, 132.9, 129.7, 132.9, 129.7, 132.9, 129.7, 132.9, 129.7, 132.9, 129.7, 132.9, 129.7, 132.9, 129.7, 132.9, 129.7, 132.9, 129.7, 132.9, 129.7, 132.9, 129.7, 132.9, 129.7, 129$ 134.8, 139.5, 142.4, 145.8; HRMS (FAB<sup>+</sup>) m/z calcd. for C<sub>16</sub>H<sub>16</sub>N<sup>79</sup>Br<sub>3</sub>S: 490.8553; found: 490.8534.

# Synthesis of 2,5-Dibromo-4-(2-bromo-4-butylphenylthio)-*N*-[2,5dibromo-4-(2-bromo-4-butylphenylthio)phenyl]-*N*-methylaniline (6)

A mixture of **4** (1.3 g, 2.7 mmol), **3** (1.8 g, 3.0 mmol),  $Pd(dba)_2$  (0.10 g, 0.17 mmol), DPPF (0.10 g, 0.17 mmol), *t*-BuONa (0.43 g, 4.5 mmol), and toluene (30 mL) was stirred at 100°C for 13 h, and the reaction was quenched by the addition of aq. NH<sub>4</sub>Cl. The mixture was extracted with Et<sub>2</sub>O, and the organic layer was dried over Na<sub>2</sub>SO<sub>4</sub>. The solvents were removed under reduced pressure, and the crude material was chromatographed (SiO<sub>2</sub>, CHCl<sub>3</sub>) to give bis(2,5-dibromo-4-(2-bromo-4-butylphenyl-thio)phenyl)amine (**5**) as colorless solid. This material was dissolved in THF (20 mL), and to this solution MeI (0.16

mL, 2.6 mmol), KOH (1.1 g, 20 mmol), and Bu<sub>4</sub>NI (0.18 g, 0.49 mmol) were added. The mixture was stirred at 40°C for 10.5 h, and the reaction was quenched by aq. NH<sub>4</sub>Cl. The mixture was extracted with Et<sub>2</sub>O and the organic layer dried over Na<sub>2</sub>SO<sub>4</sub>. The solvents were evaporated, and the crude material was subjected to column chromatography (SiO<sub>2</sub>, hexane/CHCl<sub>3</sub> = 4:1) to give **6** as colorless solid (1.7 g, 64%). **6**: colorless solid, mp 140–142°C. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>):  $\delta = 0.92$  (t, J = 7.3 Hz, 6H), 1.34 (sext, J = 7.4 Hz, 4H), 1.59 (quint, J = 7.6 Hz, 4H), 2.58 (t, J = 7.7 Hz, 4H), 3.20 (s, 3H), 7.08–7.13 (m, 4H), 7.20 (s, 2H), 7.22 (s, 2H), 7.48 (d, J = 1.4 Hz, 2H); <sup>13</sup>C NMR (126 MHz, CDCl<sub>3</sub>):  $\delta = 13.9$ , 22.3, 33.1, 34.9, 41.5, 119.2, 124.4, 126.3, 128.0, 128.6, 130.0, 131.3, 133.0, 133.6, 136.4, 145.1, 147.4; LRMS (FAB+) *m*/*z* 983 (M<sup>+</sup>). Anal. Calcd. for C<sub>33</sub>H<sub>31</sub>Br<sub>6</sub>NS<sub>2</sub>: C, 40.23; H, 3.17; N, 2.85. Found: C, 40.45; H, 3.34; N, 2.57.

#### Synthesis of Bis(benzothiaborino)carbazole (2)

To an Et<sub>2</sub>O (500 mL) solution of **6** (0.50 g, 0.51 mmol), *t*-BuLi (2.2 M in pentane, 3.0 mL, 6.6 mmol) was added at  $-75^{\circ}$ C, and the mixture was stirred for 20 min. To this mixture, MesB(OMe)<sub>2</sub> (0.37 mL, 1.8 mmol) was added, and the mixture was refluxed for 2 h. The solvent was evaporated, and the crude material was suspended in benzene. The suspension was filtered, and the solvent was removed under reduced pressure. CHCl<sub>3</sub> was added until all of the solid was dissolved, and the solution was concentrated to give a red precipitate. The precipitate was subjected to GPLC to give **2** as an orange solid (0.12 g, 31%). **2**: orange solid, mp 160–161°C (dec.). <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>):  $\delta = 0.88$  (t, J = 7.3 Hz, 6H), 1.30 (sext, J = 7.5 Hz, 4H), 1.54 (quint, J = 7.6 Hz, 4H), 1.96 (s, 12H), 2.46 (s, 6H), 2.60 (t, J = 7.7 Hz, 4H), 3.95 (s, 3H), 6.98 (s, 4H), 7.47 (dd, J = 8.2, 2.0 Hz, 2H), 7.63 (d, J = 2.0 Hz, 2H), 7.75 (d, J = 8.2 Hz, 2H), 7.84 (s, 2H), 8.60 (s, 2H); <sup>13</sup>C NMR (126 MHz, CDCl<sub>3</sub>):  $\delta = 13.9$ , 21.4, 22.2, 23.0, 29.5, 33.9, 35.2, 117.3, 118.1, 124.9, 126.0, 127.0, 132.9, 133.1, 133.3, 133.6, 136.5, 138.7, 138.8, 139.0, 140.7, 140.9, 141.9; HRMS (FAB<sup>+</sup>) *m/z* calcd for C<sub>51</sub>H<sub>53</sub><sup>11</sup>B<sub>2</sub>N<sup>32</sup>S<sub>2</sub>: 765.3806; found: 765.3799.

### REFERENCES

- 1. R. E. Martin and F. Diederich, Angew. Chem. Int. Ed., 38, 1350 (1999).
- S. Yamaguchi and C. Xu, J. Syn. Org. Chem. Jpn., 63, 1115 (2005); (b) A. Fukazawa, H. Yamada, and S. Yamaguchi, Angew. Chem. Int. Ed., 47, 5582 (2008); (c) M. Shimizu, K. Mochida, and T. Hiyama, Angew. Chem. Int. Ed., 47, 9760 (2008); (d) T. Matsuda, S. Kadowaki, T. Goya, and M. Murakami, Org. Lett., 9, 133 (2007).
- T. Agou, J. Kobayashi, and T. Kawashima, *Chem. Eur. J.*, **13**, 8051 (2007); (b) T. Agou, J. Kobayashi, and T. Kawashima, *Chem. Commun.*, 3204 (2007); (c) T. Agou, J. Kobayashi, and T. Kawashima, *Org. Lett.*, **8**, 2241 (2006).
- J. P. Wolfe, J. Ahmen, J. P. Sadighi, R. A. Singer, and S. L. Buchwald, *Tetrahedron Lett.*, 38, 6367 (1997).
- 5. J. F. Hartwig, Acc. Chem. Res., 31, 852 (1998).
- Similar oxidative C–C bond formation was reported in an attempted synthesis of a dibenzoselenasilin. In this case, dibenzoselenophene was obtained instead of the expected compound. O. G. Rodin, V. V. Redchenko, A. B. Kostitsyn, V. F. Traven, *Zh. Obshch. Khim.*, **58**, 1409 (1988).