

Design and synthesis of a reagent for solid-phase incorporation of the phosphothreonine mimetic (2*S*,3*R*)-2-amino-3-methyl-4-phosphonobutyric acid (Pmab) into peptides in a bio-reversible phosphonyl-*bis*-pivaloyloxymethyl (POM) prodrug form

Wen-Jian Qian · Terrence R. Burke Jr.

Received: 7 June 2013 / Accepted: 18 July 2013
© Springer-Verlag Wien (outside the USA) 2013

Abstract Reported herein are the synthesis and solid-phase peptide incorporation of *N*-Fmoc-(2*S*,3*R*)-2-amino-3-methyl-4-phosphonobutyric acid *bis*-pivaloyloxymethyl phosphoryl ester [Fmoc-Pmab(POM)₂-OH, **2**] as a phosphatase-stable phosphothreonine (pThr) mimetic bearing orthogonal protection suitable for the synthesis of Pmab-containing peptides having bio-reversible protection of the phosphonic acid moiety. This represents the first report of a bio-reversibly protected pThr mimetic in a form suitable for facile solid-phase peptide synthesis.

Keywords Phosphothreonine mimetic · Prodrug · Pmab · Solid-phase peptide synthesis · Signal transduction

Abbreviations

Ac	Acetyl
[α] _D	Specific rotation
BnBr	Benzyl bromide
Bu ^t	<i>tert</i> -Butyl
Cbz	Benzyloxycarbonyl
CH ₂ Cl ₂	Dimethyl chloride
CH ₃ CN	Acetonitrile
DIPEA	Diisopropylethylamine
DMF	Dimethylformamide

ESI	Electrospray ionization
EtOAc	Ethyl acetate
F ₂ Pmp	Difluorophosphonomethylphenylalanine
Fmoc	9-Fluorenylmethoxycarbonyl
Fmoc-OSu	9-Fluorenylmethyl succinimidyl carbonate
HBTU	1- <i>O</i> -Benzotriazole- <i>N,N,N',N'</i> -tetramethyluromium- hexafluoro-phosphate
HCl	Hydrochloric acid
His	Histidine
HOBt	1-Hydroxybenzotriazole
HPLC	High-performance liquid chromatography
HRMS	High resolution mass spectra
Leu	Leucine
LiOH	Lithium hydroxide
MeOH	Methanol
MgSO ₄	Magnesium sulfate
Mtt	Methyltrityl
<i>m/z</i>	Mass-to-charge ratio
NaHCO ₃	Sodium bicarbonate
NMP	<i>N</i> -Methyl-2-pyrrolidone
NMR	Nuclear magnetic resonance
PBD	Polo box domain
Pd-C	Palladium on carbon
PLE	Porcine liver esterase
Plk1	Polo-like kinase 1
Pmab	(2 <i>S</i> ,3 <i>R</i>)-2-Amino-3-methyl-4-phosphonobutyric acid
POM	Pivaloyloxymethyl
POMI	Pivaloyloxymethyl iodide
PPIs	Protein-protein interactions
pSer	Phosphoserine
pThr	Phosphothreonine
pTyr	Phosphotyrosine
Ser	Serine
SPPS	Solid-phase peptide synthesis

Electronic supplementary material The online version of this article (doi:10.1007/s00726-013-1567-0) contains supplementary material, which is available to authorized users.

W.-J. Qian · T. R. Burke Jr. (✉)
Chemical Biology Laboratory, Molecular Discovery Program,
Center for Cancer Research, National Cancer Institute-Frederick,
National Cancer Institute, National Institutes of Health,
Frederick, MD 21702, USA
e-mail: burkete@helix.nih.gov

$t_{1/2}$	Half life
TFA	Trifluoroacetic acid
THF	Tetrahydrofuran
TLC	Thin layer chromatography
TMS	Trimethylsilane
Trt	Trityl

Introduction

Protein phosphorylation is a fundamental form of post-translational modification that affects approximately one-third of all proteins. The presence of phosphothreonine (pThr), phosphoserine (pSer) and phosphotyrosine (pTyr) can introduce unique recognition features, which often facilitate specific protein–protein interactions (PPIs) (Yaffe 2002; Ladbury 2005; Elia and Yaffe 2005). Synthetic phosphopeptides modeled on recognition sequences can serve as useful pharmacological tools for studying these PPIs (Eisele et al. 1999; Lu et al. 2012). However, in cellular systems the bioavailability of phosphopeptides may be limited by the enzymatic lability of the phosphoryl ester bond toward phosphatases and poor membrane transport of the phosphoryl di-anionic species. While replacement of the phosphoryl ester oxygen by methylene or fluoromethylenes has addressed issues related to phosphatase hydrolysis for pTyr (Burke and Lee 2003), pSer (Shapiro et al. 1993; Perich 1994; Nair et al. 1995; Panigrahi et al. 2009) and pThr (Berkowitz et al. 1996; Otaka et al. 2000; Liu et al. 2009), cell membrane transit can still be limited by the di-anionic charge of the resulting phosphonic acids.

A general strategy for increasing the cell membrane permeability of phosphates and phosphonic acids is to mask their acidic functionality with neutral “prodrug” groups that can be removed enzymatically once the agent is within the cell. Although a variety of general prodrug strategies have been for phosphonic acids (Schultz 2003; Hecker and Erion 2008), there are few reports detailing incorporation of these functionalities into protected amino acid derivatives that are amenable to solid-phase peptide synthesis. Recent examples of the latter include the application of 5'-nitro-furyl-2'-methyl *N*-(4''-chlorobutyl)phosphonamido ester prodrug strategy to the pTyr mimetic difluorophosphonomethylphenylalanine (F₂Pmp) (Boutselis et al. 2007) and to the pSer mimetic, difluorophosphonoaminobutyric acid (Arrendale et al. 2012). In these two cases, the prodrug-protected reagents were used to prepare dipeptides for examination in whole cell studies. The use of these reagents for the solid-phase synthesis of longer peptides is potentially limited by chemical instability of the phosphonamido group to repetitive cycles of piperidine treatment needed for Fmoc-based solid-phase protocols (Boutselis et al. 2007). In

the broader sense, there is a paucity of reagents available for the solid-phase synthesis of polypeptides containing prodrug-protected pThr mimetics.

The pivaloyloxymethyl (POM) group is an esterase-labile moiety that has been widely applied to phosphoryl prodrug protection of nucleotides (Hecker and Erion 2008) and phosphate and phosphonic acid functionality in small molecules and peptide mimetics (Stankovic et al. 1997; Mandal et al. 2009, 2011; Zhao and Etzkorn 2007). However, in spite of its usefulness in these latter contexts, there are no prior reports of polypeptides containing pThr, pSer or their phosphonic acid-based mimetics, bearing POM protection. Given the importance of pThr in a large number of biological processes (Elia and Yaffe 2005), a reagent that would permit the solid-phase synthesis of polypeptides containing POM-protected pThr mimetics would be highly desirable.

We had previously reported the preparation of *N*-Fmoc-(2*S*,3*R*)-2-amino-3-methyl-4-phosphonobutyric acid bis-*tert*-butyl phosphoryl ester [Fmoc-Pmab(Bu^t)₂-OH, **1**, Fig. 1] as a hydrolytically-stable pThr mimetic bearing orthogonal protection suitable for the standard Fmoc-based synthesis of peptides containing Pmab. However this reagent yields peptides in which the Pmab residue has its phosphonic acid moiety in the free di-anionic form (Liu et al. 2009). To date, there have been no reports of Pmab suitably derivatized for the solid-phase synthesis of polypeptides that maintained the phosphonic acid in a prodrug-protected form. In the current paper we describe the first preparation of Fmoc-Pmab having the phosphonic acid group masked as its POM *bis*-esters [Fmoc-Pmab(POM)₂-OH, **2**, Fig. 1] and demonstrate its use in the solid-phase synthesis of a peptide bearing full POM protection of the Pmab residue.

Materials and methods

General methods

All experiments involving moisture-sensitive compounds were conducted under anhydrous conditions. Fmoc-Ser(Trt)-OH, and Fmoc-His(Mtt)-OH were purchased from




Fig. 1 Structures of key reagents

NovaChioChem. All solvents were purchased in anhydrous form (Aldrich) and used directly. Analytical TLCs were performed using Analtech precoated plates (Uniplate, silica gel GHLF, 250 nm) containing a fluorescence indicator. NMR spectra were recorded using a Varian Inova 400 MHz spectrometer. Coupling constants are reported in hertz, and peak shifts are reported in δ (ppm) relative to TMS. Low-resolution mass spectra (ESI) were measured with an Agilent 260 1200 LC/MSD-SL system. High resolution mass spectra (HRMS) were obtained by positive ion, ESI analysis on a Thermo Scientific LTQ-XL Orbitrap mass spectrometer with HPLC sample introduction using a short narrow-bore C_{18} reversed-phase column with CH_3CN-H_2O gradients. Reported m/z values are the average of eight or more scans over the chromatographic peak of interest.


Synthesis of Fmoc-Pmab(POM)₂-OH (**2**)

(2*S*,3*R*)-Benzyl 2-(((benzyloxy)carbonyl)amino)-4-(di-tert-butoxyphosphoryl)-3-methylbutanoate (**4**) To a solution of **3** (Liu et al. 2009) (0.26 g, 0.57 mmol) in THF–H₂O (4:1; 5 mL) at 0 °C was added a solution of LiOH·H₂O (48 mg, 1.14 mmol) and the mixture was stirred at room temperature until all starting material was consumed as indicated by TLC. The pH was adjusted to 2–3 by the addition of 1 N aqueous HCl and THF was removed by evaporation. The residue was extracted (EtOAc) and the combined organic phase was washed with brine, dried (MgSO₄), filtered and concentrated. To a solution of the crude residue in DMF (5.0 mL) were added sequentially, NaHCO₃ (95 mg, 1.14 mmol) and BnBr (0.10 mL, 0.85 mmol) at room temperature under argon and the mixture was stirred until the starting material was consumed as indicated by TLC.

The mixture was diluted with EtOAc and washed with H₂O and brine, dried (MgSO₄) and filtered and concentrated. Purification by silica gel column chromatography (CH₂Cl₂ : MeOH from 100:1 to 30:1) afforded **4** as a colorless oil (0.25 g, 80 % for two steps, Scheme 1). $[\alpha]_D^{21.6}$ –0.59 (c 1.0, CHCl₃); ¹H NMR (400 MHz, CDCl₃) δ 7.40–7.28 (m, 10H), 5.91 (d, J = 8.0 Hz, 1H), 5.20–5.06 (m, 4H), 4.38–4.31 (m, 1H), 2.42 (brs, 1H), 1.84–1.69 (m, 1H), 1.57–1.48 (m, 1H), 1.47–1.45 (m, 18H), 1.10 (d, J = 8.0 Hz, 3H) ppm; ¹³C NMR (100 MHz, CDCl₃) δ 171.7, 156.4, 136.5, 135.4, 128.8, 128.7, 128.6, 128.3, 82.4 (d, ² J_{CP} = 10 Hz), 82.3 (d, ² J_{CP} = 10 Hz), 67.4, 67.2, 59.8 (d, ³ J_{CP} = 10 Hz), 34.0, 32.6, 30.59, 30.56, 17.6 ppm; ESI-HRMS m/z calcd for C₂₈H₄₁NO₇P (M+H)⁺: 534.2621, found: 534.2594.

(((2*R*,3*S*)-4-(Benzyloxy)-3-(((benzyloxy)carbonyl)amino)-2-methyl-4-oxobutyl)phosphoryl)bis(oxy))bis(methylene) Bis(2,2-dimethylpropanoate) (**5**) A solution of **4** (0.27 g, 0.5 mmol) in 10 % TFA (CH₂Cl₂) was stirred at room temperature (1 h), then volatiles were removed under vacuum and the residue was taken up in DMF with pivaloyloxymethyl iodide (POMI) (Bandgar et al. 2011) (0.32 mL, 2.0 mmol) and DIPEA (0.35 mL, 2.0 mmol) and stirred at room temperature under argon (overnight). The mixture was diluted with H₂O, extracted with EtOAc and the combined organic extracts were washed with H₂O, and brine, dried (MgSO₄), filtered and concentrated. Purification by silica gel chromatography (EtOAc : hexanes from 1:4 to 1:1) afforded **5** as a white semi-solid (0.24 g, 74 % yield for two steps, Scheme 1). $[\alpha]_D^{21.9}$ 2.88 (c 0.7, CHCl₃); ¹H NMR (400 MHz, CDCl₃) δ 7.42–7.18 (m, 10H), 5.70–5.60 (m, 4H), 5.56 (d, J = 12.0 Hz, 1H), 5.19 (s, 2H), 5.11 (s, 2H), 4.42–4.34 (m, 1H), 2.45 (brs, 1H), 2.04–1.91 (m, 1H), 1.78–1.67 (m, 1H), 1.22 (s, 9H),


Scheme 1 Synthesis of title reagent **2**



1.21 (s, 9H), 1.08 (d, $J = 4.0$ Hz, 3H) ppm; ^{13}C NMR (100 MHz, CDCl_3) δ 177.1, 171.1, 156.3, 136.3, 135.2, 128.9, 128.8, 128.7, 128.44, 128.35, 81.7 (d, $^2J_{\text{CP}} = 20$ Hz), 67.6, 67.4, 59.2 (d, $^3J_{\text{CP}} = 10$ Hz), 38.9, 31.9, 30.3, 29.9, 29.6 (d, $^1J_{\text{CP}} = 140$ Hz), 27.0, 17.5 ppm; ESI-HRMS m/z calcd for 650.2725, found: 650.2704.

(2*S*,3*R*)-2-(((9*H*-Fluoren-9-yl)methoxy)carbonylamino)-4-(bis((pivaloyloxy)methoxy)phosphoryl)-3-methylbutanoic Acid [*N*-Fmoc-*Pmab*(POM)₂-OH] (**2**) A solution of **5** (0.18 g, 0.28 mmol) in MeOH was hydrogenated over 10 % Pd-C (30 mg) until the reaction was complete as indicated by TLC. The mixture was filtered, evaporated to dryness and reacted with Fmoc-OSu (0.19 g, 0.55 mmol) and NaHCO_3 (70 mg, 0.83 mmol) in dioxane : H_2O (1:1, 5.5 mL) at room temperature (overnight). The mixture was acidified with 1 N HCl, extracted with EtOAc and the combined organic phases were washed with brine, dried (MgSO_4), filtered and the filtrate concentrated. Purification by silica gel column chromatography (CH_2Cl_2 :MeOH from 20:1 to 4:1) provided **2** as a colorless oil (0.14 g, 75 % yield for two steps, Scheme 1). $[\alpha]_{\text{D}}^{21.0}$ 12.8 (c 0.66, CHCl_3); ^1H NMR (400 MHz, CDCl_3) δ 7.77 (d, $J = 8.0$ Hz, 2H), 7.60 (d, $J = 8.0$ Hz, 2H), 7.40 (t, $J = 8.0$ Hz, 2H), 7.32 (td, $J_1 = 8.0$ Hz, $J_2 = 4.0$ Hz, 2H), 5.78–5.64 (m, 5H), 4.51–4.44 (m, 1H), 4.40 (d, $J = 8.0$ Hz, 2H), 4.22 (t, $J = 8.0$ Hz, 1H), 2.54 (brs 1H), 2.18–2.03 (m, 1H), 1.97–1.83 (m, 1H), 1.24 (s, 9H), 1.23 (s, 9H), 1.13 (d, $J = 8.0$ Hz, 3H) ppm; ^{13}C NMR (100 MHz, CDCl_3) δ 177.2, 172.8, 156.4, 144.0, 143.8, 141.5, 128.0, 127.3, 125.3, 120.2, 82.0, 67.5, 58.3 (d, $^3J_{\text{CP}} = 10$ Hz), 47.3, 39.0, 31.4, 29.4 (d, $^1J_{\text{CP}} = 140$ Hz), 27.0, 17.6 (d, $^3J_{\text{CP}} = 7$ Hz) ppm; ESI-HRMS m/z calcd for $\text{C}_{32}\text{H}_{42}\text{NO}_{11}\text{PNa}$ ($\text{M}+\text{Na}$)⁺: 670.2393, found: 670.2376.

Scheme 2 Synthesis of bis-POM-protected peptide **6** using reagent **2**



Use of reagent **2** for the solid-phase synthesis of peptide **6**

Standard Fmoc-protected amino acids were purchased from Novabiochem. Peptides were synthesized on NovaSyn[®] TG Sieber resin (Novabiochem, cat. no. 01-64-0092) using Fmoc-based solid-phase protocols in *N*-methyl-2-pyrrolidone (NMP). 1-*O*-Benzotriazole-*N,N,N',N'*-tetramethyl-uronium-hexafluoro-phosphate (HBTU) (5.0 eq.), hydroxybenzotriazole (HOBT) (5.0 eq.) and *N,N*-diisopropylethylamine (DIPEA) (10.0 eq.) were used as coupling reagents. Reagent **2** was employed for introduction of the first residue. Following completion of peptide elongation and Fmoc-deprotection, amino terminal acetylation was achieved using 1-acetylimidazole. The finished resin (**7**, Scheme 2) was washed with DMF, MeOH, CH_2Cl_2 and diethyl ether and then dried under vacuum (overnight). Peptide **6** was cleaved from the resin using 1 % TFA in CH_2Cl_2 . The resin was removed by filtration and the filtrate was concentrated under vacuum, then precipitated with cold ether and the precipitate washed with cold ether. The resulting solid was dissolved in 50 % aqueous acetonitrile (5 mL) and purified by reverse phase preparative HPLC using a Phenomenex C₁₈ column (21 mm dia × 250 mm, cat. no: 00G-4436-P0) with a linear gradient from 0 % aqueous acetonitrile (0.1 % TFA) to 100 % acetonitrile (0.1 % trifluoroacetic acid) over 30 min at a flow rate of 10.0 mL/min. Lyophilization gave peptide **6** as a white powder (>99 % pure by HPLC, Scheme 2). ESI-MS m/z calcd for $\text{C}_{39}\text{H}_{66}\text{N}_8\text{O}_{14}\text{P}$ ($\text{M}+\text{H}$)⁺: 901.4, found: 901.4.

Pig liver esterase hydrolysis studies on peptide **6**

Following literature procedures (Srivastva and Farquhar 1984), 0.05 M potassium phosphate buffer (pH 7.4) was




Fig. 2 Results of pig liver esterase treatment of peptide **6**

placed in a centrifuge tube and a solution of peptide **6** in MeOH was added to achieve a concentration of 200 μM of **6** with MeOH being less than 1 %. To a 1 mL aliquot of the above solution was added pig liver esterase (57.6 units) and the reaction mixture was incubated at 37 °C with gentle agitation. At various time points, aliquots of the reaction mixture (50 μL) were transferred to an Eppendorf tube containing MeCN (50 μL). Following filtration, the hydrolysis product was monitored by LC-MS. Data are shown in Fig. 2.

Results and discussion

Preparation of reagent **2** began with Cbz-Pmab(Bu_2^t)-OMe (**3**), which is an intermediate in our previously reported synthesis of **1** (Liu et al. 2009). Benzyl *trans*-esterification of **3–4** was accomplished by initial LiOH-mediated hydrolysis of the methyl ester followed by reaction of the free acid with benzyl bromide (NaHCO_3 in DMF, 80 % yield for two steps, Scheme 1). Subsequent treatment of **4** with pivaloyloxymethyl iodide (POMI) (Bandgar et al. 2011) and diisopropylethyl amine (DIPEA) in DMF provided the corresponding bis-POM-protected intermediate (**5**). Hydrogenolytic removal of amino and carboxyl protecting groups and introduction of *N*-Fmoc-protection by treatment with 9-fluorenylmethyl succinimidyl carbonate (Fmoc-OSu) in aqueous THF with NaHCO_3 , gave the desired reagent **2** (55 % yield from **4**) (Scheme 1).

In order to demonstrate the usefulness of **2** for the incorporation of Pmab(POM)₂ into peptides by solid-phase techniques, we chose the sequence, Ac-Pro-Leu-His-Ser-Pmab-amide, which has previously shown to bind with good affinity to the polo-like kinase 1 (Plk1) polo box domain (PBD) (Yun et al. 2009; Liu et al. 2011, 2012a, b). PBD-binding ligands may potentially serve as anti-cancer agents by blocking the spatial organization required for Plk1 to function in oncogenic processes (van de Weerd

et al. 2008). The syntheses of Ac-Pro-Leu-His-Ser-Pmab(POM)₂-amide (**6**) was accomplished on NovaSyn TG Sieber resin using standard Fmoc protocols. Histidine and serine were employed in their 4-methoxytrityl (Mtt) and trityl (Trt) side chain-protected forms, respectively, to allow their cleavage under mildly acidic conditions (Scheme 2). Following synthesis completion, the resin-bound Ac-Pro-Leu-His(Mtt)-Ser(Trt)-Pmab(POM)₂-amide (**7**) was subjected to treatment with 1 % TFA, which resulted in removal of histidine and serine protecting groups and cleavage of the peptide from the resin with retention of Pmab-*bis*-POM functionality. Purification by HPLC provided Ac-Pro-Leu-His-Ser-Pmab(POM)₂-amide (**6**) as a white solid (Scheme 2).

In order to examine the bio-reversibility of the Pmab POM protection of **6**, we performed in vitro assays using porcine liver esterase (PLE) in phosphate buffer at pH 7.4 (Srivastva and Farquhar 1984). We observed that the parent Pmab *bis*-POM-containing **6** was rapidly converted to the mono-POM-containing product ($t_{1/2} \approx 5$ min) with further deprotection to the free Pmab-containing peptide occurring at a slower rate (20 % conversion over 8 h) (Fig. 2). Reduction in the rate of enzymatic cleavage of a second POM group is known (Srivastva and Farquhar 1984).

Conclusions

Our current paper presents the synthesis of a reagent [Fmoc-Pmab(POM)₂-OH (**2**)] that allows the facile synthesis of peptides containing the phosphatase-stable pThr mimetic, Pmab, bearing bio-reversible POM protection. This represents a rare example of a reagent that allows the solid-phase synthesis of polypeptides having a pThr mimetic in bio-reversible prodrug form. Reagent **2** should find utility in a variety of pharmacological applications.

Acknowledgments This research was supported by the Intramural Research Program of the NIH, Center for Cancer Research, NCI-Frederick and the National Cancer Institute, National Institutes of Health.

Conflict of interest The authors declare that they have no conflict of interest.

References

- Arrendale A, Kim K, Choi J, Li W, Geahlen RL, Borch RF (2012) Synthesis of a phosphoserine mimetic prodrug with potent 14-3-3 protein inhibitory activity. *Chem Biol* 19:764–771
- Bandgar BP, Sarangdhar RJ, Viswakarma S, Ahamed FA (2011) Synthesis and biological evaluation of orally active prodrugs of indomethacin. *J Med Chem* 54:1191–1201
- Berkowitz DB, Eggen M, Shen Q, Shoemaker RK (1996) Ready access to fluorinated phosphonate mimics of secondary phosphates. Synthesis of the (α,α -difluoroalkyl)phosphonate analogs

- of L-phosphoserine, L-phosphoallothreonine, and L-phosphothreonine. *J Org Chem* 61:4666–4675
- Boutselis IG, Yu X, Zhang Z-Y, Borch RF (2007) Synthesis and cell-based activity of a potent and selective protein tyrosine phosphatase 1B inhibitor prodrug. *J Med Chem* 50:856–864
- Burke TR Jr, Lee K (2003) Phosphotyrosyl mimetics in the development of signal transduction inhibitors. *Acc Chem Res* 36:426–433
- Eisele F, Owen DJ, Waldmann H (1999) Peptide conjugates as tools for the study of biological signal transduction. *Bioorg Med Chem* 7:193–224
- Eliu AEH, Yaffe MB (2005) Phosphoserine/threonine binding domains. In: Cesare G (ed) *Modular protein domains*. Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim, pp 163–179. doi:10.1002/3527603611.ch8
- Hecker SJ, Erion MD (2008) Prodrugs of phosphates and phosphonates. *J Med Chem* 51:2328–2345
- Ladbury JE (2005) Protein–protein recognition in phosphotyrosine-mediated intracellular signaling. *Protein Rev* 3:165–184
- Liu F, Park J-E, Lee KS, Burke TR Jr (2009) Preparation of orthogonally protected (2S,3R)-2-amino-3-methyl-4-phosphonobutyric acid (Pmab) as a phosphatase-stable phosphothreonine mimetic and its use in the synthesis of polo-box domain-binding peptides. *Tetrahedron* 65:9673–9679
- Liu F, Park J-E, Qian W-J, Lim D, Graber M, Berg T, Yaffe MB, Lee KS, Burke TR Jr (2011) Serendipitous alkylation of a Plk1 ligand uncovers a new binding channel. *Nat Chem Biol* 7:595–601
- Liu F, Park J-E, Qian W-J, Lim D, Scharow A, Berg T, Yaffe MB, Lee KS, Burke TR (2012a) Identification of high affinity polo-like kinase 1 (Plk1) polo-box domain binding peptides using oxime-based diversification. *ACS Chem Biol* 7:805–810
- Liu F, Park J-E, Qian W-J, Lim D, Scharow A, Berg T, Yaffe MB, Lee KS, Burke TR (2012b) Peptoid-peptide hybrid ligands targeting the polo box domain of polo-like kinase 1. *Chem Biochem* 13:1291–1296
- Lu CHS, Liu K, Tan LP, Yao SQ (2012) Current chemical biology tools for studying protein phosphorylation and dephosphorylation. *Chem Eur J* 18:28–39
- Mandal PK, Liao WSL, McMurray JS (2009) Synthesis of phosphatase-stable, cell-permeable peptidomimetic prodrugs that target the SH2 domain of Stat3. *Org Lett* 11:3394–3397
- Mandal PK, Gao F, Lu Z, Ren Z, Ramesh R, Birtwistle JS, Kaluarachchi KK, Chen X, Bast RC, Liao WS, McMurray JS (2011) Potent and selective phosphopeptide mimetic prodrugs targeted to the Src homology 2 (SH2) domain of signal transducer and activator of transcription 3. *J Med Chem* 54:3549–3563
- Nair SA, Lee B, Hangauer DG (1995) Synthesis of orthogonally protected L-homocysteine and L-2-amino-4-phosphonobutanoic acid from L-homoserine. *Synthesis* (7):810–814
- Otaka A, Mitsuyama E, Kinoshita T, Tamamura H, Fujii N (2000) Stereoselective synthesis of CF₂-substituted phosphothreonine mimetics and their incorporation into peptides using newly developed deprotection procedures. *J Org Chem* 65:4888–4899
- Panigrahi K, Eggen M, Maeng J-H, Shen Q, Berkowitz DB (2009) The alpha, alpha-difluorinated phosphonate L-pSer-analogue: an accessible chemical tool for studying kinase-dependent signal transduction. *Chem Biol* 16:928–936
- Perich JW (1994) Efficient Fmoc/solid-phase synthesis of Abu(P)-containing peptides using Fmoc-Abu(PO₃Me₂)-OH. *Int J Pept Protein Res* 44:288–294
- Schultz C (2003) Prodrugs of biologically active phosphate esters. *Bioorg Med Chem* 11:885–898
- Shapiro G, Buechler D, Ojea V, Pombo-Villar E, Ruiz M, Weber HP (1993) Synthesis of both D- and L-Fmoc-Abu[PO(OCH₂CH:CH₂)₂]-OH for solid phase phosphonopeptide synthesis. *Tetrahedron Lett* 34:6255–6258
- Srivastva DN, Farquhar D (1984) Bioreversible phosphate protective groups: synthesis and stability of model acyloxymethyl phosphates. *Bioorg Chem* 12:118–129
- Stankovic CJ, Surendran N, Lunney EA, Plummer MS, Para KS, Shahripour A, Fergus JH, Marks JS, Herrera R, Hubbell SE, Humblet C, Saltiel AR, Stewart BH, Sawyer TK (1997) The role of 4-phosphonodifluoromethyl- and 4-phosphono-phenylalanine in the selectivity and cellular uptake of SH2 domain ligands. *Bioorg Med Chem Lett* 7:1909–1914
- van de Weerd BCM, Littler DR, Klompmaker R, Huseinovic A, Fish A, Perrakis A, Medema RH (2008) Polo-box domains confer target specificity to the Polo-like kinase family. *Biochim Biophys Acta Mol Cell Res* 1783:1015–1022
- Yaffe MB (2002) Phosphotyrosine-binding domains in signal transduction. *Nat Rev Mol Cell Biol* 3:177–186
- Yun S-M, Moulaei T, Lim D, Bang JK, Park J-E, Shenoy SR, Liu F, Kang YH, Liao C, Soung N-K, Lee S, Yoon D-Y, Lim Y, Lee D-H, Otaka A, Appella E, McMahon JB, Nicklaus MC, Burke TR Jr, Yaffe MB, Wlodawer A, Lee KS (2009) Structural and functional analyses of minimal phosphopeptides targeting the polo-box domain of polo-like kinase 1. *Nat Struct Mol Biol* 16:876–882
- Zhao S, Etkorn FA (2007) A phosphorylated prodrug for the inhibition of Pin1. *Bioorg Med Chem Lett* 17:6615–6618