


This article was downloaded by: [Dalhousie University]

On: 13 November 2013, At: 07:24

Publisher: Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Nucleosides, Nucleotides and Nucleic Acids

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/Incn20>

Synthesis and Properties of Pyrrolidine-Based Negatively Charged Dna Mimics

Vladimir A. Efimov^a, Valeryi N. Klykov^a & Oksana G.

Chakhmakhcheva^a

^a Shemyakin-Ovchinnikov Institute of Bioorganic Chemistry, Moscow, Russia

Published online: 05 Dec 2007.

To cite this article: Vladimir A. Efimov, Valeryi N. Klykov & Oksana G. Chakhmakhcheva (2007) Synthesis and Properties of Pyrrolidine-Based Negatively Charged Dna Mimics, *Nucleosides, Nucleotides and Nucleic Acids*, 26:10-12, 1595-1599, DOI: [10.1080/15257770701548345](https://doi.org/10.1080/15257770701548345)

To link to this article: <http://dx.doi.org/10.1080/15257770701548345>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

SYNTHESIS AND PROPERTIES OF PYRROLIDINE-BASED NEGATIVELY CHARGED DNA MIMICS

Vladimir A. Efimov, Valeryi N. Klykov, and Oksana G. Chakhmakhcheva

□ *Shemyakin-Ovchinnikov Institute of Bioorganic Chemistry, Moscow, Russia*

□ *A set of novel chiral pyrrolidine-based nucleotide mimics, in which nucleobase, hydroxyl group and phosphonic acid residue were attached to different carbon atoms of the pyrrolidine ring, was synthesized. These monomers were used for the synthesis of the corresponding oligomers, and their physico-chemical properties were evaluated.*

Keywords DNA mimics; pyrrolidine; hybridization; antisense reagents

INTRODUCTION

In the last years, a search for modified oligonucleotides and their analogues with improved physico-chemical and biological properties attracted our attention in connection with their application in molecular biology and medicine as diagnostic probes and antisense agents. With this aim in view, we have undertaken the investigations on the synthesis of several types of negatively charged oligonucleotide mimics constructed from pyrrolidine-based chiral monomer units connected with phosphonate ester bonds. Particularly, we reported pHypNA mimics containing 1-acetyl-4-hydroxypyrrolidine-2-phosphonic acid backbone^[1] (Figure 1). *Trans-L* type of pHypNAs, similarly to HypNA-pPNA hetero-oligomers developed by us earlier,^[2] have demonstrated high hybridization and discrimination characteristics as well as the ability to hybridize to complementary nucleic acids in low salt concentrations.^[1–3] Analysis of biological properties of pHypNA oligomers has shown that this type of DNA mimics combined high biological stability with the ability to penetrate cell membranes. They do not activate

The authors thank Russian Foundation of Basic Research for financial support.

Address correspondence to V. A. Efimov, ul. Miklukho-Maklaya 16/10, Moscow 117997, Russia.
E-mail: eva@mail.ibch.ru




FIGURE 1 General chemical structures of HypNA-pPNA and pHypNA mimic oligomers and four chiral pHypNA monomers.

RNase H and inhibit mRNA translation by hybridization arrest. Other properties of pHypNA mimics include ability to bind RNA targets stronger than DNA, good water solubility and cellular uptake with distribution in cytosol and preferentially in nucleus as well as a long shelf-life at room temperature. Thus, these compounds can be evaluated as useful tools for application in molecular biology, particularly in nucleic acid based diagnostics, isolation and analysis of nucleic acids and in the study of gene expression.^[3–5] Gene knockdown experiments in vivo revealed that their effect lasts over a period of several days. So, the use of these mimics can represent a potent technology for administering antisense-based drugs for future therapeutic applications.^[3,4] To continue these investigations, we obtained several novel types of monomers on the base of pyrrolidine (PyrRNAs).

RESULTS AND DISCUSSION

In pyrrolidine-based monomers obtained, nucleobase, hydroxyl group and phosphonic acid residue were attached to different carbon atoms of the pyrrolidine ring (Figure 2). As examples, the synthetic routes to obtain *trans*-L PyrRNA-2 and PyrRNA-3 monomers are depicted in Scheme 1. The derivatives of *trans*-4-OH-*N*-acetyl-L-prolinol bearing heterocyclic bases,




FIGURE 2 General chemical structures of chiral monomer units synthesized for the construction of the corresponding PyrrNA mimic oligomers.

which were used as starting compounds, were obtained as described.^[6] Schemes for the synthesis of other PyrrNA monomers will be published elsewhere. The synthesis of the corresponding PyrrNA oligomers was carried out by the modified solid-phase phosphotriester method with O-nucleophilic intramolecular catalysis similar to the synthesis of pHypNA and HypNA-pPNA oligomers.^[2,4] Preliminary results obtained in the study of the hybridization properties of PyrrNA-2 and PyrrNA-3 mimics revealed that oligomers constructed from monomers with *cis-L* configuration are not able to form stable complexes with the complementary oligonucleotides, whereas oligomers constructed from *trans-L* monomers are able to form complexes with the complementary targets, but the stability of these complexes is lower than

TABLE 1 Stability of complexes formed by homo-Thy₁₅ PyrRNA DNA mimics with the complementary DNA, or RNA, targets in 5–10 μ M concentration in 10 mM Tris-HCl (pH 7.5)/0.5 M NaCl/10 mM MgCl₂

Oligomer type	T _m , °C	
	DNA target	RNA target
DNA	47	35
pHypNA	<i>trans</i> -L 76	64
	<i>trans</i> -D 53	47
PyrRNA-1	<i>trans</i> -L ND ^a	ND ^a
	<i>cis</i> -L <20	<20
	<i>trans</i> -D ND ^a	ND ^a
	<i>cis</i> -D <20	<20
PyrRNA-2	<i>trans</i> -L 35	30
	<i>cis</i> -L <20	<20
PyrRNA-3	<i>trans</i> -L 31	27
	<i>cis</i> -L <20	<20
PyrRNA-4	<i>trans</i> -L ND ^a	ND ^a
	<i>cis</i> -L <20	<20
	<i>trans</i> -D <20	<20
	<i>cis</i> -D ND ^a	ND ^a
PyrRNA-5	<i>trans</i> -L ND ^a	ND ^a
	<i>cis</i> -D ND ^a	ND ^a

^aExamination is currently in progress.

the stability of natural DNA/DNA and DNA/RNA duplexes and significantly lower than the stability of complexes between *trans*-L pHypNA oligomers and DNA (or RNA) targets (Table 1). Also, *cis*-L and *cis*-D PyrRNA-1 oligomers as well as *cis*-L and *trans*-D PyrRNA-4 oligomers were not able to form complexes with the complementary targets. The investigations on the properties of other types of PyrRNA mimics are currently in progress.

REFERENCES

1. Efimov, V.; Klykov, V.; Chakhmakhcheva, O. Phosphono peptide nucleic acids with a constrained hydroxyproline-based backbone, *Nucleosides, Nucleotides Nucleic Acids* **2003**, 22, 593–599.
2. Efimov, V.; Buryakova, A.; Chakhmakhcheva, O. Synthesis of polyacrylamides N-substituted with PNA-like oligonucleotide mimics for molecular diagnostic applications. *Nucleic Acids Res.* **1999**, 27, 4416–4426.
3. Efimov, V.; Chakhmakhcheva, O.; Wickstrom, E. Synthesis and application of negatively charged PNA analogues. *Nucleosides, Nucleotides, Nucleic Acids* **2005**, 24, 1853–1874.
4. Efimov, V.; Birikh, K.; Staroverov, D.; Lukyanov, S.; Tereshina, M.; Zarskiy, A.; Chakhmakhcheva, O. Hydroxyproline-based DNA mimics provide an efficient gene silencing in vitro and in vivo. *Nucleic Acids Res.* **2006**, 34, 2247–2257.
5. Efimov, V.; Chakhmakhcheva, O. Hydroxyproline-based DNA mimics: a review on synthesis and properties. *Collect. Czech. Chem. Commun.* **2006**, 71, 929–955.
6. Ceulemans, G.; van Aerschot, A.; Wroblowski, B.; Rozenski, J.; Hendrix, C.; Herdewijn, P. Oligonucleotide analogues with 4-hydroxy-N-acetylprolinol as sugar substitute. *Chem. Eur. J.* **1997**, 3, 1997–2010.