





Figure 1. Molecular structure of **4b**; selected bond lengths (Å) and angles (deg).

Hiyama reported that hydrosilation of 1,1,4,4-tetrakis(trimethylsilyl)butatriene catalyzed by rhodium complexes gave 1,2-addition products in which a silyl group attached at the terminal position.⁸ When **1** was treated with diphenylsilane in the presence of a catalytic amount of $\text{RhCl}(\text{PPh}_3)_3$, hydrosilated products were obtained in moderate yields (Scheme 3).²² Contrary to the previous results, allenes **5** were obtained as major products. These have a silyl group at the internal position as a result of 2,1-addition, while 2,3-addition gave 1,3-dienes **6** as minor products. Interestingly, **6** is formally an *anti*-addition product. A reaction of **1a** using Ph_2SiD_2 (97%D) gave deuterated products **7** and **8**. Deuterium was incorporated at C1 in **7** and C3 in **8** (96 and 97%D). It should be noted that D was not attached at the other carbons. The possibility that (*Z*)-**1** isomerized to 1,3-enyne **9** and this accepted hydrosilation to give **6** can be ruled out. However, it is not clear yet whether **6** formed via direct *anti*-addition of Si-H on (*Z*)-**1** or *syn*-addition to (*E*)-**1** that was isomerized from (*Z*)-**1**. Further investigation is now in progress.



Scheme 3. Hydrosilation of butatrienes catalyzed by Rh complexes.

The authors are grateful to Ms. Keiko Yamada for elemental analysis. Dr. Hiroyuki Koshino is appreciated for characterization of hydrosilation products by NMR. A part of this work was financially supported by Mitsubishi Chemical Corporation Fund.

References and Notes

- # On leave from Dae-gu Christian University, Korea (JISTEC Winter Institute Program, 2002)
- ## Current address: RIKEN Venture OM Chemtech Ltd., Wako, Saitama 351-0198

- 1 A. Nakamura, P.-J. Kim, and N. Hagihara, *J. Organomet. Chem.*, **3**, 7 (1965).
- 2 a) K. K. Joshi, *J. Chem. Soc. A*, **1966**, 594. b) D. Bright and O. S. Mills, *Chem. Commun.*, **1966**, 211.
- 3 L. Hagelee, R. West, J. Calabrese, and J. Norman, *J. Am. Chem. Soc.*, **101**, 4888 (1979).
- 4 U. Schubert and J. Grönen, *Organometallics*, **6**, 2459 (1987).
- 5 A. Maercker and A. Groos, *Angew. Chem., Int. Ed. Engl.*, **35**, 210 (1996).
- 6 L. Stehling and G. Wilke, *Angew. Chem., Int. Ed. Engl.*, **27**, 571 (1988).
- 7 M. Iyoda, S. Tanaka, H. Otani, M. Nose, and M. Oda, *J. Am. Chem. Soc.*, **110**, 8494 (1988).
- 8 T. Kusumoto and T. Hiyama, *Bull. Chem. Soc. Jpn.*, **63**, 3103 (1990).
- 9 L. Stehling and G. Wilke, *Angew. Chem., Int. Ed. Engl.*, **24**, 496 (1985).
- 10 a) A. Nakamura, *Bull. Chem. Soc. Jpn.*, **38**, 1868 (1965). b) R. O. Angus, Jr., M. N. Janakiraman, R. A. Jacobson, and R. P. Johnson, *Organometallics*, **6**, 1909 (1987).
- 11 Y. Wakatsuki, H. Yamazaki, N. Kumegawa, T. Satoh, and J. Y. Satoh, *J. Am. Chem. Soc.*, **113**, 9604 (1991).
- 12 Y. Wakatsuki, H. Yamazaki, N. Kumegawa, and P. S. Johar, *Bull. Chem. Soc. Jpn.*, **66**, 987 (1993).
- 13 a) Y. Suzuki, R. Hirotani, H. Komatsu, and H. Yamazaki, *Chem. Lett.*, **1999**, 1299. b) M. Ogasawara, H. Ikeda, K. Ohtsuki, and T. Hayashi, *Chem. Lett.*, **2000**, 776. c) T. Ohmura, S. Yorozyu, Y. Yamamoto, and N. Miyaura, *Organometallics*, **19**, 365 (2000).
- 14 N. Suzuki, M. Nishiura, and Y. Wakatsuki, *Science*, **295**, 660 (2002).
- 15 Cumulenylidene complexes are also known, see: M. I. Bruce, *Chem. Rev.*, **98**, 2797 (1998) and the references cited therein.
- 16 **3a**: ^1H NMR (C_6D_6 , Me_4Si) δ = -0.14 (s, 18H), 6.71 (d, $^3J_{\text{Rh-H}}$ = 2.6 Hz, 2H), 7.01–7.13 (m, 18H), 7.90–7.97 (m, 12H). ^{13}C NMR (C_6D_6 , Me_4Si) δ = -0.67, 124.34, 127.77 ($^2J_{\text{P-C}}$ = 5.0 Hz), 129.45, 132.95 ($^1J_{\text{P-C}}$ = 21.2 Hz), 135.15 ($^2J_{\text{P-C}}$ = 6.1 Hz), 156.97 (q, $^1J_{\text{Rh-C}}$ = 12.9, $^2J_{\text{P-C}}$ = 3.4 Hz). ^{31}P NMR (C_6D_6 , H_3PO_4) δ = 31.2 (d, $^1J_{\text{P-Rh}}$ = 138.1 Hz). ^{29}Si NMR (C_6D_6 , Me_4Si) δ = -11.5 (d, $^3J_{\text{Si-Rh}}$ = 3.2 Hz). Anal. Calcd for $\text{C}_{46}\text{H}_{50}\text{ClP}_2\text{RhSi}_2$: C 64.29, H 5.86, Found: C 64.72, H 5.84. Yield 83% by ^1H NMR (35% isol.).
- 17 P. J. Stang, M. R. White, and G. Maas, *Organometallics*, **2**, 720 (1983).
- 18 a) H. Werner, M. Laubender, R. Wiedemann, and B. Windmüller, *Angew. Chem., Int. Ed. Engl.*, **35**, 1237 (1996). b) K. Ilg and H. Werner, *Chem.—Eur. J.*, **7**, 4633 (2001).
- 19 J.-D. van Loon, P. Seiler, and F. Diederich, *Angew. Chem., Int. Ed. Engl.*, **32**, 1187 (1993).
- 20 **4a**: ^1H NMR (C_6D_6 , Me_4Si) δ = -0.18 (s, 9H), 0.33 (s, 9H), 5.71 (dd, $^5J_{\text{H-H}}$ = 3.0, $^3J_{\text{Rh-H}}$ = 3.0 Hz, 1H), 6.02 (dd, $^5J_{\text{H-H}}$ = 3.0, $^3J_{\text{Rh-H}}$ = 3.0 Hz, 1H), 7.06 (m, 18H), 7.90 (m, 12H). ^{13}C NMR (C_6D_6 , Me_4Si) δ = -0.40, 0.40, 116.22, 122.16, 127.66 ($^3J_{\text{P-C}}$ = 4.5 Hz), 129.69, 131.85 ($^1J_{\text{P-C}}$ = 21.2 Hz), 135.21 ($^2J_{\text{P-C}}$ = 5.9 Hz), 154.66 (q, $^1J_{\text{Rh-C}}$ = 13.4, $^2J_{\text{P-C}}$ = 3.7 Hz), 157.07 (q, $^1J_{\text{Rh-C}}$ = 15.1, $^2J_{\text{P-C}}$ = 3.9 Hz). ^{31}P NMR (C_6D_6 , H_3PO_4) δ = 39.3 (d, $^1J_{\text{P-Rh}}$ = 134.3 Hz). ^{29}Si NMR (C_6D_6 , Me_4Si) δ = -4.9 (d, $^3J_{\text{Si-Rh}}$ = 3.7 Hz), -8.0 (d, $^3J_{\text{Si-Rh}}$ = 1.8 Hz). Anal. Calcd for $\text{C}_{46}\text{H}_{50}\text{ClP}_2\text{RhSi}_2$: C 64.29, H 5.86, Found: C 64.04, H 5.87. Yield 58% by ^1H NMR (17% isol.).
- 21 **4b**: Crystal data: $\text{C}_{52}\text{H}_{62}\text{ClP}_2\text{RhSi}_2$, fw=943.50, orthorhombic, space group = $Pna2(1)$, a = 33.305(4), b = 11.4630(12), c = 12.9333(13) Å, V = 4937.5(9) Å³, Z = 4, D_{calcd} = 1.269 g/cm³, R = 0.057, R_w = 0.125. Data were collected on Bruker APEX diffractometer with a CCD area detector, using graphite monochromated Mo $K\alpha$ radiation at 110 K. A total of 37042 reflections was measured. The structure was solved by direct methods and refined on F^2 by full-matrix least-squares techniques. The final cycle of least-squares refinement was based on 14067 observed reflections ($I \geq 2\sigma(I)$) with 535 variable parameters. The data were deposited in Cambridge Crystallographic Data Centre (CCDC 191256).
- 22 Typically, to a solution of $\text{RhCl}(\text{PPh}_3)_3$ (18.5 mg, 0.02 mmol) in diethyl ether (0.5 mL) were added **1a** (99 mg, 0.5 mmol) and diphenylsilane (184 mg, 1 mmol). The mixture was stirred at room temperature for 1 h. Usual workup gave **5a** and **6a** (Y. 35% and 16% isolated). Yields were determined by GC. **5a**: ^1H NMR (C_6D_6) δ = -0.01 (s, 9H), 0.06 (s, 9H), 1.37 (dd, J = 15, 2.8 Hz, 1H), 1.48 (dd, J = 15, 2.8 Hz, 1H), 4.5 (t, J = 2.8 Hz, 1H), 5.38 (s, 1H, Si-H), 7.1–7.2 (m, 6H), 7.65–7.69 (m, 4H). ^{13}C NMR (C_6D_6) δ = -0.77, -0.64, 17.57, 75.96, 78.15, 128.24, 129.98, 133.98, 135.99, 211.41. **6a**: ^1H NMR (C_6D_6) δ = 0.07 (s, 9H), 0.22 (s, 9H), 5.33 (s, 1H, Si-H), 6.05 (d, J = 19.0 Hz, 1H), 6.49 (d, J = 1.0 Hz, 1H), 7.16 (dd, 3J = 19.0, 4J = 1.0 Hz, 1H), 7.39 (m, 4H), 7.42 (m, 2H), 7.58 (dd, J = 7.8, 1.5 Hz, 4H). ^{13}C NMR (C_6D_6) δ = -1.32, 0.46, 128.32, 130.00, 134.23, 136.22, 136.25, 147.44, 153.73, 155.65.